

Decompositions of Edge-Colored Digraphs: A New Technique in the Construction of Constant-Weight Codes and Related Families

Yeow Meng Chee*, Fei Gao†, Han Mao Kiah§, Alan Chi Hung Ling‡, Hui Zhang*, Xiande Zhang*

*School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore,
emails: ymchee@ntu.edu.sg, huizhang@ntu.edu.sg, xiandezhang@ntu.edu.sg

†Institute of High Performance Computing, Agency for Science, Technology and Research, Singapore, email: gaofei@ihpc.a-star.edu.sg

§Coordinated Science Lab, University of Illinois, USA, email:hmkih@illinois.edu

‡Department of Computer Science, University of Vermont, USA, email: aling@emba.uvm.edu

Abstract—We demonstrate that certain Johnson-type bounds are asymptotically exact for a variety of classes of codes, namely, constant-composition codes, nonbinary constant-weight codes and multiply constant-weight codes. This was achieved via an interesting application of the theory of decomposition of edge-colored digraphs.

Key words and phrases: Johnson-type bounds, constant-composition codes, constant-weight codes, multiply constant-weight codes

I. INTRODUCTION

In 1970s, Wilson [1]–[4] demonstrated that the elementary necessary conditions for the existence of balanced incomplete block designs are asymptotically sufficient. His work therefore determined the size of an optimal binary constant-weight code of weight w and distance $2w - 2$, provided that the length n is sufficiently large and satisfies certain congruences. Moreover, these optimal codes meet the Johnson bound and his results show that Johnson bound is asymptotically exact for fixed weight w and distance $2w - 2$.

Since Wilson's seminal work, there have been interesting developments in both areas of combinatorial design and coding theory. In the former, Wilson's ideas matured into Lamken and Wilson's theory of decomposition of edge-colored digraphs [5] and the theory has been used extensively in establishing the asymptotic existence of many classes of combinatorial designs. Theoretical developments also extend Lamken and Wilson's results to other classes of decompositions. Of particular interest is the class of superpure decompositions [6].

On the other hand, binary constant-weight codes have been generalized to constant-composition codes and nonbinary constant-weight codes and have found applications in power-line communications [7], [8], frequency hopping [9], coding for bandwidth-limited channels [10]. More recently, multiply constant-weight codes are introduced in an application for physically unclonable functions [11]. Not surprisingly, in these generalizations, Johnson-type upper bounds have been derived.

Therefore, a natural question is whether the advanced techniques of decompositions of edge-colored digraphs are relevant in constructing optimal codes and whether Johnson-type bounds are asymptotically exact for these generalizations. We answer both questions in the affirmative for certain distances where the weight is fixed.

II. PRELIMINARY

The ring $\mathbb{Z}/q\mathbb{Z}$ is denoted by \mathbb{Z}_q . For positive integer n , the set $\{1, 2, \dots, n\}$ is denoted by $[n]$.

Let \mathbb{Z}_q^X denote the set of vectors whose elements belong to \mathbb{Z}_q and are indexed by X . A q -ary code of length n is then a set $\mathcal{C} \subseteq \mathbb{Z}_q^X$ with $|X| = n$. The support of a vector $u \in \mathbb{Z}_q^X$, denoted $\text{supp}(u)$, is the set $\{x \in X : u_x \neq 0\}$. The Hamming weight of $u \in \mathbb{Z}_q^X$ is defined as $\|u\| = |\text{supp}(u)|$. The distance induced by this norm is called the Hamming distance, so that $d(u, v) = \|u - v\|$, for $u, v \in \mathbb{Z}_q^X$. A code \mathcal{C} is said to have distance d if $d(u, v) \geq d$ for all distinct $u, v \in \mathcal{C}$. The composition of a vector $u \in \mathbb{Z}_q^X$ is the tuple $\bar{w} = [w_1, \dots, w_{q-1}]$, where $w_i = |\{x \in X : u_x = i\}|$, where $i \in \mathbb{Z}_q \setminus \{0\}$. Unless mentioned otherwise, we always assume $w_1 \geq w_2 \geq \dots \geq w_{q-1}$.

A. Constant-Weight Codes and Constant-Composition Codes

A code \mathcal{C} is said to have constant-weight w if every codeword in \mathcal{C} has weight w , and has constant-composition \bar{w} if every codeword in \mathcal{C} has composition \bar{w} . We refer to a q -ary code of length n , distance d , and constant-weight w as a $\text{CWC}(n, d, w)_q$. If in addition, the code has constant-composition \bar{w} , then it is referred to as a $\text{CCC}(n, d, \bar{w})_q$. The maximum size of a $\text{CWC}(n, d, w)_q$ is denoted $A_q(n, d, w)$ while the maximum size of a $\text{CCC}(n, d, \bar{w})_q$ is denoted $A_q(n, d, \bar{w})$. Any $\text{CWC}(n, d, w)_q$ or $\text{CCC}(n, d, \bar{w})_q$ attaining the maximum size is called *optimal*.

Consider a composition \bar{w} and let $w = \sum_{i=1}^{q-1} w_i$. Then a $\text{CCC}(n, d, \bar{w})_q$ is also a $\text{CWC}(n, d, w)_q$.

Johnson-type bounds for constant-weight codes and constant-composition codes have been derived.

Lemma 1 (Svanström [12], Svanström *et al.* [13]).

$$A_q(n, d, \bar{w}) \leq \left\lfloor \frac{n}{w_1} A_q(n-1, d, [w_1-1, \dots, w_{q-1}]) \right\rfloor,$$

$$A_q(n, d, w) \leq \left\lfloor \frac{(q-1)n}{w} A_q(n-1, d, w-1) \right\rfloor.$$

Apply the fact that $A_q(n, 2w, \bar{w}) = \lfloor n/w \rfloor$ and $A_q(n, 2w, w) = \lfloor n/w \rfloor$ (see Fu *et al.* [14], Chee *et al.* [15]) to Lemma 1, we have the following upper bounds:

$$A_q(n, 2w-2, \bar{w}) \leq \left\lfloor \frac{n}{w_1} \left\lfloor \frac{n-1}{w_1-1} \right\rfloor \right\rfloor, \quad (1)$$

$$A_q(n, 2w-3, \bar{w}) \leq \begin{cases} \left\lfloor \frac{n}{w_1} \left\lfloor \frac{n-1}{w_1-1} \right\rfloor \right\rfloor, & \text{if } w_1 > w_2, \\ \left\lfloor \frac{n}{w_1} \left\lfloor \frac{n-1}{w_1} \right\rfloor \right\rfloor, & \text{otherwise,} \end{cases} \quad (2)$$

$$A_q(n, 2w - 2, w) \leq \left\lfloor \frac{(q-1)n}{w} \left\lfloor \frac{n-1}{w-1} \right\rfloor \right\rfloor, \quad (3)$$

$$A_q(n, 2w - 3, w) \leq \left\lfloor \frac{(q-1)n}{w} \left\lfloor \frac{(q-1)(n-1)}{w-1} \right\rfloor \right\rfloor. \quad (4)$$

In this paper, we show that the above inequalities are exact provided that n is sufficiently large and satisfies certain congruence conditions. To do so, we apply the theory of decompositions of edge-colored digraphs to construct optimal codes meeting the above bounds.

We list some previous asymptotic or exact results.

- (i) Results for $A_q(n, d, \bar{w})$ are known
 - (a) for all \bar{w} and $d = 2w - 1$ [16];
 - (b) for all d where $w \leq 3$ [15]–[17];
 - (c) for $(q, d, w) = (3, 5, 4)$ [18].
- (ii) Results for $A_q(n, d, w)$ are known
 - (a) for all w and $d = 2w - 1$ [19];
 - (b) for all q and $(d, w) \in \{(3, 2), (4, 3), (5, 3)\}$ [19]–[21];
 - (c) for $(q, d, w) \in \{(3, 5, 4), (3, 6, 4), (4, 5, 4)\}$ [22]–[24].

B. Multiply Constant-Weight Codes

Consider a binary code $\mathcal{C} \subseteq \mathbb{Z}_2^{[m] \times [n]}$ of constant-weight mw and distance d . The code \mathcal{C} is said to be of *multiply constant-weight* w if for $u \in \mathcal{C}$, $i \in [m]$, the subword $(u_{i,j})_{j \in [n]}$ is of constant-weight w . Denote such a code by $\text{MCWC}(m, n, d, w)$. Similarly, the maximum size of a $\text{MCWC}(m, n, d, w)$ is given by $M(m, n, d, w)$ and a multiply constant-weight code attaining the maximum size is said to be *optimal*. A Johnson-type bound can again be derived.

Lemma 2 (Chee *et al.* [11]).

$$M(m, n, 2mw - 2, w) \leq \left\lfloor \frac{n}{w} \left\lfloor \frac{n}{w} \right\rfloor \right\rfloor. \quad (5)$$

Again, we verify (5) is exact provided n is sufficiently large and satisfies certain congruence conditions. Next, we describe the main tool in our constructions of optimal codes.

C. Decomposition of Edge-Colored Complete Digraphs

Denote the set of all ordered pairs of a finite set X with distinct components by $\overline{\binom{X}{2}}$. An *edge-colored digraph* is a triple $G = (V, C, E)$, where V is a finite set of *vertices*, C is a finite set of *colors* and E is a subset of $\overline{\binom{V}{2}} \times C$. Members of E are called *edges*. The *complete edge-colored digraph* on n vertices with r colors, denoted by $K_n^{(r)}$, is the edge-colored digraph (V, C, E) , where $|V| = n$, $|C| = r$ and $E = \overline{\binom{V}{2}} \times C$.

A family \mathcal{F} of edge-colored subgraphs of an edge-colored digraph K is a *decomposition* of K if every edge of K belongs to exactly one member of \mathcal{F} . Given a family of edge-colored digraphs \mathcal{G} , a decomposition \mathcal{F} of K is a \mathcal{G} -*decomposition* of K if each edge-colored digraph in \mathcal{F} is isomorphic to some $G \in \mathcal{G}$. Furthermore, a \mathcal{G} -decomposition of K is said to be *superpure* if any two distinct edge-colored subgraphs in \mathcal{F} share at most two vertices.

Lamken and Wilson [5] exhibited the asymptotic existence of decompositions of $K_n^{(r)}$ for a fixed family of digraphs. Hartmann [6] later extended their results to superpure decompositions. To state the theorems, we require more concepts.

Consider an edge-colored digraph $G = (V, C, E)$ with $|C| = r$. Let $((u, v), c) \in E$ denote a directed edge from u to v , colored by c . For any vertex u and color c , define the *indegree* and *outdegree* of u with respect to c , to be the number of directed edges of color c entering and leaving u respectively. Then for vertex u , we define the *degree vector* of u in G , denoted by $\tau(u, G)$, to be the vector of length $2r$, $\tau(u, G) \triangleq (\text{in}_1(u, G), \text{out}_1(u, G), \dots, \text{in}_r(u, G), \text{out}_r(u, G))$. Define $\alpha(\mathcal{G})$ to be the greatest common divisor of the integers t such that the $2r$ -vector (t, t, \dots, t) is a nonnegative integral linear combination of the degree vectors $\tau(u, G)$ as u ranges over all vertices of all digraphs $G \in \mathcal{G}$.

For each $G = (V, C, E) \in \mathcal{G}$, let $\mu(G)$ be the *edge vector* of length r given by $\mu(G) \triangleq (m_1(G), m_2(G), \dots, m_r(G))$ where $m_i(G)$ is the number of edges with color i in G . We denote by $\beta(\mathcal{G})$ the greatest common divisor of the integers m such that (m, m, \dots, m) is a nonnegative integral linear combination of the vectors $\mu(G)$, $G \in \mathcal{G}$. Then \mathcal{G} is said to be *admissible* if $(1, 1, \dots, 1)$ can be expressed as a positive rational combination of the vectors $\mu(G)$, $G \in \mathcal{G}$.

Below is the main theorem we allude to in our proofs.

Theorem 3 (Lamken and Wilson [5], Hartmann [6]). *Let \mathcal{G} be an admissible family of edge-colored digraphs with r colors. Then there exists a constant n_0 (resp. n_1) such that a (resp. superpure) \mathcal{G} -decomposition of $K_n^{(r)}$ exists for every $n \geq n_0$ (resp. $n \geq n_1$) satisfying: $n(n-1) \equiv 0 \pmod{\beta(\mathcal{G})}$ and $n-1 \equiv 0 \pmod{\alpha(\mathcal{G})}$.*

We apply Theorem 3 to construct codes that meet the upper bounds (1)–(5). To do so, we have two main steps.

- (A) Define a family \mathcal{G} of edge-colored digraphs on a set of r colors. We then show that a \mathcal{G} -decomposition of $K_n^{(r)}$ results in a code of length n satisfying certain weight and distance properties.
- (B) Compute $\alpha(\mathcal{G})$ and $\beta(\mathcal{G})$ and hence determine the congruences that n needs to satisfy.

In this paper, we focus on step (A), describing the construction of \mathcal{G} and establishing the correspondence to certain codes. As the computations in step (B) are usually tedious and analogous, we exhibit one computation for illustrative purposes and defer other computations to the full paper [25].

III. ASYMPTOTICALLY EXACT JOHNSON-TYPE BOUNDS FOR CONSTANT-COMPOSITION CODES

Fix $\bar{w} = [w_1, w_2, \dots, w_{q-1}]$ with $w_1 \geq \dots \geq w_{q-1} > 0$ and let $w = \sum_{i=1}^{q-1} w_i$. In this section, we construct infinite families of optimal $\text{CCC}(n, d, \bar{w})_q$ for $d = 2w - 2$ or $d = 2w - 3$, and establish the asymptotic exactness of (1) and (2).

A. When Distance $d = 2w - 2$

Consider the following characterization of codes of constant-weight w with distance $2w - 2$. Note that since the constant-composition codes are constant-weight codes, the lemma is applicable for both classes of codes.

Lemma 4. *The following are necessary and sufficient for a code \mathcal{C} of constant-weight w to have distance $2w - 2$:*

- (C1) *for $i \in [q-1]$, the ordered pairs in the set $\{(x, y) : u_x = i, y \in \text{supp}(u) \setminus \{x\}, \text{ for any } u \in \mathcal{C}\}$ are distinct; and*
- (C2) *for any $u, v \in \mathcal{C}$, $|\text{supp}(u) \cap \text{supp}(v)| \leq 2$.*

Note that condition (C2) ensures that the supports of any two codewords intersect in at most two positions. In the case when the supports of two codewords intersect at exactly two positions, condition (C1) ensures that both words differ in *at least one coordinate*. Hence, the distance between any two codewords is at least $2w - 2$.

Definition of the family $\mathcal{G}(\bar{w})$. For fixed \bar{w} , define an edge-colored digraph $G(\bar{w}) = (V[\bar{w}], C[\bar{w}], E[\bar{w}])$, where


$$V[\bar{w}] \triangleq \{x_{ij} : i \in [q-1], j \in [w_i]\};$$

$$C[\bar{w}] \triangleq [q-1];$$

$$E[\bar{w}] \triangleq \left\{ ((x_{ij}, x_{ij'}), i) : i \in [q-1], (j, j') \in \binom{[w_i]}{2} \right\} \cup \left\{ ((x_{ij}, x_{i'j'}), i) : (i, i') \in \binom{[q-1]}{2}, j \in [w_i], j' \in [w_{i'}] \right\}.$$

Let s be the largest integer such that $w_1 = w_2 = \dots = w_s$. For each $s+1 \leq i \leq q-1$, let G_i be an edge-colored digraph with two vertices y_i, z_i and one directed edge with color i from y_i to z_i . Then $\mathcal{G}(\bar{w}) = \{G(\bar{w})\} \cup \{G_i : s+1 \leq i \leq q-1\}$.

Example 1. Let $q = 3$ and $\bar{w} = [w_1, w_2] = [3, 2]$. The edge-colored digraph $G(\bar{w})$ is given below, where the solid lines and dotted lines denote directed edges with color 1 and 2, respectively, and “ \longleftrightarrow ” (“ \dashrightarrow ”) denotes the two directed edges with the same color with one in each direction.




Then G_2 is the digraph $y_2 \dashrightarrow z_2$ and the family of digraphs is given by $\mathcal{G}(\bar{w}) = \{G(\bar{w}), G_2\}$.

Construction of an optimal $\text{CCC}(n, 2w-2, \bar{w})_q$. Suppose a superpure $\mathcal{G}(\bar{w})$ -decomposition of $K_n^{(q-1)}$ exists. For each F isomorphic to $G(\bar{w})$, there is a unique partition of the vertex set $V(F) = \bigcup_{i=1}^{q-1} S_i$ so that all outgoing edges from x in F has color i if and only if $x \in S_i$. Then construct one codeword u with support $V(F)$ such that $u_x = i$ for $x \in S_i$. Hence, u has composition \bar{w} .¹

Since we have a $\mathcal{G}(\bar{w})$ -decomposition of $K_n^{(q-1)}$, then (C1) of Lemma 4 is satisfied. Furthermore, since the decomposition is superpure, (C2) is also met. Hence, we obtain a $\text{CCC}(n, 2w-2, \bar{w})_q$ code of size $n(n-1)/(w_1(w-1))$. The code meets the upper bound in (1) and is therefore optimal.

Example 2. Let $\bar{w} = [2, 1]$. Consider the following superpure $\mathcal{G}(\bar{w})$ -decomposition of $K_5^{(2)}$:

¹While we only use the $G(\bar{w})$'s in the construction of the codewords, the G_i 's are needed to ensure that every edge of $K_n^{(q-1)}$ appears in the $\mathcal{G}(\bar{w})$ -decomposition.



$$\begin{array}{ccccccccc} 1 & \dashrightarrow & 2 & & 2 & \dashrightarrow & 1 & & 3 & \dashrightarrow & 1 & & 4 & \dashrightarrow & 2 & & 5 & \dashrightarrow & 3 \\ 1 & \dashrightarrow & 3 & & 2 & \dashrightarrow & 4 & & 3 & \dashrightarrow & 5 & & 4 & \dashrightarrow & 5 & & 5 & \dashrightarrow & 4 \end{array}$$

The corresponding code is then given by $\{(0, 1, 2, 1, 0), (0, 2, 1, 0, 1), (1, 0, 1, 2, 0), (1, 1, 0, 0, 2), (2, 0, 0, 1, 1)\}$, which is indeed a $\text{CCC}(5, 4, [2, 1])_3$.

Computation of $\alpha(\mathcal{G}(\bar{w}))$ and $\beta(\mathcal{G}(\bar{w}))$. First consider the digraph $G(\bar{w})$. Observe that for $i \in [q-1], k \in [w_i]$ we have $\text{in}_i(x_{ik}, G(\bar{w})) = w_i - 1$, $\text{out}_i(x_{ik}, G(\bar{w})) = w - 1$, $\text{in}_j(x_{ik}, G(\bar{w})) = w_j$ and $\text{out}_j(x_{ik}, G(\bar{w})) = 0$ for $j \neq i$. Consider G_i for $s+1 \leq i \leq q$. Then $\text{in}_i(z_i, G_i) = \text{out}_i(y_i, G_i) = 1$ and all other indegrees and outdegrees are zero.

Let $a = \text{gcd}(w_1, w)$. Pick $t = \lfloor w/w_1 \rfloor$ so that $0 \leq w - tw_1 < w_1$. Observe also that $t \geq s$. Consider the vector

$$v = \frac{w - tw_1}{a} \tau(x_{(t+1)1}, G(\bar{w})) + \frac{w_1}{a} \sum_{i=1}^t \tau(x_{i1}, G(\bar{w})).$$

For $j \in [q-1]$, let $\text{in}_j(v)$ and $\text{out}_j(v)$ denote the coordinates in v corresponding to the summation of the indegrees or outdegrees with respect to color j . Then we have

$$\text{in}_j(v) = \begin{cases} \frac{w w_j - w_1}{a} = \frac{w_1(w-1)}{a}, & \text{for } j \in [s], \\ \frac{w w_j - w_1}{a} < \frac{w_1(w-1)}{a}, & \text{for } s+1 \leq j \leq t, \\ \frac{w w_{t+1} - w + t w_1}{a} < \frac{w_{t+1}(w-1)}{a}, & \text{for } j = t+1, \\ \frac{w w_j}{a}, & \text{otherwise,} \end{cases}$$

$$\text{out}_j(v) = \begin{cases} \frac{w_1(w-1)}{a}, & \text{for } j \in [t], \\ \frac{(w - t w_1)(w-1)}{a} < \frac{w_1(w-1)}{a}, & \text{for } j = t+1, \\ 0, & \text{otherwise.} \end{cases}$$

Observe that the first $2s$ coordinates of v are $w_1(w-1)/a$ and all other coordinates have values at most $w_1(w-1)/a$. Adding to v a suitable nonnegative integral combinations of $\tau(y_i, G_i)$'s and $\tau(z_i, G_i)$'s, we get $w_1(w-1)/a \cdot (1, 1, \dots, 1)$. Hence, we conclude that $\alpha(\mathcal{G}(\bar{w})) = w_1(w-1)/a$.

Next, consider the edge vector $\mu(G)$ with $G \in \mathcal{G}(\bar{w})$. For $G(\bar{w})$, $m_i(G(\bar{w})) = w_i(w-1)$ for $i \in [q-1]$, while for G_i with $s+1 \leq i \leq q-1$, $m_i(G_i) = 1$ and $m_j(G_i) = 0$ for $j \neq i$. Hence, $\beta(\mathcal{G}(\bar{w})) = w_1(w-1)$.

Applying Theorem 3, we obtain our first asymptotic result.

Proposition 5. Fix \bar{w} and let $w = \sum_{i=1}^{q-1} w_i$. There exists an integer n_0 such that

$$A_q(n, 2w-2, \bar{w}) = \frac{n(n-1)}{w_1(w-1)}$$

for all $n \geq n_0$ satisfying $n(n-1) \equiv 0 \pmod{w_1(w-1)}$ and $n-1 \equiv 0 \pmod{w_1(w-1)/a}$, where $a = \text{gcd}(w_1, w)$.

B. When Distance $d = 2w - 3$

We have the following analogous characterization of codes of constant-weight w with distance $2w - 3$.

Lemma 6. The following are sufficient for a code \mathcal{C} of weight w to have distance $2w - 3$:

- (C3) for $i, j \in [q-1]$, the ordered pairs in the set $\{(x, y) : u_x = i, u_y = j, x \neq y, \text{ for any } u \in \mathcal{C}\}$ are distinct; and
 (C4) for any $u, v \in \mathcal{C}$, $|\text{supp}(u) \cap \text{supp}(v)| \leq 2$.

Again observe that condition (C4) ensures that the supports of any two codewords intersect in at most two positions. In the case when the supports of two codewords intersect exactly at two positions, condition (C3) ensures that the codewords differ at *both coordinates*. Hence, the distance between any two codewords is at least $2w-3$.


Definition of $\mathcal{G}^*(\bar{w})$. For fixed \bar{w} , define an edge-colored digraph $G^*(\bar{w}) = (V(\bar{w}), C(\bar{w}), E(\bar{w}))$, where

$$\begin{aligned} V(\bar{w}) &\triangleq \{x_{ij} : i \in [q-1], j \in [w_i]\}; \\ C(\bar{w}) &\triangleq [q-1] \times [q-1]; \\ E(\bar{w}) &\triangleq \left\{ ((x_{ij}, x_{ij'}), (i, i')) : i \in [q-1], (j, j') \in \binom{[w_i]}{2} \right\} \cup \\ &\quad \left\{ ((x_{ij}, x_{i'j'}), (i, i')) : (i, i') \in \binom{[q-1]}{2}, j \in [w_i], j' \in [w_{i'}] \right\}. \end{aligned}$$

For $i, j \in [q-1]$, let G_{ij}^* be a digraph with vertices y_i, z_j and one directed edge with color (i, j) from y_i to z_j . To define $\mathcal{G}^*(\bar{w})$, we have two cases depending on whether $w_1 = w_2$:

- (i) When $w_1 > w_2$, let r be the largest integer such that $w_2 = \dots = w_r = w_1 - 1$. Then set $\mathcal{G}^*(\bar{w}) = \{G^*(\bar{w})\} \cup \{G_{ij}^* : (i, j) \in [q-1] \times [q-1] \setminus \{(1, 1), (1, 2), (1, 3), \dots, (1, r), (2, 1), (3, 1), \dots, (r, 1)\}\}$.
- (ii) When $w_1 = w_2$, let r be the largest integer such that $w_1 = \dots = w_r$. Then set $\mathcal{G}^*(\bar{w}) = \{G^*(\bar{w})\} \cup \{G_{ij}^* : (i, j) \in [q-1] \times [q-1] \setminus \binom{[r]}{2}\}$.

Example 3. Let $\bar{w} = [3, 2]$. The edge-colored digraph $G^*(\bar{w})$ is given below, where the lines “ \longrightarrow ”, “ \dashrightarrow ”, “ $\cdots \dashrightarrow$ ” and “ \rightsquigarrow ” denote directed edges with color $(1, 1)$, $(1, 2)$, $(2, 1)$ and $(2, 2)$ respectively.



Then G_{22}^* is the digraph $y_2 \rightsquigarrow z_2$ and the family of digraphs is given by $\mathcal{G}^*(\bar{w}) = \{G^*(\bar{w}), G_{22}^*\}$.

Construction of an optimal CCC($n, 2w-3, \bar{w}$)_q. Suppose a superpure $\mathcal{G}^*(\bar{w})$ -decomposition of $K_n^{\binom{(q-1)^2}$ exists. For F isomorphic to $G^*(\bar{w})$, there is a unique partition $V(F) = \bigcup_{i=1}^{q-1} S_i$ so that the edges from x to y in F has color (i, j) if $x \in S_i$ and $y \in S_j$. Construct a codeword u with support $V(F)$ such that $u_x = i$ for $x \in S_i$. So, u has composition \bar{w} .

Since we have a superpure $\mathcal{G}^*(\bar{w})$ -decomposition of $K_n^{\binom{(q-1)^2}$, (C3) and (C4) of Lemma 6 are satisfied. Hence, we obtain a CCC($n, 2w-3, \bar{w}$)_q code. Let M be the number of digraphs isomorphic to $G^*(\bar{w})$. It is easy to see that $M = n(n-1)/(w_1(w_1-1))$ if $w_1 \neq w_2$ and $M = n(n-1)/w_1^2$, otherwise. The code is optimal by the upper bound (2).

We compute $\alpha(\mathcal{G}^*(\bar{w}))$ and $\beta(\mathcal{G}^*(\bar{w}))$ in the full paper.

$$\begin{aligned} \alpha(\mathcal{G}^*(\bar{w})) &= \begin{cases} w_1(w_1-1), & \text{if } w_1 > w_2, \\ w_1, & \text{otherwise.} \end{cases} \\ \beta(\mathcal{G}^*(\bar{w})) &= \begin{cases} w_1(w_1-1), & \text{if } w_1 > w_2, \\ w_1^2, & \text{otherwise.} \end{cases} \end{aligned}$$

Applying Theorem 3 we obtain the following proposition.

Proposition 7. Fix \bar{w} and let $w = \sum_{i=1}^{q-1} w_i$. There exists an integer n_0 such that

$$A_q(n, 2w-3, \bar{w}) = \begin{cases} \frac{n(n-1)}{w_1(w_1-1)}, & \text{if } w_1 > w_2, \\ \frac{n(n-1)}{w_1^2}, & \text{otherwise,} \end{cases}$$

for all $n \geq n_0$ satisfying

- (i) $n-1 \equiv 0 \pmod{w_1(w_1-1)}$, if $w_1 > w_2$,
- (ii) $n-1 \equiv 0 \pmod{w_1^2}$, otherwise.

IV. ASYMPTOTICALLY EXACT JOHNSON-TYPE BOUNDS FOR q -ARY CONSTANT-WEIGHT CODES

In this section, we construct infinite families of optimal CWC(n, d, w)_q codes for $d = 2w-2$ or $d = 2w-3$, and establish the asymptotic exactness of (3) and (4).

Note that constant-composition codes are special instances of constant-weight codes. Hence, we make use of the digraphs defined in Section III to form the desired families for constant-weight codes. Specifically, consider the set of compositions,

$$W = \left\{ [w_1, w_2, \dots, w_{q-1}] : 0 \leq w_i \leq w \text{ for } i \in [q-1], \sum_{i=1}^{q-1} w_i = w \right\}.$$

Note here we do not require all values in the composition to be positive and the composition to be monotone decreasing. Furthermore, we can extend the constructions in Section III to define $G(\bar{w})$ and $G^*(\bar{w})$ for all $\bar{w} \in W$. Define the following families of digraphs

$$\mathcal{G}(w) \triangleq \bigcup_{\bar{w} \in W} G(\bar{w}) \text{ and } \mathcal{G}^*(w) \triangleq \bigcup_{\bar{w} \in W} G^*(\bar{w}).$$

Similar to constructions in Section III, we have that a superpure $\mathcal{G}(w)$ -decomposition of $K_n^{\binom{(q-1)}{2}}$ and a superpure $\mathcal{G}^*(w)$ -decomposition of $K_n^{\binom{(q-1)^2}$ yield a CWC($n, 2w-2, w$)_q and a CWC($n, 2w-3, w$)_q respectively.

The number of digraphs in a superpure $\mathcal{G}(w)$ -decomposition of $K_n^{\binom{(q-1)}{2}}$ is $(q-1)n(n-1)/(w(w-1))$ since the total number of directed edges is $(q-1)n(n-1)$. In the full paper we used the methods of Lamken and Wilson to establish the following.

Proposition 8. Fix w . There exists an integer n_0 such that

$$A_q(n, 2w-2, w) = \frac{(q-1)n(n-1)}{w(w-1)}$$

for all $n \geq n_0$ satisfying $(q-1)n(n-1) \equiv 0 \pmod{w(w-1)}$ and $n-1 \equiv 0 \pmod{w-1}$.

On the other hand, $\mathcal{G}^*(w)$ corresponds to the family of digraphs constructed in the proof of [5, Theorem 8.1]. Therefore, we have the following proposition.

Proposition 9. Fix w . There exists an integer n_0 such that

$$A_q(n, 2w - 3, w) = \frac{(q - 1)^2 n(n - 1)}{w(w - 1)}$$

for all $n \geq n_0$ satisfying $(q - 1)^2 n(n - 1) \equiv 0 \pmod{w(w - 1)}$ and $(q - 1)(n - 1) \equiv 0 \pmod{w - 1}$.

V. ASYMPTOTICALLY EXACT JOHNSON-TYPE BOUNDS FOR MULTIPLY CONSTANT-WEIGHT CODES

In this section, we construct an infinite family of optimal MCWC($m, n, 2mw - 2, w$) and establish the asymptotic exactness of (5). Fix m and w . Using digraphs from Section III, we define the edge-colored digraph $H^*(m, w) \triangleq G^*(\bar{w})$, with $w_1 = w_2 = \dots = w_m = w$ (here $q - 1 = m$). Define $\mathcal{H}^*(m, w) = \{H^*(m, w)\} \cup \{G_{i_i}^* : i \in [m]\}$.

Construction of an optimal MCWC($m, n, 2mw - 2, w$). Suppose an $\mathcal{H}^*(m, w)$ -decomposition of $K_n^{(m^2)}$ exists. For each F isomorphic to $H^*(m, w)$, there is a unique partition $V(F) = \bigcup_{i=1}^m S_i$ so that the edges from x to y in F has color (i, j) if and only if $x \in S_i$ and $y \in S_j$. Then construct one codeword u such that $u_{(i,x)} = 1$ for $i \in [m]$ and $x \in S_i$.

Since we have an $\mathcal{H}^*(m, w)$ -decomposition of $K_n^{(m^2)}$, the distance of the code is $2mw - 2$. Hence, we have an MCWC($m, n, 2mw - 2, w$) of size $n(n - 1)/w^2$ that meets the upper bound given by (5). Similar computations yield $\alpha(\mathcal{H}^*(m, w)) = w$ and $\beta(\mathcal{H}^*(m, w)) = w^2$. Applying Theorem 3 we obtain the following.

Proposition 10. Fix m and w . There exists an integer n_0 so that

$$M(m, n, 2mw - 2, w) = \frac{n(n - 1)}{w^2}$$

for all $n \geq n_0$ satisfying $n - 1 \equiv 0 \pmod{w^2}$.

VI. CONCLUSION

We verify that Johnson-type bounds are asymptotically exact for several generalizations of binary constant-weight codes. This was achieved via an interesting application of superpure decompositions of edge-colored digraphs.

Observe that in Propositions 5, 7, 8, 9 and 10 Johnson-type bounds are shown to be exact for sufficiently large n satisfying certain congruences. We hypothesize that the Johnson-type bounds are tight up to an additive constant for sufficiently large n .

Conjecture. Fix q, \bar{w}, w, m and let $d \in \{2w - 2, 2w - 3\}$. Define $U_q(n, d, \bar{w}), U_q(n, d, w), U(m, n, d, w)$ to be the upper bounds given by (1) – (5). Then

$$A_q(n, d, \bar{w}) = U_q(n, d, \bar{w}) - O(1),$$

$$A_q(n, d, w) = U_q(n, d, w) - O(1),$$

$$M(m, n, 2mw - 2, w) = U(m, n, 2mw - 2, w) - O(1).$$

The case for binary constant-weight codes, that is, $q = 2$ and $d = 2w - 2$, has been verified by Chee *et al.* [26].

ACKNOWLEDGEMENT

This work was done while H. M. Kiah was a graduate student at Nanyang Technological University.

The authors thank the anonymous reviewers and the TPC member for their constructive comments.

REFERENCES

- [1] R. M. Wilson, "An existence theory for pairwise balanced designs. I. Composition theorems and morphisms," *J. Combin. Theory Ser. A*, vol. 13, pp. 220–245, 1972.
- [2] —, "An existence theory for pairwise balanced designs. II. The structure of PBD-closed sets and the existence conjectures," *J. Combin. Theory Ser. A*, vol. 13, pp. 246–273, 1972.
- [3] —, "Cyclotomy and difference families in elementary abelian groups," *J. Number Theory*, vol. 4, pp. 17–47, 1972.
- [4] —, "An existence theory for pairwise balanced designs. III. Proof of the existence conjectures," *J. Combin. Theory Ser. A*, vol. 18, pp. 71–79, 1975.
- [5] E. R. Lamken and R. M. Wilson, "Decompositions of edge-colored complete graphs," *J. Combin. Theory Ser. A*, vol. 89, no. 2, pp. 149–200, 2000.
- [6] S. Hartmann, "Superpure digraph designs," *J. Combin. Des.*, vol. 10, no. 4, pp. 239–255, 2002.
- [7] W. Chu, C. J. Colbourn, and P. Dukes, "Constructions for permutation codes in powerline communications," *Des. Codes Cryptogr.*, vol. 32, no. 1–3, pp. 51–64, 2004.
- [8] Y. M. Chee, H. M. Kiah, and P. Purkayastha, "Estimates on the size of symbol weight codes," *IEEE Trans. Inform. Theory*, vol. 59, no. 1, pp. 301–314, January 2013.
- [9] W. Chu, C. J. Colbourn, and P. Dukes, "On constant composition codes," *Discrete Appl. Math.*, vol. 154, no. 6, pp. 912–929, 2006.
- [10] D. J. Costello and G. D. Forney, "Channel coding: the road to channel capacity," *Proc. IEEE*, vol. 95, no. 6, pp. 1150–1177, 2007.
- [11] Y. M. Chee, Z. Cherif, J.-L. Danger, S. Guillely, H. M. Kiah, J.-L. Kim, P. Solé, and X. Zhang, "Multiply constant weight codes and the reliability of loop physically unclonable functions," 2013, preprint.
- [12] M. Svanström, "Ternary codes with weight constraints," Ph.D. dissertation, Linköpings Universitet, Sweden, 1999.
- [13] M. Svanström, P. R. J. Östergård, and G. T. Bogdanova, "Bounds and constructions for ternary constant-composition codes," *IEEE Trans. Inform. Theory*, vol. 48, no. 1, pp. 101–111, 2002.
- [14] F.-W. Fu, A. J. H. Vinck, and S.-Y. Shen, "On the constructions of constant-weight codes," *IEEE Trans. Inform. Theory*, vol. 44, no. 1, pp. 328–333, 1998.
- [15] Y. M. Chee, G. Ge, and A. C. H. Ling, "Group divisible codes and their application in the construction of optimal constant-composition codes of weight three," *IEEE Trans. Inform. Theory*, vol. 54, no. 8, pp. 3552–3564, 2008.
- [16] Y. M. Chee, S. H. Dau, A. C. H. Ling, and S. Ling, "Linear size optimal q -ary constant-weight codes and constant-composition codes," *IEEE Trans. Inform. Theory*, vol. 56, no. 1, pp. 140–151, 2010.
- [17] M. Svanström, "Constructions of ternary constant-composition codes with weight three," *IEEE Trans. Inform. Theory*, vol. 46, no. 7, pp. 2644–2647, 2000.
- [18] F. Gao and G. Ge, "Optimal ternary constant-composition codes of weight four and distance five," *IEEE Trans. Inform. Theory*, vol. 57, no. 6, pp. 3742–3757, 2011.
- [19] Y. M. Chee and S. Ling, "Constructions for q -ary constant-weight codes," *IEEE Trans. Inform. Theory*, vol. 53, no. 1, pp. 135–146, 2007.
- [20] Y. M. Chee, S. H. Dau, A. C. H. Ling, and S. Ling, "The sizes of optimal q -ary codes of weight three and distance four: a complete solution," *IEEE Trans. Inform. Theory*, vol. 54, no. 3, pp. 1291–1295, 2008.
- [21] Y. M. Chee, G. Ge, H. Zhang, and X. Zhang, "Hanani triple packings and optimal q -ary codes of constant weight three," *Des. Codes Cryptogr.*, to appear.
- [22] H. Zhang and G. Ge, "Optimal ternary constant-weight codes of weight four and distance six," *IEEE Trans. Inform. Theory*, vol. 56, no. 5, pp. 2188–2203, 2010.
- [23] H. Zhang, X. Zhang, and G. Ge, "Optimal ternary constant-weight codes with weight 4 and distance 5," *IEEE Trans. Inform. Theory*, vol. 58, no. 5, pp. 2706–2718, 2012.
- [24] H. Zhang and G. Ge, "Optimal quaternary constant-weight codes with weight four and distance five," *IEEE Trans. Inform. Theory*, vol. 59, no. 3, pp. 1617–1629, 2013.
- [25] Y. M. Chee, F. Gao, H. M. Kiah, A. C. H. Ling, H. Zhang, and X. Zhang, "Decompositions of edge-colored digraphs: A new technique in the construction of constant-weight codes and related families," 2014, preprint.
- [26] Y. M. Chee, C. J. Colbourn, A. C. H. Ling, and R. M. Wilson, "Covering and packing for pairs," *J. Comb. Theory Ser. A*, vol. 120, no. 7, pp. 1440–1449, 2013.