

Combinatorics, 2016 Fall, USTC
Homework 8

- The due is on Tuesday, Nov. 22, at beginning of the class.
- Solve all problems.

1. Find the maximum number of line segments that a Hasse diagram of poset (X, \prec) with $|X| = n$ can have. Then define a poset which achieves this maximum number and draw its Hasse diagram.

2. For two natural numbers a, b , the symbol $a|b$ means that “ a divides b ”. In other words, there exists a natural number c such that $b = ac$. First verify that the relation “ $|$ ” is a partial ordering on the set \mathbf{N} of natural numbers. Then prove that every finite poset can be embedded into $(\mathbf{N}, |)$.

3. For any point $p \in R^d$ in d -dimension, write $p = (p_1, p_2, \dots, p_d)$. A set \mathcal{P} of points in R^d is called *good*, if for each $i \in [d]$, the i^{th} coordinates of these points are distinct. Given two points $p, q \in R^d$, define $\text{box}(p, q) := \{x \in R^d : \min\{p_i, q_i\} \leq x_i \leq \max\{p_i, q_i\} \text{ for each } i\}$ as the box determined by points p, q .

Prove that in any good set \mathcal{P} of $2^{2^{d-1}} + 1$ points of R^d , there is a point $x \in \mathcal{P}$ which is in the box determined by two of the other points in \mathcal{P} .

4. For any integers $k, l \geq 1$, construct a sequence of kl distinct integers with no increasing subsequence of length $k + 1$ and with no decreasing subsequence of length $l + 1$.

5. Let $P = (X, \prec)$ be a finite partial ordered set. Show that X can be expressed as a disjoint union of at most $\alpha(P)$ chains.

Hint: by induction on $|X|$.

6. Construct an explicit 2-edge-coloring of K_{kl} to show that $R(k + 1, l + 1) \geq kl + 1$.

7. Use Ramsey’s theorem to prove: for every integer $k \geq 2$, there is an integer n such that every sequence of n distinct real numbers contains a monotone subsequence of k real numbers.

(You must use Ramsey’s theorem and cannot use the Erdős-Szekeres theorem.)

8. For integers $k \geq 2$ and $s_1, s_2, \dots, s_k \geq 2$, the *Ramsey number* $R_k(s_1, s_2, \dots, s_k)$ is the least number of integer n such that any k -edge-coloring of K_n has a monochromatic clique K_{s_i} in color i . Prove that for any integers $s_1, s_2, \dots, s_k \geq 2$, the Ramsey Number

$$R_k(s_1, s_2, \dots, s_k) < +\infty.$$

9. Prove that $2^k \leq R_k(3, 3, \dots, 3) \leq (k + 1)!$