

Combinatorics, 2016 Fall, USTC
Homework 9

- The due is on **Tuesday**, Nov. 29, at beginning of the class.

1. Let V be a finite set and $r \geq 3$ be an integer. An r -uniform hypergraph is a pair (V, E) , where $E \subseteq \binom{V}{r}$. And $K_n^{(r)}$ denotes the complete r -uniform hypergraph on n vertices, i.e., $K_n^{(r)} = (V, \binom{V}{r})$ for $|V| = n$. Let hypergraph Ramsey number $R^{(r)}(s, t)$ be the least integer n such that any 2-edge-coloring of $K_n^{(r)}$ has a blue $K_s^{(r)}$ or a red $K_t^{(r)}$.

Prove that for any $s, t \geq r$, $R^{(r)}(s, t) < \infty$.

2. (a). A fixed point of a permutation π is an integer i such that $\pi(i) = i$. Let X_π be the number of fixed points of permutation π . What is the expectation $E[X_\pi]$ for a random permutation π chosen from the set S_n ? (Recall that S_n is the set of all permutations of $[n]$.)

(b). We toss a fair coin n times. What is the expected number of “runs”? “Runs” are consecutive tosses with the same result. For example, the sequence “TTTHHHHTT” has 3 runs, where ‘T’ means ‘tail’ and ‘H’ means ‘Head’.

3. Let $n \geq 6$. We color the edges of complete graph K_n with colors blue and red in such a way that each edge is in at most one blue triangle. Let X be a maximal subset of the vertices with the property that X does not contain a blue triangle. Then

(a) show that X contains at least $|X|/2$ red edges, and

(b) show that $|X| \geq \sqrt{2n}$.

4. Let $m \geq n^2/4$. Show that any graph with n vertices and with m edges has at least

$$\frac{1}{3} \left(\frac{4m^2}{n} - mn \right)$$

triangles.