

On problems about judicious bipartitions of graphs

Yuliang Ji* Jie Ma† Juan Yan‡ Xingxing Yu§

Abstract: Bollobás and Scott [5] conjectured that every graph G has a balanced bipartite spanning subgraph H such that for each $v \in V(G)$, $d_H(v) \geq (d_G(v) - 1)/2$. In this paper, we show that every graphic sequence has a realization for which this Bollobás-Scott conjecture holds, confirming a conjecture of Hartke and Seacrest [10]. On the other hand, we give an infinite family of counterexamples to this Bollobás-Scott conjecture, which indicates that $\lfloor (d_G(v) - 1)/2 \rfloor$ (rather than $(d_G(v) - 1)/2$) is probably the correct lower bound. We also study bipartitions V_1, V_2 of graphs with a fixed number of edges. We provide a (best possible) upper bound on $e(V_1)^\lambda + e(V_2)^\lambda$ for any real $\lambda \geq 1$ (the case $\lambda = 2$ is a question of Scott [13]) and answer a question of Scott [13] on $\max\{e(V_1), e(V_2)\}$.

Keywords: bipartition; bisection; degree sequence; complete k -partite graph; ℓ_λ -norm

AMS classification: 05C07, 05C70

*School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, China. Email: jjyl@mail.ustc.edu.cn.

†School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, China. Email: jiema@ustc.edu.cn. Partially supported by NSFC projects 11501539 and 11622110.

‡College of Mathematics and Systems Science, Xinjiang University, Urumqi, Xinjiang 830046, China. Email: yanjuan207@163.com. Partially supported by NSFC project 11501486.

§School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA. Email: yu@math.gatech.edu. Partially supported by NSF grants DMS-1265564 and DMS-1600387.

1 Introduction

For any positive integer k , let $[k] := \{1, \dots, k\}$. Let G be a graph and V_1, \dots, V_k be a partition of $V(G)$. When $k = 2$, such a partition is said to be a *bipartition* of G . A subgraph H of a graph G is said to be a *bisection* of G if H is a bipartite spanning subgraph of G and the partition sets of H differ in size by at most one. For $i, j \in [k]$, we use $e(V_i)$ to denote the number of edges of G with both ends in V_i and use $e(V_i, V_j)$ to denote the number of edges between V_i and V_j . *Judicious partitioning problems* for graphs ask for partitions of graphs that bound a number of quantities simultaneously, such as all $e(V_i)$ and $e(V_i, V_j)$. There has been extensive research on this type of problems over the past two decades.

As an attempt to better understand how edges of a graph are distributed, we study several judicious bipartitioning problems. Specifically, we study a conjecture of Bollobás and Scott [5] and its degree sequence version conjectured by Hartke and Seacrest [10]. We also study two questions of Scott [13] on bipartitions V_1, V_2 of a graph with m edges, bounding $e(V_1)^2 + e(V_2)^2$ and $\max\{e(V_1), e(V_2)\}$ in terms of m .

For a graph G and for any $v \in V(G)$, we use $d_G(v)$ to denote the degree of the vertex v in G . It is well known that if H is a maximum bipartite spanning subgraph of a graph G , then $d_H(v) \geq d_G(v)/2$ for each $v \in V(G)$. This, however, may not be true if one requires H to be a bisection, as observed by Bollobás and Scott [5] by considering the complete bipartite graphs $K_{2\ell+1, m}$ for $m \geq 2\ell + 3$. In an attempt to obtain a similar result for bisections, Bollobás and Scott [5] conjectured that every graph G has a bisection H such that

$$d_H(v) \geq (d_G(v) - 1)/2 \quad \text{for all } v \in V(G). \quad (1)$$

This conjecture for regular graphs was made by Häggkvist [8] in 1978, and variations of this problem were studied by Ban and Linial [2].

Hartke and Seacrest [10] studied a degree sequence version of this Bollobás-Scott conjecture. A nondecreasing sequence π (of nonnegative integers) is said to be *graphic* if it is the degree sequence of some finite simple graph G ; and such G is called a *realization* of the sequence π . Hartke and Seacrest [10] proved that for any graphic sequence π with even length, π has a realization G which admits a bisection H such that for all $v \in V(G)$, $d_H(v) \geq \lfloor (d_G(v) - 1)/2 \rfloor$. They further conjectured that for any graphic sequence π with even length, π has a realization G for which (1) holds. We prove this Hartke-Seacrest conjecture for all graphic sequences.

For a graph G and a labeling of its vertices $V(G) = \{v_1, \dots, v_n\}$, we define the *parity bisection* of G to be the bisection with partition sets V_1 and V_2 , where $V_i = \{v_j \in V(G) : j \equiv i \pmod{2}\}$ for each $i \in [2]$, and $E(H) = \{uv \in E(G) : u \in V_1 \text{ and } v \in V_2\}$.

Theorem 1.1. *Let $\pi = (d_1, \dots, d_n)$ be any graphic sequence with $d_1 \geq \dots \geq d_n$. Then there exists a realization G of π with $V(G) = \{v_1, \dots, v_n\}$ and $d_G(v_i) = d_i$ for $i \in [n]$, such that if H denotes the parity bisection of G then $d_H(v_i) \geq (d_G(v_i) - 1)/2$ for $i \in [n]$.*

The bound in Theorem 1.1 is best possible, as shown by the following example given by Hartke and Seacrest [10]. Let G be the join of a clique K on k vertices and an independent set I on $n - k$ vertices, where n is even and $k < n/2$ is odd. It is not hard to show that G in fact is the unique realization of the sequence $\pi = (d_1, \dots, d_n)$ with $d_1 = \dots = d_k = n - 1$ and $d_{k+1} = \dots = d_n = k$. Let H be an arbitrary bisection of G with parts A, B and,

without loss of generality, assume that $|A \cap V(K)| \leq k/2$. Since $k < n/2$, there must exist a vertex $v \in B \cap I$. So $d_H(v) = |A \cap V(K)| \leq \lfloor k/2 \rfloor$. Since $d_G(v) = k$ and k is odd, we see that $d_H(v) \leq (d_G(v) - 1)/2$.

The second result in this paper gives indication that perhaps the lower bound in the original Bollobás-Scott conjecture was meant to be $\lfloor (d_G(v) - 1)/2 \rfloor$ (rather than $(d_G(v) - 1)/2$).

Proposition 1.2. *Let r_1, r_2, r_3 be pairwise distinct odd integers such that for every $i \in [3]$, $r_i \notin \{1, \lfloor (r_1 + r_2 + r_3)/2 \rfloor, \lceil (r_1 + r_2 + r_3)/2 \rceil\}$. Then for any bisection H of the complete 3-partite graph $G := K_{r_1, r_2, r_3}$, there always exists a vertex v with $d_H(v) < (d_G(v) - 1)/2$.*

This result will follow from a more general result, Proposition 3.3, on all complete multipartite graphs. We remark here that, for each complete multipartite graph G , it is easy (as we will see in Section 3) to find a bisection H of G such that $d_H(v) \geq \lfloor (d_G(v) - 1)/2 \rfloor$ for all $v \in V(G)$. However, for general graphs, even the following weaker version of the Bollobás-Scott conjecture seems quite difficult to prove (or disprove).

Conjecture 1.3. *There exists some absolute constant $c > 0$ such that every graph G has a bisection H with $d_H(v) \geq (d_G(v) - c)/2$ for all $v \in V(G)$.*

We now turn our discussion to problems on general bipartitions. Answering a question of Erdős, Edwards [6] showed in 1973 that every graph with m edges admits a bipartition V_1, V_2 such that $e(V_1, V_2) \geq m/2 + t(m)/2$, where

$$t(m) := \sqrt{m/2 + 1/16} - 1/4.$$

This bound is best possible for the complete graphs of odd order. Bollobás and Scott [3] extended Edwards' bound by showing that every graph G with m edges has a bipartition V_1, V_2 simultaneously satisfying $e(V_1, V_2) \geq m/2 + t(m)/2$ and $\max\{e(V_1), e(V_2)\} \leq m/4 + t(m)/4$, where both bounds are tight for the complete graphs of odd order.

Scott [13] provided an interesting viewpoint by introducing norm for partitions. For a real number $\lambda > 0$ and a bipartition V_1, V_2 of a graph G , define the ℓ_λ -norm of (V_1, V_2) to be $(e(V_1)^\lambda + e(V_2)^\lambda)^{1/\lambda}$. Then to maximize $e(V_1, V_2)$ is equivalent to minimize the ℓ_1 -norm of (V_1, V_2) , while minimizing $\max\{e(V_1), e(V_2)\}$ is the same as minimizing the ℓ_∞ -norm of (V_1, V_2) . It is natural to consider other norms. In particular, Scott asked for the maximum of

$$\min_{V(G)=V_1 \cup V_2} e(V_1)^2 + e(V_2)^2$$

over graphs G with m edges, see Problem 3.18 in [13]. We provide an answer to this question by proving the following general result.

Theorem 1.4. *Let m be any positive integer and $\lambda \geq 1$ be any real number. Then, for any graph G with m edges,*

$$\min_{V(G)=V_1 \cup V_2} e(V_1)^\lambda + e(V_2)^\lambda \leq \binom{t(m)}{2}^\lambda + \binom{t(m) + 1}{2}^\lambda.$$

Moreover, the equality holds if and only if G is a complete graph of odd order.

We also consider analogous questions for k -partitions in Section 4.

Though Edward's bound is tight for all integers $m = \binom{n}{2}$, Erdős [7] conjectured that the difference between Edwards' bound and the truth can still be arbitrarily large for other m . This was confirmed by Alon [1]: every graph with $m = n^2/2$ edges admits a bipartition V_1, V_2 such that $e(V_1, V_2) \geq m/2 + t(m)/2 + \Omega(m^{1/4})$. Bollobás and Scott [5, 13] made a similar conjecture for $\max\{e(V_1), e(V_2)\}$: for certain m , $\max\{e(V_1), e(V_2)\}$ can be arbitrary far from $m/4 + t(m)/4$. Ma and Yu [12] proved that every graph with $m = n^2/2$ edges admits a bipartition V_1, V_2 such that $\max\{e(V_1), e(V_2)\} \leq m/4 + t(m)/4 - \Omega(m^{1/4})$. Another result in the same spirit was given by Hofmeister and Lefmann [9] that any graph with $\binom{kn}{2}$ edges has a k -partition V_1, \dots, V_k with $\sum_{i=1}^k e(V_i) \leq k \binom{n}{2}$, which beats the trivial upper bound $\frac{1}{k} \binom{kn}{2}$.

Motivated by these results, Scott asked the following question: does every graph G with $\binom{kn}{2}$ edges have a vertex partition into k sets, each of which contains at most $\binom{n}{2}$ edges? (See Problem 3.9 in [13].) We show that the answer to this question is negative for $k = 2$.

Theorem 1.5. *There exist infinitely many positive integers n and for each such n there is a graph G with $\binom{2n}{2}$ edges, such that, for every bipartition V_1, V_2 of G , $\max\{e(V_1), e(V_2)\} \geq \binom{n}{2} + 5n/48$.*

This paper is organized as follows. We prove Theorem 1.1 in Section 2, and then investigate complete multipartite graphs for the Bollobás-Scott conjecture in Section 3. In Section 4, we discuss the questions of Scott and complete the proofs of Theorems 1.4 and 1.5.

2 Hartke-Seacrest conjecture

In this section, we prove Theorem 1.1. We need two operations on a sequence of non-negative integers. Let $\pi = (d_1, \dots, d_n)$ with $d_1 \geq \dots \geq d_n$. By removing d_i from π and subtracting 1 from the d_i remaining elements of π with lowest indices, we obtain a new sequence $\pi' = (d'_1, \dots, d'_{i-1}, d'_{i+1}, \dots, d'_n)$, and we say that π' is obtained from π by *laying off* d_i . This operation was introduced by Kleitman and Wang [11], and they proved the following.

Lemma 2.1 (Kleitman-Wang [11]). *For any $i \in [n]$, the sequence $\pi = (d_1, \dots, d_n)$ with $d_1 \geq \dots \geq d_n$ is graphic if and only if the sequence π' obtained from π by laying off d_i is graphic.*

It is easy to see that the sequence π' obtained from π by laying off d_i need not be non-increasing. To avoid this issue, Hartke and Seacrest [10] introduced a variation of the above operation. Choose a fixed $i \in [n]$. Let s be the smallest value among the d_i largest elements in π , not including the i th element of π (namely, d_i itself). Let $S = \{j \in [n] - \{i\} : d_j > s\}$. Note that $|S| < d_i$. Let T be the set of $d_i - |S|$ largest indices j with $j \neq i$ and $d_j = s$. Then by *laying off d_i with order*, we remove d_i from π and subtract 1 from d_j for all $j \in S \cup T$. If $\pi' = (d'_1, \dots, d'_{i-1}, d'_{i+1}, \dots, d'_n)$ denotes the new sequence, then it has the monotone property $d'_1 \geq \dots \geq d'_n$.

Clearly, the sequence obtained from π by laying off d_i with order is just a permutation of the sequence obtained from π by laying off d_i . So the following is true.

Lemma 2.2 (Hartke and Seacrest [10]). *For any $i \in [n]$, the sequence $\pi = (d_1, \dots, d_n)$ with $d_1 \geq \dots \geq d_n$ is graphic if and only if the sequence obtained from π by laying off d_i with order is graphic.*

We give a brief outline of our proof of Theorem 1.1. We choose two consecutive elements d_ℓ and $d_{\ell+1}$ of π . Using Lemma 2.2 we obtain a new graphic sequence π'' of length $n - 2$ by first laying off $d_{\ell+1}$ with order and then laying off d_ℓ with order. By induction, π'' has an $(n - 2)$ -vertex realization F whose parity bisection J has the desired property. We then show that one can form G from F by adding two new vertices (for d_ℓ and $d_{\ell+1}$) and choosing their neighbors, so that the parity bisection of G satisfies Theorem 1.1.

Proof of Theorem 1.1. We apply induction on the length n of the graphic sequence $\pi = (d_1, \dots, d_n)$ with $d_1 \geq \dots \geq d_n$. The assertion is trivial when $n = 1, 2$. So we may assume that $n \geq 3$ and the assertion holds for all graphic sequences with length less than n . Then there exist two consecutive elements of π that are identical; so let $\ell \in [n - 1]$ be fixed such that

$$d_\ell = d_{\ell+1} = k.$$

Let $\pi' = (d'_1, \dots, d'_\ell, d'_{\ell+2}, \dots, d'_n)$ be the sequence obtained from π by laying off $d_{\ell+1}$ with order. Let $\pi'' = (d''_1, \dots, d''_{\ell-1}, d''_{\ell+2}, \dots, d''_n)$ be the sequence obtained from π' by laying off d'_ℓ with order. By Lemma 2.2, π' and π'' both are graphic sequences.

Let $\omega = (f_1, \dots, f_{n-2})$ be the sequence obtained from π with d_ℓ and $d_{\ell+1}$ removed, and re-indexed so that the indices are consecutive, i.e., $f_i = d_i$ for $i \in [\ell - 1]$ and $f_i = d_{i+2}$ for $i \in [n - 2] \setminus [\ell - 1]$. Let $\omega' = (f'_1, \dots, f'_{n-2})$ be the sequence obtained from π' with d'_ℓ removed, and re-indexed so that the indices are consecutive. Also, let $\omega'' = (f''_1, \dots, f''_{n-2})$ be the sequence obtained from π'' by re-indexing so that the indices are consecutive. Note that ω'' is a graphic sequence.

To turn a realization of ω'' to a realization of π , we need to track the changes between f_i and f''_i for all $i \in [n - 2]$. Note that $0 \leq f_i - f''_i \leq 2$. Let

$$X_1 = \{i \in [n - 2] : f''_i = f_i - 1\}, \quad X_2 = \{i \in [n - 2] : f''_i = f_i - 2\}$$

and

$$K = d'_\ell = \begin{cases} k - 1 & \text{if } d'_\ell = d_\ell - 1, \\ k & \text{if } d'_\ell = d_\ell. \end{cases}$$

So $K = \sum_{i \in [n-2]} |f_i - f'_i| = \sum_{i \in [n-2]} |f'_i - f''_i|$; hence

$$|X_1| + 2|X_2| = 2K. \tag{2}$$

We now prove two claims asserting certain properties on X_1 and X_2 . For convenience, we introduce some notation. For nonempty sets A and B of integers, we write $A < B$ if the maximum integer in A is less than the minimum integer in B . A set S of integers is *consecutive* if it consists of consecutive integers. A sequence of pairwise disjoint sets, A_1, \dots, A_t , of integers is said to be *consecutive* if $A_1 \cup \dots \cup A_t$ is consecutive and, for any $i, j \in [t]$ with $i < j$ and A_i and A_j nonempty, we have $A_i < A_j$.

Claim 1. There exist consecutive sets R_1, R_2, R'_1, R'_2, Q such that $X_1 = R'_1 \cup R'_2$ and $X_2 = R_1 \cup R_2$ such that

- (a) the sequence R_1, R'_1, Q, R'_2 is consecutive,
- (b) either $R_2 = \emptyset$ or $R_2 = Q$, and
- (c) $f''_i = f''_j + 1$ for all $i \in R'_1, j \in R'_2$.

Proof of Claim 1. Let s be the minimum of the largest K numbers in $\omega = (f_1, \dots, f_{n-2})$. (Note that this s is the same as the s in the definition of laying off $d_{\ell+1}$ with order from π .) In order to keep track whether $f'_i = f_i$ or $f'_i = f_i - 1$ and whether $f''_i = f'_i$ or $f''_i = f'_i - 1$, we divide $[n-2]$ into six pairwise disjoint sets:

$$\begin{aligned} A &= \{i \in [n-2] : f_i \geq s+2\}, & D &= \{i \in [n-2] : f_i = s, f'_i = f_i - 1\}, \\ B &= \{i \in [n-2] : f_i = s+1\}, & E &= \{i \in [n-2] : f_i = s-1\}, \\ C &= \{i \in [n-2] : f_i = s, f'_i = f_i\}, & F &= \{i \in [n-2] : f_i \leq s-2\}. \end{aligned}$$

By the definitions of π' and ω' , we see that A, B, C, D, E, F is consecutive and

$$\begin{aligned} \forall i \in A, & \quad f'_i = f_i - 1 \geq s+1, \\ \forall i \in B, & \quad f'_i = f_i - 1 = s, \\ \forall i \in C, & \quad f'_i = f_i = s, \\ \forall i \in D, & \quad f'_i = f_i - 1 = s-1, \\ \forall i \in E, & \quad f'_i = f_i = s-1, \\ \forall i \in F, & \quad f'_i = f_i \leq s-2. \end{aligned}$$

Thus, it is easy to see that $A \cup B \cup D = \{i \in [n-2] : f'_i = f_i - 1\}$; so $|A| + |B| + |D| = K$.

Let $Y = \{i \in [n-2] : f''_i = f'_i - 1\}$. Then it follows that

$$A \subseteq Y \quad \text{and} \quad |Y| = K = |A| + |B| + |D|.$$

To complete our proof of Claim 1, we distinguish four cases based on relations among the sizes of B, C, D, E .

First, suppose $|C| \geq |B| + |D|$. Let C'' consist of the last $|B| + |D|$ integers in C , and $C' := C \setminus C''$. Then we see that $Y = A \cup C''$. Let $R_1 = A, R_2 = \emptyset, R'_1 = B, R'_2 = C'' \cup D$ and $Q = C'$. It is easy to check that $X_1 = R'_1 \cup R'_2$ and $X_2 = R_1 \cup R_2$, and (a) and (b) holds. Note that $f''_i = s$ for $i \in R'_1$, and $f''_j = s-1$ for $j \in R'_2$; so (c) holds.

Next, suppose $|D| \leq |C| < |B| + |D|$. Let B'' consist of the last $|B| + |D| - |C|$ integers in B , and $B' = B \setminus B''$. Then $Y = A \cup B'' \cup C$. Let $R_1 = A, R_2 = Q = B'', R'_1 = B'$ and $R'_2 = C \cup D$. It is easy to check that $X_1 = R'_1 \cup R'_2$ and $X_2 = R_1 \cup R_2$, and that (a) and (b) holds. Note that $f''_i = s$ for $i \in R'_1$, and $f''_j = s-1$ for $j \in R'_2$; so (c) holds.

Now assume $|C| < |D| \leq |C| + |E|$. Let E'' consist of the last $|D| - |C|$ integers in E , and $E' = E \setminus E''$. Then $Y = A \cup B \cup C \cup E''$. Let $R_1 = A \cup B, R_2 = \emptyset, R'_1 = C \cup D, R'_2 = E''$, and $Q = E'$. It is easy to check that $X_1 = R'_1 \cup R'_2$ and $X_2 = R_1 \cup R_2$, and (a) and (b) holds. Note that $f''_i = s-1$ for $i \in R'_1$ and $f''_j = s-2$ for $j \in R'_2$; so (c) holds.

Finally we consider the case when $|D| > |C| + |E|$. Let D'' consist of the last $|D| - |C| - |E|$ integers in D , and $D' = D \setminus D''$. Then $Y = A \cup B \cup C \cup D'' \cup E$. Let $R_1 = A \cup B,$

$R_2 = Q = D''$, $R'_1 = C \cup D'$ and $R'_2 = E$. It is easy to check that $X_1 = R'_1 \cup R'_2$ and $X_2 = R_1 \cup R_2$, and (a) and (b) holds. Note that $f''_i = s - 1$ for $i \in R'_1$ and $f''_j = s - 2$ for $j \in R'_2$; so (c) holds. \blacksquare

Let $I_1 = \{i \in [n - 2] : i \equiv 1 \pmod{2}\}$ and $I_2 = \{i \in [n - 2] : i \equiv 0 \pmod{2}\}$.

Claim 2. $|X_1 \cap I_1| - |X_1 \cap I_2| \in \{-2, 0, 2\}$. Moreover, $|X_1 \cap I_1| - |X_1 \cap I_2| = 0$ implies $|X_2 \cap I_1| - |X_2 \cap I_2| \in \{-1, 0, 1\}$.

Proof of Claim 2. By (2), we see $|X_1|$ must be even. So $|R'_1|$ and $|R'_2|$ are of the same parity. Since both R'_1 and R'_2 are consecutive, $|X_1 \cap I_1| - |X_1 \cap I_2| \in \{-2, 0, 2\}$.

Now suppose $|X_1 \cap I_1| - |X_1 \cap I_2| = 0$. If $R_2 = \emptyset$, then $X_2 = R_1$ is a consecutive set and thus $|X_2 \cap I_1|$ and $|X_2 \cap I_2|$ differ by at most one. So we may assume $R_2 \neq \emptyset$. Then $R_2 = Q$ by Claim 1. As the sequence R_1, R'_1, Q, R'_2 is consecutive, we see that $X_1 \cup X_2$ is a consecutive set; so $|(X_1 \cup X_2) \cap I_1|$ and $|(X_1 \cup X_2) \cap I_2|$ differ by at most one. Hence, since $|X_1 \cap I_1| - |X_1 \cap I_2| = 0$, $||X_2 \cap I_1| - |X_2 \cap I_2|| \leq 1$. \blacksquare

We are ready to construct a realization of $\pi = (d_1, \dots, d_n)$. Recall that $\omega'' = (f''_1, \dots, f''_{n-2})$ is a graphic sequence. By induction hypothesis, there exists a realization F of ω'' with $V(F) = \{w_1, \dots, w_{n-2}\}$ and $d_F(w_i) = f''_i$ for $i \in [n - 2]$, such that the parity bisection J of F satisfies

$$d_J(w_i) \geq (d_F(w_i) - 1)/2 \quad \text{for all } i \in [n - 2]. \quad (3)$$

Let $W_j = \{w_i : i \equiv j \pmod{2}\}$ for $j \in [2]$.

In what follows, we will construct a graph G as the realization of π such that its parity bisection H of G satisfies $d_H(v) \geq (d_G(v) - 1)/2$ for all $v \in V(G)$, by adding two new vertices a, b (so $V(G) = V(F) \cup \{a, b\}$) and some edges from these two vertices to F (which we will describe in three separate cases). Notice that if $K = k - 1$, then we would add the edge ab as well; so for convenience, let

$$\epsilon = \begin{cases} 1, & \text{if } K = k - 1, \\ 0, & \text{if } K = k. \end{cases}$$

We write $V(G) = \{v_1, \dots, v_n\}$ such that $v_i = w_i$ for $i < \ell$, $\{v_\ell, v_{\ell+1}\} = \{a, b\}$, and $v_i = w_{i-2}$ for $\ell + 1 < i \leq n$.

In view of Claim 2, we consider the following three cases. In each of these three cases, we use a to represent the vertex in $\{v_\ell, v_{\ell+1}\}$ with odd index. So the parity partition of $V(G)$ is

$$V_1 = W_1 \cup \{a\} \quad \text{and} \quad V_2 = W_2 \cup \{b\}.$$

Case 1. $|X_1 \cap I_1| - |X_1 \cap I_2| = 0$.

We know $F \subseteq G$ and $V(G) = V(F) \cup \{a, b\}$, and we need to add edges at a and b to form G , a realization of π . Add ab if $\epsilon = 1$, av_i for all $i \in X_2 \cup (X_1 \cap I_2)$, and bv_j for all $j \in X_2 \cup (X_1 \cap I_1)$. Since $|X_1 \cap I_1| = |X_1 \cap I_2|$, G is a realization of π . Let H denote the parity bisection of G ; so V_1, V_2 are the partition sets of H . We need to show that $d_H(v) \geq (d_G(v) - 1)/2$ for all $v \in V(G)$.

For each w_i with $i \notin X_1 \cup X_2$, its neighborhoods in F, G are the same; so by (3), $d_H(w_i) = d_J(w_i) \geq (d_F(w_i) - 1)/2 = (d_G(w_i) - 1)/2$.

For vertices w_i with $i \in X_2$, we have $d_G(w_i) = d_F(w_i) + 2$ and $d_H(w_i) = d_J(w_i) + 1$; so by (3), $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 = (d_G(w_i) - 1)/2$.

For vertices w_i with $i \in X_1$, we have $d_G(w_i) = d_F(w_i) + 1$ and $d_H(w_i) = d_J(w_i) + 1$; so by (3), $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 > (d_G(w_i) - 1)/2$.

For the vertex a , we have $d_G(a) = |X_2| + |X_1 \cap I_2| + \epsilon$ and $d_H(a) = |X_2 \cap I_2| + |X_1 \cap I_2| + \epsilon$. Note that in this case, by Claim 2, we have $||X_2 \cap I_1| - |X_2 \cap I_2|| \leq 1$, which implies that

$$2d_H(a) - d_G(a) = (|X_2 \cap I_2| - |X_2 \cap I_1|) + |X_1 \cap I_2| + \epsilon \geq -1.$$

Hence, $d_H(a) \geq (d_G(a) - 1)/2$.

Similarly, for the vertex b , we have $d_G(b) = |X_2| + |X_1 \cap I_1| + \epsilon$ and $d_H(b) = |X_2 \cap I_1| + |X_1 \cap I_1| + \epsilon$. Note, by Claim 2, $||X_2 \cap I_1| - |X_2 \cap I_2|| \leq 1$; so

$$2d_H(b) - d_G(b) = (|X_2 \cap I_1| - |X_2 \cap I_2|) + |X_1 \cap I_1| + \epsilon \geq -1.$$

Hence, $d_H(b) \geq (d_G(b) - 1)/2$.

Case 2. $|X_1 \cap I_2| - |X_1 \cap I_1| = 2$.

Recall that $X_1 = R'_1 \cup R'_2$, where each R'_i is consecutive. Thus it follows that $|R'_i \cap I_2| = |R'_i \cap I_1| + 1$ for $i \in [2]$. Since the sequence R_1, R'_1, Q, R'_2 is consecutive and starts from the integer 1, we see that $|R_1 \cap I_2| = |R_1 \cap I_1| - 1$ and $|Q \cap I_2| = |Q \cap I_1| - 1$. Therefore, since $R_2 = \emptyset$ or $R_2 = Q$ (by (b) of Claim 1), we have

$$-2 \leq |X_2 \cap I_2| - |X_2 \cap I_1| \leq -1. \quad (4)$$

We claim that there exists some $z \in X_1 \cap I_2$ with $d_J(w_z) \geq d_F(w_z)/2$. To see this, choose $x \in R'_1 \cap I_2$ and $y \in R'_2 \cap I_2$. By (3), we have $d_J(w_x) \geq (d_F(w_x) - 1)/2$ and $d_J(w_y) \geq (d_F(w_y) - 1)/2$. By (c) of Claim 1, $d_F(w_x) = d_F(w_y) + 1$. Observe that for any vertex u of F , $d_F(u)$ and $2d_J(u) - d_F(u)$ are of the same parity; so $2d_J(w_x) - d_F(w_x)$ and $2d_J(w_y) - d_F(w_y)$ must have different parities. Therefore there exists $z \in \{x, y\}$ such that $d_J(w_z) \geq d_F(w_z)/2$, proving the claim.

We now add edges at a and b to form G from F : add av_i for all $i \in X_2 \cup (X_1 \cap I_2) \setminus \{z\}$, and bv_j for all $j \in X_2 \cup (X_1 \cap I_1) \cup \{z\}$. Since $|X_1 \cap I_2| = |X_1 \cap I_1| + 2$, G is a realization of π . Next we show that the parity bisection H of G satisfies the property that $d_H(v) \geq (d_G(v) - 1)/2$ for all $v \in V(G)$.

For each w_i with $i \notin X_1 \cup X_2$, its neighborhoods in F, G are the same; so by (3), $d_H(w_i) = d_J(w_i) \geq (d_F(w_i) - 1)/2 = (d_G(w_i) - 1)/2$.

For each w_i with $i \in X_2$, $d_G(w_i) = d_F(w_i) + 2$ and $d_H(w_i) = d_J(w_i) + 1$. Hence by (3) and the way we choose z , $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 = (d_G(w_i) - 1)/2$.

For w_i with $i \in X_1 \setminus \{z\}$, we have $d_G(w_i) = d_F(w_i) + 1$ and $d_H(w_i) = d_J(w_i) + 1$; so by (3), $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 > (d_G(w_i) - 1)/2$.

The vertex w_z satisfies $d_G(w_z) = d_F(w_z) + 1$ and $d_H(w_z) = d_J(w_z)$. Hence by (3), $d_H(w_z) = d_J(w_z) \geq d_F(w_z)/2 = (d_G(w_z) - 1)/2$.

For the vertex a , by definition we have $d_G(a) = |X_2| + |X_1 \cap I_2| - 1 + \epsilon$ and $d_H(a) = |X_2 \cap I_2| + |X_1 \cap I_2| - 1 + \epsilon$. By (4) and the fact that $|X_1 \cap I_2| \geq 2$,

$$2d_H(a) - d_G(a) = (|X_2 \cap I_2| - |X_2 \cap I_1|) + |X_1 \cap I_2| - 1 + \epsilon \geq -1.$$

Hence, $d_H(a) \geq (d_G(a) - 1)/2$.

For the vertex b , we have $d_G(b) = |X_2| + |X_1 \cap I_1| + 1 + \epsilon$ and $d_H(b) = |X_2 \cap I_1| + |X_1 \cap I_1| + \epsilon$. This, together with (4), imply that

$$2d_H(b) - d_G(b) = (|X_2 \cap I_1| - |X_2 \cap I_2|) + |X_1 \cap I_1| - 1 + \epsilon \geq 0.$$

Hence, $d_H(b) \geq (d_G(b) - 1)/2$.

Case 3. $|X_1 \cap I_1| - |X_1 \cap I_2| = 2$.

In this case, we have $|R'_i \cap I_1| = |R'_i \cap I_2| + 1$ for $i \in [2]$ (as R'_1 and R'_2 are consecutive). Because R_1, R'_1, Q, R'_2 is consecutive, it follows that $|R_1 \cap I_1| = |R_1 \cap I_2|$ and $|Q \cap I_1| = |Q \cap I_2| - 1$. Since $R_2 = \emptyset$ or $R_2 = Q$ (by (b) of Claim 1),

$$0 \leq |X_2 \cap I_2| - |X_2 \cap I_1| \leq 1. \quad (5)$$

Since $|X_1|$ is even and $|R'_i \cap I_1| = |R'_i \cap I_2| + 1$ for $i \in [2]$, there exist $x \in R'_1 \cap I_1$ and $y \in R'_2 \cap I_1$. By (3) and (c) of Claim 1, we have $d_J(w_x) \geq (d_F(w_x) - 1)/2$, $d_J(w_y) \geq (d_F(w_y) - 1)/2$, and $d_F(w_x) = d_F(w_y) + 1$. Since for any vertex u of F , $d_F(u)$ and $2d_J(u) - d_F(u)$ are of the same parity, $2d_J(w_x) - d_F(w_x)$ and $2d_J(w_y) - d_F(w_y)$ must have different parities. Therefore there exists $z \in \{x, y\}$ such that $d_J(w_z) \geq d_F(w_z)/2$.

We now add edges at a and b to form the graph G : add ab if $\epsilon = 1$, av_i for all $i \in X_2 \cup (X_1 \cap I_2) \cup \{z\}$, and bv_j for all $j \in X_2 \cup (X_1 \cap I_1) \setminus \{z\}$. Since $|X_1 \cap I_1| = |X_1 \cap I_2| + 2$, G is a realization of π . We need to verify that $d_H(v) \geq (d_G(v) - 1)/2$ for all $v \in V(G)$.

If $v = w_i$ for some $i \notin X_1 \cup X_2$, then $d_G(w_i) = d_F(w_i)$ and $d_H(w_i) = d_J(w_i)$; so by (3), $d_H(w_i) = d_J(w_i) \geq (d_F(w_i) - 1)/2 = (d_G(w_i) - 1)/2$.

If $v = w_i$ for some $i \in X_2$, then $d_G(w_i) = d_F(w_i) + 2$ and $d_H(w_i) = d_J(w_i) + 1$; again by (3), $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 = (d_G(w_i) - 1)/2$.

If $v = w_i$ for some $i \in X_1 \setminus \{z\}$, then $d_G(w_i) = d_F(w_i) + 1$ and $d_H(w_i) = d_J(w_i) + 1$; so by (3), $d_H(w_i) = d_J(w_i) + 1 \geq (d_F(w_i) - 1)/2 + 1 > (d_G(w_i) - 1)/2$.

Now suppose $v = w_z$. Note that $d_G(w_z) = d_F(w_z) + 1$ and $d_H(w_z) = d_J(w_z)$. So $d_H(w_z) = d_J(w_z) \geq d_F(w_z)/2 = (d_G(w_z) - 1)/2$.

Suppose $v = a$. Note that $d_G(a) = |X_2| + |X_1 \cap I_2| + 1 + \epsilon$ and $d_H(a) = |X_2 \cap I_2| + |X_1 \cap I_2| + \epsilon$. By (5),

$$2d_H(a) - d_G(a) = (|X_2 \cap I_2| - |X_2 \cap I_1|) + |X_1 \cap I_2| - 1 + \epsilon \geq -1.$$

So $d_H(a) \geq (d_G(a) - 1)/2$.

Finally, suppose $v = b$. We have $d_G(b) = |X_2| + |X_1 \cap I_1| - 1 + \epsilon$ and $d_H(b) = |X_2 \cap I_1| + |X_1 \cap I_1| - 1 + \epsilon$. By (5) and the fact that $|X_1 \cap I_1| \geq 2$,

$$2d_H(b) - d_G(b) = (|X_2 \cap I_1| - |X_2 \cap I_2|) + |X_1 \cap I_1| - 1 + \epsilon \geq 0.$$

So $d_H(b) \geq (d_G(b) - 1)/2$. ■

3 Complete multipartite graphs

For convenience, we say that a bisection H of a graph G is *good* if for each $v \in V(G)$, $2d_H(v) \geq d_G(v) - 1$. Thus the Bollobás-Scott conjecture says that every graph contains a good bisection. Here we discuss which complete multipartite graphs have good bisections. Throughout the rest of this section, let $G := K_{r_1, \dots, r_k}$, and let X_1, \dots, X_k denote the partition sets of G with $|X_i| = r_i$ for all $i \in [k]$.

First, we note that whenever $|V(G)|$ is even, G has a good bisection. Since $|V(G)|$ is even, $V(G)$ has a partition V_1, V_2 such that $|V_1| = |V_2|$, $||X_i \cap V_1| - |X_i \cap V_2|| = 1$ if $|X_i|$ is odd, and $|X_i \cap V_1| = |X_i \cap V_2|$ if $|X_i|$ is even. Let H denote the maximum bisection of G with partition sets V_1 and V_2 . For any $v \in V(G)$, $v \in X_i$ for some $i \in [k]$. Note that $d_G(v) = |V(G)| - |X_i|$ and $d_H(v) \geq (|V(G)| - |X_i| - 1)/2 = (d_G(v) - 1)/2$.

We will see that this is not always the case when $|V(G)|$ is odd. The main result of this section is a necessary and sufficient condition for a complete multipartite graph with odd order to contain a good bisection. As a consequence, we show that for many complete multipartite graphs G , G (and even G minus an edge) does not have a good bisection. (However, it is not hard to see that such G does have a bisection H such that for each $v \in V(G)$, $d_H(v) \geq \lfloor (d_G(v) - 1)/2 \rfloor$.)

For a bisection H of G with partition sets V_1 and V_2 , we say that X_i *crosses* H if $X_i \cap V_j \neq \emptyset$ for $j \in [2]$. For a subset $W \subseteq V(G)$, let $\overline{W} = V(G) \setminus W$. We need two easy lemmas.

Lemma 3.1. *Let $G = K_{r_1, \dots, r_k}$ and let X_i , $i \in [k]$, be the partition sets of G . Suppose H is a good bisection of G with partition sets V_1 and V_2 . Then the following statements hold for $i \in [k]$.*

- (i) *If X_i crosses H and $|\overline{X_i}|$ is even then $|\overline{X_i} \cap V_1| = |\overline{X_i} \cap V_2|$ and $||X_i \cap V_1| - |X_i \cap V_2|| \leq 1$.*
- (ii) *If X_i crosses H and $|\overline{X_i}|$ is odd then $||\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2|| = 1$ and $||X_i \cap V_1| - |X_i \cap V_2|| \leq 2$.*

Proof. Suppose X_i crosses H . Then there exist $v_j \in X_i \cap V_j$ for $j \in [2]$. Note that $d_G(v_1) = d_G(v_2) = |\overline{X_i}|$. Thus, since H is a good bisection, $d_H(v_j) = |\overline{X_i} \cap V_{3-j}| \geq \lfloor |\overline{X_i}|/2 \rfloor$ for $j \in [2]$. Notice that $|\overline{X_i} \cap V_1| + |\overline{X_i} \cap V_2| = |\overline{X_i}|$. So $||\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2|| \leq 1$.

If $|\overline{X_i}|$ is even then $|\overline{X_i} \cap V_1| = |\overline{X_i} \cap V_2|$, and (i) holds. So assume $|\overline{X_i}|$ is odd. Since $||V_1| - |V_2|| \leq 1$, $||X_i \cap V_1| - |X_i \cap V_2|| \leq 2$, and (ii) holds. ■

Lemma 3.2. *Let $G = K_{r_1, \dots, r_k}$ with $|V(G)|$ odd, let X_1, \dots, X_k be the partition sets of G , and let H be a good bisection of G with partition sets V_1, V_2 such that $|V_1| = |V_2| + 1$. Let*

$$\mathcal{X}'_0 = \{X_i : i \in [k], X_i \text{ crosses } H, |X_i| \equiv 0 \pmod{2}, \text{ and } ||X_i \cap V_1| - |X_i \cap V_2|| = 2\}$$

and

$$\mathcal{X}_1 = \{X_i : i \in [k], X_i \text{ crosses } H, \text{ and } |X_i| \equiv 1 \pmod{2}\}.$$

Let $W_1 = \bigcup_{X_i \in \mathcal{X}_1} X_i$, $W_0 = \bigcup_{X_i \in \mathcal{X}'_0} X_i$, $|\mathcal{X}_1| = t$, and $|\mathcal{X}'_0| = t'$. Then

- (i) $|W_1 \cap V_1| - |W_1 \cap V_2| = t$,

(ii) $|W_0 \cap V_1| - |W_0 \cap V_2| = 2t'$, and

(iii) $|\overline{W_1 \cup W_0} \cap V_1| - |\overline{W_1 \cup W_0} \cap V_2| = -(t + 2t' - 1)$.

Proof. First, we prove (i). If $W_1 = \emptyset$, then $|W_1 \cap V_1| - |W_1 \cap V_2| = 0 = t$. So assume $W_1 \neq \emptyset$ and let $X_i \in \mathcal{X}_1$. Then $|X_i|$ is odd and X_i crosses H . Hence, since $|V(G)|$ is odd, $|\overline{X_i}|$ is even. By Lemma 3.1, $|\overline{X_i} \cap V_1| = |\overline{X_i} \cap V_2|$. Because $|V_1| = |V_2| + 1$, $|X_i \cap V_1| - |X_i \cap V_2| = 1$. Hence, $|W_1 \cap V_1| - |W_1 \cap V_2| = \sum_{X_i \in \mathcal{X}_1} (|X_i \cap V_1| - |X_i \cap V_2|) = t$.

We now prove (ii). If $W_0 = \emptyset$ then $t' = 0$ and the result holds trivially. So assume $W_0 \neq \emptyset$ and let $X_i \in \mathcal{X}'_0$. Then $|X_i|$ is even and X_i crosses H . Since $|V(G)|$ is odd, $|\overline{X_i}|$ is odd. By Lemma 3.1, $||\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2|| = 1$, and by the definition of \mathcal{X}'_0 , $||X_i \cap V_1| - |X_i \cap V_2|| = 2$. Therefore, because $|V_1| = |V_2| + 1$, $|X_i \cap V_1| - |X_i \cap V_2| = 2$ (as well as $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = -1$). Hence $|W_0 \cap V_1| - |W_0 \cap V_2| = \sum_{X_i \in \mathcal{X}'_0} (|X_i \cap V_1| - |X_i \cap V_2|) = 2t'$.

It is easy to see that (iii) follows from (i), (ii) and the assumption $|V_1| = |V_2| + 1$. \blacksquare

We now give a necessary and sufficient condition for a complete multipartite graph with odd order to admit a good bisection. Let $G = K_{r_1, \dots, r_k}$ with partition sets X_i , $i \in [k]$, such that $|X_i| = r_i$. Let $\mathcal{X} = \{X_i : i \in [k]\}$, $\mathcal{S}_1 = \{X_i : i \in [k] \text{ and } |X_i| \equiv 1 \pmod{2}\}$ and $\mathcal{S}_0 = \{X_i : i \in [k] \text{ and } |X_i| \equiv 0 \pmod{2}\}$. For any $\mathcal{A} \subseteq \mathcal{X}$, let $s(\mathcal{A}) = \sum_{X_i \in \mathcal{A}} |X_i|$. We say that \mathcal{A} is *good* if there exists $\mathcal{A}' \subseteq \mathcal{A}$ such that

$$s(\mathcal{A}') = s(\mathcal{A})/2 + (m + 2n - 1)/2,$$

where $m = |\mathcal{S}_1 \setminus \mathcal{A}|$ and n is a nonnegative integer with $n \leq |\mathcal{S}_0 \setminus \mathcal{A}|$.

Proposition 3.3. *Let $G = K_{r_1, \dots, r_k}$ with partition sets X_1, \dots, X_k , and assume $|V(G)|$ is odd. Let $\mathcal{X} = \{X_i : i \in [k]\}$. Then G has a good bisection if and only if \mathcal{X} has a good subset.*

Proof. First, we prove that if G has a good bisection, then \mathcal{X} has a good subset. Let H be a good bisection of G and let V_1, V_2 be the corresponding partition sets of H . Since $|V(G)|$ is odd, we may assume that $|V_1| = |V_2| + 1$. Let $\mathcal{S} = \{X_i : i \in [k] \text{ and } X_i \text{ crosses } H\}$, $\mathcal{X}_1 = \{X_i : i \in [k], X_i \text{ crosses } H, \text{ and } |X_i| \equiv 1 \pmod{2}\}$, $\mathcal{X}'_0 = \{X_i : i \in [k], X_i \text{ crosses } H, |X_i| \equiv 0 \pmod{2}, \text{ and } ||X_i \cap V_1| - |X_i \cap V_2|| = 2\}$, and $\mathcal{X}''_0 = \{X_i : i \in [k], X_i \text{ crosses } H, |X_i| \equiv 0 \pmod{2}, \text{ and } |X_i \cap V_1| = |X_i \cap V_2|\}$.

By Lemma 3.1 (ii) and the assumption $|V_1| = |V_2| + 1$, for every $X_i \in \mathcal{X}$ crossing H and $|X_i|$ even, $||X_i \cap V_1| - |X_i \cap V_2|| = 2$ or $|X_i \cap V_1| = |X_i \cap V_2|$. Hence, $\mathcal{S} = \mathcal{X}_1 \cup \mathcal{X}'_0 \cup \mathcal{X}''_0$. Let $\mathcal{A} = \mathcal{X} \setminus \mathcal{S}$.

Moreover, let $W_1 = \bigcup_{X_i \in \mathcal{X}_1} X_i$, $W_0 = \bigcup_{X_i \in \mathcal{X}'_0} X_i$, $W'_0 = \bigcup_{X_i \in \mathcal{X}''_0} X_i$ and $W_2 = \bigcup_{X_i \in \mathcal{A}} X_i$. Then $V(H) = W_1 \cup W_0 \cup W'_0 \cup W_2$. By the definition of \mathcal{X}''_0 , $|W'_0 \cap V_1| = |W'_0 \cap V_2|$. Let $|\mathcal{X}_1| = t$ and $|\mathcal{X}'_0| = t'$; then by Lemma 3.2 (iii), $|(W'_0 \cup W_2) \cap V_2| - |(W'_0 \cup W_2) \cap V_1| = t + 2t' - 1$. Combining these two equalities, we get

$$|W_2 \cap V_2| - |W_2 \cap V_1| = |(W'_0 \cup W_2) \cap V_2| - |(W'_0 \cup W_2) \cap V_1| = t + 2t' - 1. \quad (6)$$

Since X_i does not cross H for any $X_i \subseteq \mathcal{A}$, there exists $\mathcal{A}' \subseteq \mathcal{A}$ such that $W_2 \cap V_2 = \bigcup_{X_i \in \mathcal{A}'} X_i$ and $W_2 \cap V_1 = \bigcup_{X_i \in \mathcal{A} \setminus \mathcal{A}'} X_i$. Now, $|W_2 \cap V_2| = s(\mathcal{A}')$ and $|W_2 \cap V_1| = s(\mathcal{A}) - s(\mathcal{A}')$.

Thus $s(\mathcal{A}') = s(\mathcal{A})/2 + (t + 2t' - 1)/2$ by (6). Note that $t = |\mathcal{X}_1| = |\mathcal{S}_1 \setminus \mathcal{A}|$ and $t' = |\mathcal{X}'_0| \leq |\mathcal{X}'_0 \cup \mathcal{X}''_0| = |\mathcal{S}_0 \setminus \mathcal{A}|$. So \mathcal{A} is a good subset of \mathcal{X} .

Now, we prove that if \mathcal{X} has a good subset, then G has a good bisection. Let \mathcal{A} be a good subset of \mathcal{X} . Then there exists $\mathcal{A}' \subseteq \mathcal{A}$ such that $s(\mathcal{A}') = s(\mathcal{A})/2 + (m + 2n - 1)/2$, where $m = |\mathcal{S}_1 \setminus \mathcal{A}|$ and $n \leq |\mathcal{S}_0 \setminus \mathcal{A}|$. Let $\mathcal{S}'_0 \subseteq \mathcal{S}_0 \setminus \mathcal{A}$ with $|\mathcal{S}'_0| = n$, and let $\mathcal{S}''_0 = (\mathcal{S}_0 \setminus \mathcal{A}) \setminus \mathcal{S}'_0$. We partition $V(G)$ into V_1 and V_2 such that

- $|X_i \cap V_1| - |X_i \cap V_2| = 1$ if $X_i \in \mathcal{S}_1 \setminus \mathcal{A}$,
- $|X_i \cap V_1| - |X_i \cap V_2| = 2$ if $X_i \in \mathcal{S}'_0$,
- $|X_i \cap V_1| - |X_i \cap V_2| = 0$ if $X_i \subseteq \mathcal{S}''_0$, and
- $X_i \subseteq V_1$ if $X_i \in \mathcal{A} \setminus \mathcal{A}'$, and $X_i \subseteq V_2$ if $X_i \in \mathcal{A}'$.

Then

$$|V_1| - |V_2| = 2|\mathcal{S}'_0| + |\mathcal{S}_1 \setminus \mathcal{A}| + s(\mathcal{A}) - 2s(\mathcal{A}') = 1.$$

Let H be the bisection of G with partition sets V_1 and V_2 and edge set $E(H) = \{uv \in E(G) : u \in V_1 \text{ and } v \in V_2\}$. Next, we show that H is a good bisection of G . Note that, for each $v \in X_i \subseteq V(G)$, $d_G(v) = |\overline{X_i}|$, $d_H(v) = |\overline{X_i} \cap V_1|$ if $v \in V_2$, and $d_H(v) = |\overline{X_i} \cap V_2|$ if $v \in V_1$. Also note that

$$|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = (|V_1| - |V_2|) - (|X_i \cap V_1| - |X_i \cap V_2|) = 1 - (|X_i \cap V_1| - |X_i \cap V_2|).$$

If $v \in X_i$ for some $X_i \in \mathcal{S}_1 \setminus \mathcal{A}$, then $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = 1 - 1 = 0$; so $d_H(v) = d_G(v)/2$.

If $v \in X_i$ for some $X_i \in \mathcal{S}'_0$ then $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = 1 - 2 = -1$; so $d_H(v) \geq (d_G(v) - 1)/2$.

If $v \in X_i$ for some $X_i \in \mathcal{S}''_0$ then $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = 1 - 0 = 1$; so $d_H(v) \geq (d_G(v) - 1)/2$.

If $v \in X_i \cap V_2$ for some $X_i \in \mathcal{A}$ then $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = 1 + |X_i| > 0$; so $d_H(v) = |\overline{X_i} \cap V_1| \geq d_G(v)/2$.

Finally, suppose $v \in X_i \cap V_1$ for some $X_i \in \mathcal{A}$. Then $|\overline{X_i} \cap V_1| - |\overline{X_i} \cap V_2| = 1 - |X_i|$, i.e., $|\overline{X_i} \cap V_2| - |\overline{X_i} \cap V_1| = |X_i| - 1 \geq 0$. This implies that $d_H(v) = |\overline{X_i} \cap V_2| \geq d_G(v)/2$. ■

Proof of Proposition 1.2. Let $G = K_{r_1, r_2, r_3}$ and $\mathcal{X} = \{X_1, X_2, X_3\}$ such that X_1, X_2, X_3 are the partition sets of G and $|X_i| = r_i$ for $i \in [3]$. Let $\mathcal{S}_0 = \{X_i : i \in [3] \text{ and } |X_i| \equiv 0 \pmod{2}\}$ and $\mathcal{S}_1 = \{X_i : i \in [3] \text{ and } |X_i| \equiv 1 \pmod{2}\}$. Then $\mathcal{S}_0 = \emptyset$ and $\mathcal{S}_1 = \mathcal{X}$.

If \mathcal{X} has no good subset then the assertion follows from Proposition 3.3. So assume that \mathcal{A} is a good subset of \mathcal{X} with $\mathcal{A}' \subseteq \mathcal{A}$ such that $s(\mathcal{A}') = s(\mathcal{A})/2 + (m + 2n - 1)/2$, where $m = |\mathcal{S}_1 \setminus \mathcal{A}| = 3 - |\mathcal{A}|$ and $n \leq |\mathcal{S}_0 \setminus \mathcal{A}| = 0$. So

$$s(\mathcal{A}') = s(\mathcal{A})/2 - |\mathcal{A}|/2 + 1.$$

It is easy to see that $\mathcal{A} \neq \emptyset$. Since $r_i \geq 3 \geq |\mathcal{A}|$, $\mathcal{A}' \neq \emptyset$ and, hence, $|\mathcal{A}| \neq 1$. Since r_1, r_2, r_3 are all distinct, $|\mathcal{A}| \neq 2$. So $|\mathcal{A}| = 3$. Now a straightforward analysis shows that for some $i \in [3]$, $r_i \in \{\lfloor (r_1 + r_2 + r_3)/2 \rfloor, \lceil (r_1 + r_2 + r_3)/2 \rceil\}$. This is a contradiction. ■

Proposition 3.3 characterizes those complete multipartite graphs which do not have a good bisection. The next result says that there are more such examples.

Proposition 3.4. *Let $G = K_{r_1, \dots, r_k}$ where $r_i \geq 7$ for every $i \in [k]$. Suppose G does not have a good bisection. Then for any edge $e \in E(G)$, $G - e$ does not have a good bisection.*

Proof. Assume, to the contrary, that $G' = G - e$ has a good bisection H' with partition sets V_1, V_2 . We may assume that $E(H') = \{xy \in E(G') : x \in V_1 \text{ and } y \in V_2\}$. Then for every vertex $v \in V(G')$, $d_{H'}(v) \geq (d_{G'}(v) - 1)/2$. Let H be the bisection of G with partition sets V_1 and V_2 such that $E(H) = \{xy \in E(G) : x \in V_1 \text{ and } y \in V_2\}$. Let $e = uw$.

Then $d_G(v) = d_{G'}(v)$ for all $v \in V(G) \setminus \{u, w\}$, and $d_G(v) = d_{G'}(v) + 1$ for $v \in \{u, w\}$. Also, we have $d_H(v) = d_{H'}(v)$ for all $v \in V(G) \setminus \{u, w\}$, and $d_{H'}(v) \leq d_H(v) \leq d_{H'}(v) + 1$ for $v \in \{u, w\}$.

Since H is not a good bisection of G , there exists a vertex $v \in V(G)$ such that $d_H(v) < (d_G(v) - 1)/2$. So we have

$$(d_{G'}(v) - 1)/2 \leq d_{H'}(v) \leq d_H(v) < (d_G(v) - 1)/2 \leq d_{G'}(v)/2$$

which implies that $d_{G'}(v)$ is odd (since $d_H(v)$ is an integer), $d_G(v) = d_{G'}(v) + 1$, and $(d_{G'}(v) - 1)/2 = d_{H'}(v) = d_H(v)$. Since $d_G(v) = d_{G'}(v) + 1$, $v \in \{u, w\}$.

Assume, without loss of generality, that $v = u \in X_i \cap V_1$, where X_1, \dots, X_k are the partition sets of G . (Then $w \notin X_i$ as $uw \in E(G)$.) So $d_{G'}(u) = |\overline{X}_i| - 1$, $|\overline{X}_i|$ is even, and $w \in V_1$. Thus, $|V_2 \cap \overline{X}_i| = d_{H'}(u) = (d_{G'}(u) - 1)/2 = |\overline{X}_i|/2 - 1$. Therefore, $|V_1 \cap \overline{X}_i| = |\overline{X}_i| - |V_2 \cap \overline{X}_i| = |\overline{X}_i|/2 + 1$. So $|V_1 \cap \overline{X}_i| - |V_2 \cap \overline{X}_i| = 2$. Because $||V_1| - |V_2|| = 1$, $||V_1 \cap X_i| - |V_2 \cap X_i|| \leq 3$. Since $|X_i| \geq 7$, $|V_1 \cap X_i| \geq 2$. Therefore, there exists a vertex $v_1 \in V_1 \cap X_i$ such that $v_1 \neq u$. Also, $v_1 \neq w$ since $w \notin X_i$. Then $d_{G'}(v_1) = d_G(v_1) = |\overline{X}_i|$ is even. Thus $d_{H'}(v_1) = |V_2 \cap \overline{X}_i| = |\overline{X}_i|/2 - 1 < (d_{G'}(v_1) - 1)/2$. This contradicts the assumption that H' is a good bisection of G' . \blacksquare

4 Scott's questions on bipartitions

In this section, we address two questions of Scott [13] on bipartitions of graphs. First, we prove Theorem 1.4 on ℓ_λ -norm of bipartitions (with $\lambda \geq 1$), for which we need a result of Bollobás and Scott [3] on judicious bipartitions. Recall the definition of $t(m) = \sqrt{m/2 + 1/16} - 1/4$.

Lemma 4.1 (Bollobás and Scott). *Let G be a graph with m edges. Then there exists a bipartition $V(G) = V_1 \cup V_2$ such that $e(V_1, V_2) \geq m/2 + t(m)/2$ and $\max\{e(V_1), e(V_2)\} \leq m/4 + t(m)/4$. Moreover, if for every such bipartition $V(G) = V_1 \cup V_2$ it always holds that $\max\{e(V_1), e(V_2)\} = m/4 + t(m)/4$, then G must be a complete graph of odd order.*

Proof of Theorem 1.4. Let $\lambda \geq 1$ and let G be a graph with m edges. By Lemma 4.1, there is a bipartition V_1, V_2 of $V(G)$ such that $e(V_1, V_2) \geq m/2 + t(m)/2$ and, for $i \in [2]$, $e(V_i) \leq m/4 + t(m)/4 = \binom{t(m)+1}{2}$. Note that

$$e(V_1) + e(V_2) = m - e(V_1, V_2) \leq m/2 - t(m)/2 = t(m)^2.$$

Without loss of generality, we assume that $e(V_1) \geq e(V_2)$. Then $e(V_2) \leq t(m)^2/2$.

We claim that $e(V_1)^\lambda + e(V_2)^\lambda \leq \binom{t(m)}{2}^\lambda + \binom{t(m)+1}{2}^\lambda$. This is true if $e(V_2) \leq \binom{t(m)}{2}$. So we may assume $\binom{t(m)}{2} \leq e(V_2) \leq t(m)^2/2$. For $\lambda \geq 1$, the function $f(x) = (t(m)^2 - x)^\lambda + x^\lambda$ is strictly decreasing when $\binom{t(m)}{2} \leq x \leq t(m)^2/2$. Therefore,

$$e(V_1)^\lambda + e(V_2)^\lambda \leq (t(m)^2 - e(V_2))^\lambda + e(V_2)^\lambda \leq \binom{t(m)}{2}^\lambda + \binom{t(m)+1}{2}^\lambda.$$

Now assume that for every bipartition V_1, V_2 of $V(G)$, we have $e(V_1)^\lambda + e(V_2)^\lambda = \binom{t(m)}{2}^\lambda + \binom{t(m)+1}{2}^\lambda$. Then it follows from the above arguments, $e(V_2) = \binom{t(m)}{2}$ and $e(V_1) + e(V_2) = t(m)^2$. So $\max\{e(V_1), e(V_2)\} = e(V_1) = t(m)^2 - \binom{t(m)}{2} = m/4 + t(m)/4$. By Lemma 4.1, G is a complete graph of odd order. \blacksquare

Remark. From the above proof, we see that actually $V(G)$ has a bipartition V_1, V_2 such that $e(V_1)^\lambda + e(V_2)^\lambda \leq \binom{t(m)}{2}^\lambda + \binom{t(m)+1}{2}^\lambda$ for all $\lambda \geq 1$.

To extend Theorem 1.4 to k -partitions for $k \geq 3$, we need the following result of Xu and Yu [14] on k -partitions.

Lemma 4.2 (Xu and Yu). *Let G be a graph with m edges and let $k \geq 3$ be a positive integer. Then there exists a k -partition $V(G) = V_1 \cup \dots \cup V_k$ such that*

$$e(V_1, \dots, V_k) \geq \frac{k-1}{k}m + \frac{k-1}{k}t(m) - \frac{17k}{8},$$

and for $i \in [k]$,

$$e(V_i) \leq \frac{m}{k^2} + \frac{k-1}{k^2}t(m).$$

We now determine the ℓ_λ -norm (where $\lambda \geq 1$) for k -partitions, up to an additive term $O(m^{\lambda-1})$. The proof is similar to the bipartition case.

Theorem 4.3. *Let $k \geq 3$ be an integer and $\lambda \geq 1$ be a real number. Then any graph G with m edges has a k -partition $V(G) = V_1 \cup \dots \cup V_k$ such that*

$$\sum_{i=1}^k e(V_i)^\lambda \leq \frac{1}{k^{2\lambda-1}}m^\lambda - \frac{(k-1)\lambda}{k^{2\lambda-1}}m^{\lambda-1}t(m) + O(m^{\lambda-1}).$$

Proof. Let G be a graph with m edges. By Lemma 4.2, there exists a k -partition $V(G) = V_1 \cup \dots \cup V_k$ such that

$$\sum_{i=1}^k e(V_i) = m - e(V_1, \dots, V_k) \leq \frac{m}{k} - \frac{k-1}{k}t(m) + \frac{17k}{8}$$

and for $i \in [k]$,

$$e(V_i) \leq \frac{m}{k^2} + \frac{k-1}{k^2}t(m).$$

Without loss of generality, let $e(V_1) \geq e(V_2) \geq \dots \geq e(V_k)$ and let $\alpha := e(V_1) - m/k^2$.

If $\alpha \leq -\frac{k-1}{k^2}t(m)$ then

$$\sum_{i=1}^k e(V_i)^\lambda \leq k \left(\frac{m}{k^2} - \frac{k-1}{k^2}t(m) \right)^\lambda \leq \frac{1}{k^{2\lambda-1}}m^\lambda - \frac{(k-1)\lambda}{k^{2\lambda-1}}m^{\lambda-1}t(m) + O(m^{\lambda-1}).$$

So we may assume that

$$-\frac{k-1}{k^2}t(m) \leq \alpha \leq \frac{k-1}{k^2}t(m). \quad (7)$$

Note that we may assume $\sum_{i=1}^k e(V_i) = \frac{m}{k} - \frac{k-1}{k}t(m) + \frac{17k}{8}$. Also note that $\sum_{i=1}^k e(V_i)^\lambda$ increases if we replace $e(V_k)$ by $e(V_k) - 1$ and replaces $e(V_i)$ by $e(V_i) + 1$, for any $i \in [k-1]$. Therefore, we may further assume that $e(V_1) = \dots = e(V_{k-1}) = \frac{m}{k^2} + \alpha$ and $e(V_k) = \frac{m}{k^2} - \frac{k-1}{k}t(m) + \frac{17k}{8} - (k-1)\alpha$. So by (7), we have

$$\begin{aligned} \sum_{i=1}^k e(V_i)^\lambda &\leq (k-1) \left(\frac{m}{k^2} + \alpha \right)^\lambda + \left(\frac{m}{k^2} - \frac{k-1}{k}t(m) + \frac{17k}{8} - (k-1)\alpha \right)^\lambda \\ &= (k-1) \left(\frac{m}{k^2} + \alpha \right)^\lambda + \left(\frac{m}{k^2} - \frac{k-1}{k}t(m) - (k-1)\alpha \right)^\lambda + O(m^{\lambda-1}) \\ &\leq (k-1) \left(\frac{m}{k^2} + \frac{k-1}{k^2}t(m) \right)^\lambda + \left(\frac{m}{k^2} - \frac{k-1}{k}t(m) - \frac{(k-1)^2}{k^2}t(m) \right)^\lambda + O(m^{\lambda-1}) \\ &= \frac{1}{k^{2\lambda-1}}m^\lambda - \frac{(k-1)\lambda}{k^{2\lambda-1}}m^{\lambda-1}t(m) + O(m^{\lambda-1}), \end{aligned}$$

where the second inequality holds because the expression in the second line is an increasing function of α , for $-\frac{k-1}{k^2}t(m) \leq \alpha \leq \frac{k-1}{k^2}t(m)$. \blacksquare

We remark that the bound in the above theorem is tight up to the term $O(m^{\lambda-1})$, by considering the complete graph K_{ks} which has $m = \binom{ks}{2}$ edges. Thus $s = (2t(m)+1)/k$. The minimum $\sum_{i=1}^k e(V_i)^\lambda$ over all k -partitions V_1, \dots, V_k of $V(K_{ks})$ is attained when $|V_i| = s$ for $i \in [k]$, and this minimum value equals

$$k \binom{s}{2}^\lambda = \frac{k}{2^\lambda} \left(\frac{2t(m)+1}{k} \right)^\lambda \left(\frac{2t(m)+1-k}{k} \right)^\lambda.$$

Using $2t(m)^2 + t(m) = m$ and $t(m) = \Theta(\sqrt{m})$, we see that

$$k \binom{s}{2}^\lambda = \frac{1}{k^{2\lambda-1}}m^\lambda - \frac{(k-1)\lambda}{k^{2\lambda-1}}m^{\lambda-1}t(m) + \Theta(m^{\lambda-1}).$$

It would be interesting to find the optimal upper bound in Theorem 4.3. We believe that the extremal graphs for ℓ_λ -norms of k -partitions (where $\lambda \geq 1$) should be the complete graphs $K_{kn+\lfloor k/2 \rfloor}$. We formulate the following question. For fixed $\lambda \geq 1$ and integer $k \geq 2$, let $s := s(m)$ be such that $m = \binom{ks+\lfloor k/2 \rfloor}{2}$ and let

$$f_{\lambda,k}(m) := \left\lfloor \frac{k}{2} \right\rfloor \binom{s+1}{2}^\lambda + \left\lceil \frac{k}{2} \right\rceil \binom{s}{2}^\lambda.$$

Question 4.4. Fix any real $\lambda \geq 1$ and integer $k \geq 2$. For any positive integer m , is it true that

$$\min_{V(G)=V_1 \cup \dots \cup V_k} \sum_{i=1}^k e(V_i)^\lambda \leq f_{\lambda,k}(m)$$

for all graphs G with m edges, with equality if and only if $m = \binom{ks + \lfloor k/2 \rfloor}{2}$ for some integer s ? Does the equality hold only for $K_{ks + \lfloor k/2 \rfloor}$ (modulo some isolated vertices)?

A result of Bollobás and Scott [4] shows that this is true for $\lambda = 1$ and any k . Theorem 1.4 provides an affirmative answer for the case $k = 2$.

We now turn to the following question of Scott [13].

Question 4.5. Does every graph G with $\binom{kn}{2}$ edges have a vertex partition into k sets, each of which contains at most $\binom{n}{2}$ edges?

We give a negative answer to this question in the case $k = 2$. For this we need to show that there exist an infinite sequence of pairs of integers with certain properties.

Lemma 4.6. There are pairs (a_i, b_i) of integers for all $i \geq 0$ such that

(i) $a_i \geq 36$ and a_i is even, and $b_i \geq 21$ and b_i is odd,

(ii) $3b_i(b_i - 1) = a_i(a_i - 1)$, and

(iii) $b_i \leq 7a_i/12$.

Proof. We recursively define integer pairs (n_i, t_i) as follows, such that the desired sequence $\{(a_i, b_i)\}_{i \geq 0}$ will be a subsequence of $\{(n_i, t_i)\}_{i \geq 0}$.

Let

$$(n_0, t_0) = (36, 21) \quad \text{and} \quad (n_1, t_1) = (133, 77)$$

and, for $i \geq 1$, let

$$n_{i+1} = 4n_i - n_{i-1} - 1 \quad \text{and} \quad t_{i+1} = 4t_i - t_{i-1} - 1. \tag{8}$$

For convenience, we write

$$\alpha_i := n_i(n_i - 1) - 3t_i(t_i - 1)$$

for $i \geq 0$, and

$$\beta_i := 2n_i n_{i-1} - n_i - n_{i-1} - 6t_i t_{i-1} + 3t_i + 3t_{i-1} + 1$$

for $i \geq 1$.

We claim that $\alpha_i = 0$ for $i \geq 0$ and that $\beta_i = 0$ for $i \geq 1$. By a direct calculation, we see that $\alpha_0 = 0$, $\alpha_1 = 0$ and $\beta_1 = 0$. Now assume for some $i \geq 1$, we have $\alpha_j = 0$ for

$j \in [i] \cup \{0\}$, and $\beta_j = 0$ for $j \in [i]$. Using (8) and the definition of α_{i+1} and β_{i+1} , we have

$$\begin{aligned}\alpha_{i+1} &= (4n_i - n_{i-1} - 1)(4n_i - n_{i-1} - 2) - 3(4t_i - t_{i-1} - 1)(4t_i - t_{i-1} - 2) \\ &= 16n_i^2 - 8n_i n_{i-1} + n_{i-1}^2 - 12n_i + 3n_{i-1} + 2 \\ &\quad - 48t_i^2 + 24t_i t_{i-1} - 3t_{i-1}^2 + 36t_i - 9t_{i-1} - 6 \\ &= 16\alpha_{i-1} + \alpha_{i-2} - 4\beta_{i-1} = 0,\end{aligned}$$

$$\begin{aligned}\beta_{i+1} &= 2(4n_i - n_{i-1} - 1)n_i - (4n_i - n_{i-1} - 1) - n_i \\ &\quad - 6(4t_i - t_{i-1} - 1)t_i + 3(4t_i - t_{i-1} - 1) + 3t_i + 1 \\ &= 8n_i^2 - 2n_{i-1}n_i - 7n_i + n_{i-1} - 24t_i^2 + 6t_{i-1}t_i + 21t_i - 3t_{i-1} - 1 \\ &= 8\alpha_{i-1} - \beta_{i-1} = 0.\end{aligned}$$

Thus, the claim follows from induction.

From (8), we see that both $\{n_i\}_{i \geq 0}$ and $\{t_i\}_{i \geq 0}$ are increasing sequences; so $n_i \geq 36$ and $t_i \geq 21$ for $i \geq 0$. Moreover, $t_i \leq 7n_i/12$ for $i \geq 0$. For otherwise, $t_i > 7n_i/12$ for some i . Then $i \geq 1$ and

$$3t_i(t_i - 1) > 3(7n_i/12)(7n_i/12 - 1) = 49n_i^2/48 - 7n_i/4,$$

which is larger than $n_i(n_i - 1)$ (since $n_i \geq 36$), a contradiction.

Using (8), it is easy to observe that n_i is even if and only if $i \equiv 0, 3 \pmod{4}$, and that t_i is odd if and only if $i \equiv 0, 1 \pmod{4}$. Therefore letting $a_i = n_{4i}$ and $b_i = t_{4i}$ for $i \geq 0$, we see that the pairs (a_i, b_i) satisfy all requirements (i), (ii) and (iii). \blacksquare

We now give the proof of Theorem 1.5.

Proof of Theorem 1.5. By Lemma 4.6, there exist infinitely many pairs $(2n, t)$ of positive integers such that t is odd, $t \leq 7n/6$, and $3t(t - 1) = 2n(2n - 1)$.

Let G be the union of three pairwise disjoint copies of the clique K_t . Then $|V(G)| = 3t$ and

$$e(G) = \frac{3t(t - 1)}{2} = n(2n - 1) = \binom{2n}{2}.$$

Let V_1, V_2 be a bipartition of $V(G)$. Without loss of generality, we assume that $|V_1| \geq |V_2|$.

Then $G[V_1]$ is the disjoint union of three cliques, say K_a, K_b , and K_c . (We set $K_0 = \emptyset$.) Hence, $G[V_2]$ is the disjoint union of cliques K_{t-a}, K_{t-b} and K_{t-c} . As t is odd, we have

$$a + b + c \geq \lceil |V(G)|/2 \rceil = (3t + 1)/2.$$

Choose integers a', b', c' such that $a' \leq a, b' \leq b, c' \leq c$ and

$$a' + b' + c' = (3t + 1)/2.$$

We also need an easy property of binomial coefficients that for any integers $m - n \geq 2$,

$$\binom{m}{2} + \binom{n}{2} > \binom{m-1}{2} + \binom{n+1}{2}. \quad (9)$$

Then we have

$$\begin{aligned}
e(V_1) &= \binom{a}{2} + \binom{b}{2} + \binom{c}{2} \\
&\geq \binom{a'}{2} + \binom{b'}{2} + \binom{c'}{2} \\
&\geq \binom{(t+1)/2}{2} + \binom{(t+1)/2}{2} + \binom{(t-1)/2}{2} \quad (\text{by (9)}) \\
&= \frac{3t^2 - 4t + 1}{8} \\
&= \frac{4n^2 - 2n - t + 1}{8} \quad (\text{as } 3t(t-1) = 2n(2n-1)) \\
&\geq \binom{n}{2} + \frac{5}{48}n \quad (\text{as } t \leq 7n/6).
\end{aligned}$$

This completes the proof of Theorem 1.5. ■

It seems likely that similar result holds for general k -partitions, though we are not able to construct such graphs due to difficulties in proving a more general version of Lemma 4.6.

References

- [1] N. Alon, Bipartite subgraphs, *Combinatorica* **16** (1996), 301–311.
- [2] A. Ban and N. Linial, Internal partitions of regular graphs, *J. Graph Theory* **83** (2016), 5–18.
- [3] B. Bollobás and A.D. Scott, Exact bounds for judicious partitions of graphs, *Combinatorica* **19** (1999), 473–486.
- [4] B. Bollobás and A.D. Scott, Better bounds for max cut, in: *Contemporary Combinatorics*, in: *Bolyai Soc. Math. Stud.*, vol. 10, 2002, pp. 185–246.
- [5] B. Bollobás and A. D. Scott, Problems and results on judicious partitions, *Random Struct. Alg.* **21** (2002), 414–430.
- [6] C.S. Edwards, Some extremal properties of bipartite subgraphs, *Canad. J. Math.* **25** (1973), 475–485.
- [7] P. Erdős, Some recent problems in Combinatorics and Graph Theory, in *Combinatorics and Computing (Proc. 26th Southeastern International Conference on Graph Theory, Boca Raton, 1995) Congressus Numerantium* 112, (1995).
- [8] R. Häggkvist, Problems, Combinatorics, North-Holland, Amsterdam, New York, 1978, p. 1203.
- [9] T. Hofmeister and H. Lefmann, On k -partite subgraphs, *Ars Combin.* **50** (1998), 303–308.
- [10] S. G. Hartke and T. Seacrest, Graphic sequences have realizations containing bisections of large degree, *J. Graph Theory* **71** (2012), 386–401.
- [11] D. J. Kleitman and D. L. Wang, Algorithms for constructing graphs and digraphs with given valences and factors, *Discrete Math* **6** (1973), 79–88.
- [12] J. Ma and X. Yu, On judicious bipartitions of graphs, *Combinatorica* **36** (2016), 537–556.
- [13] A. Scott, Judicious partitions and related problems, in: *Surveys in Combinatorics*, in: London Math. Lecture Note Ser. vol. 327, Cambridge Univ. Press, Cambridge, 2005, pp. 95–117.
- [14] B. Xu and X. Yu, Better bounds for k -partitions of graphs, *Combin. Probab. Comput.* **20** (2011), 631–640.