

COMPLEXITY OF NULL DYNAMICAL SYSTEMS AND SAUER–SHELAH LEMMAS

GUORONG GAO, JIE MA, MINGYUAN RONG, AND TUAN TRAN

ABSTRACT. The topological entropy of a topological dynamical system, introduced in a foundational paper by Adler, Konheim and McAndrew [Trans. Am. Math. Soc., 1965], is a nonnegative number that measures the uncertainty or disorder of the system. Comparing with positive entropy systems, zero entropy systems are much less understood. In order to distinguish between zero entropy systems, Huang and Ye [Adv. Math., 2009] introduced the concept of maximal pattern entropy of a topological dynamical system. At the heart of their analysis is a Sauer–Shelah type lemma. In the present paper, we provide a shorter and more conceptual proof of a strengthening of this lemma, and discuss its surprising connection between dynamical system, combinatorics and a recent breakthrough in communication complexity. We also improve one of the main results of Huang and Ye on the maximal pattern entropy of zero-dimensional systems, by proving a new Sauer–Shelah type lemma, which unifies and enhances various extremal results on VC-dimension, Natarajan dimension and Steele dimension.

1. INTRODUCTION

Ergodic theory and topological dynamics are two sister branches of the theory of dynamical systems whose origins can be traced back to Poincaré’s work on the three-body problem. The central object of study in topological dynamics is a **topological dynamical system (TDS)** (X, T) , where X is a nonempty compact metrisable space and $T: X \rightarrow X$ is a continuous map.

Ever since Kolmogorov [23] introduced entropy into ergodic theory 65 years ago, it has played a very central role in the study of dynamical systems. The spectacular success of entropy in ergodic theory led Adler, Konheim and McAndrew [1] to develop its topological counterpart in 1965. They associated to any topological dynamical system (X, T) a topological invariant $h_{\text{top}}(T) \in \mathbb{R}_+ \cup \{\infty\}$, called the **topological entropy** of (X, T) , which measures the uncertainty or disorder of the system. This concept has revolutionised topological dynamics, and is especially important for applications to classical mechanics and statistical physics (see, for example, the surveys [14, 20]).

Systems with positive topological entropy are random in certain sense, and systems with zero topological entropy are said to be **deterministic** even though they may exhibit complicated behaviours. Comparing with positive entropy systems, deterministic systems are much less understood. In order to distinguish between deterministic systems, Huang and Ye [16] introduced the concept of maximal pattern entropy $h_{\text{top}}^*(T)$ of a topological dynamical system (X, T) . It is known that both $h_{\text{top}}(T)$ and $h_{\text{top}}^*(T)$ take value in $\{0, \log 2, \log 3, \dots\} \cup \{\infty\}$, and that $h_{\text{top}}(T) > 0$ implies $h_{\text{top}}^*(T) = \infty$ (see [1, 16]). Hence maximal pattern entropy is especially useful for deterministic systems.¹

We now proceed to give more details. Let (X, T) be a TDS. Given two finite open covers \mathcal{U}, \mathcal{V} (of X), their **joint** is defined as $\mathcal{U} \vee \mathcal{V} := \{A \cap B: A \in \mathcal{U}, B \in \mathcal{V}\}$. Clearly, $\mathcal{U} \vee \mathcal{V}$ is also an open cover of size at most $|\mathcal{U}||\mathcal{V}|$, and at least 1. For a finite open cover \mathcal{U} , let $p_{X, \mathcal{U}}^*: \mathbb{N} \rightarrow \mathbb{N}$ be the function given by

$$p_{X, \mathcal{U}}^*(n) = \max_{S \subset \mathbb{N} \cup \{0\}, |S|=n} N\left(\bigvee_{i \in S} T^{-i}\mathcal{U}\right),$$

where $N(\mathcal{V})$ denotes the minimum size of a subcover chosen from an open cover \mathcal{V} . It is easy to see that the sequence $\{\log p_{X, \mathcal{U}}^*(n)\}_{n \in \mathbb{N}}$ is sub-additive, and thus by Fekete’s lemma, the limit $\lim_{n \rightarrow +\infty} \frac{1}{n} \log p_{X, \mathcal{U}}^*(n)$ exists. Denote this limit by $h_{\text{top}}^*(T, \mathcal{U})$. The **maximal pattern entropy** of (X, T)

¹We note that there are other ways to distinguish deterministic systems (cf. [11, 5]).

is then defined as

$$h_{\text{top}}^*(T) := \sup_{\mathcal{U}} h_{\text{top}}^*(T, \mathcal{U}),$$

where the supremum is taken over all finite open covers.²

We say a topological dynamical system (X, T) is **null** if $h_{\text{top}}^*(T) = 0$, that is, $h_{\text{top}}^*(T)$ attains the minimum possible value. From the definition of $h_{\text{top}}^*(T)$, we find that (X, T) is null if and only if $p_{X, \mathcal{U}}^*(n)$ grows sub-exponentially in n for each finite open cover \mathcal{U} . For such systems, an intriguing conjecture of Huang and Ye [16] further rules out the intermediate growth between polynomial and exponential.

Conjecture 1.1 ([16]). *If (X, T) is a null TDS, then $p_{X, \mathcal{U}}^*$ is of polynomial order for each finite open cover \mathcal{U} .*

The conjecture was repeated in the survey on local entropy theory by Glasner and Ye [10]. It was shown to be true for interval maps by Li [24], circle maps by Yang [30], and most interestingly for zero-dimensional systems by Huang and Ye [16]. In this paper, we study the conjecture and discuss its surprising connections between dynamical system, combinatorics, and theoretical computer science.

1.1. General systems. For general topological spaces, the following result, due to Huang and Ye [16, a special case of Theorem 4.5], is the state of the art of Conjecture 1.1.

Theorem 1.2 ([16]). *If (X, T) is a null TDS, then for each finite open cover \mathcal{U} there is a constant $c = c(\mathcal{U})$ such that $p_{X, \mathcal{U}}^*(n) \leq n^{c \log n}$ for every $n \geq 2$.*

For completeness, we provide a (simplified) proof of Theorem 1.2 in appendix. At the heart of Huang and Ye's argument is a Sauer–Shelah type lemma, namely Theorem 1.3 below. This combinatorial lemma and its predecessors [29, 26, 27, 19, 21] play a major role in all aspects of local entropy theory for topological dynamical systems. For a thorough discussion we refer the reader to the survey by Glasner and Ye [10], and Chapter 12 of the book by Kerr and Li [22].

Before stating Theorem 1.3, we borrow some terminologies from [2, 16]. We write Y^Z for the class of all functions from a set Z to a set Y . For $h \in Y^Z$ and $S \subseteq Z$, the restriction $h|_S$ of h to S is the map in Y^S defined by $i \mapsto h(i)$. Given $\mathcal{H} \subseteq Y^Z$ and $S \subseteq Z$, the restriction of \mathcal{H} to S is

$$\mathcal{H}|_S = \{h|_S : h \in \mathcal{H}\}.$$

Given an integer $r \geq 2$ and a nonempty set Z , we consider partial classes $\mathcal{H} \subseteq \{1, \dots, r, \star\}^Z$, where each $h \in \mathcal{H}$ is a partial function; specifically if $i \in Z$ is such that $h(i) = \star$ then h is undefined at i . The (non-traditional) **VC-dimension** of \mathcal{H} , denoted by $\text{VC}(\mathcal{H})$, is the maximum size of a shattered set $S \subseteq Z$, where S is **shattered** if $\mathcal{H}|_S$ contains $\{1, \dots, r\}^S$. A total class $\mathcal{F} \subseteq \{1, \dots, r\}^Z$ is called a **net** of \mathcal{H} if for every $h \in \mathcal{H}$ there exists $f \in \mathcal{F}$ such that $h(i) \in \{1, \dots, r, \star\} \setminus \{f(i)\}$ for all $i \in Z$. The smallest possible size of a net of \mathcal{H} is the **covering number** of \mathcal{H} and is denoted $C(\mathcal{H})$. We shall use $\binom{n}{\leq d}$ to denote the sum $\binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{d}$.

Using the above language, one can restate the combinatorial lemma of Huang and Ye [16, Lemma 4.1] as follows.

Theorem 1.3 ([16]). *Given $r \geq 2$ and $d \geq 1$, there exist positive constants C_r and D_r such that the following holds for $n \geq \max\{dC_r, D_r\}$. If $\mathcal{H} \subseteq \{1, \dots, r, \star\}^n$ is a partial class with $\text{VC}(\mathcal{H}) \leq d$, then*

$$C(\mathcal{H}) \leq r^2 2^m \left(\frac{n}{m}\right)^{2m},$$

where $m := \log_{r+1} \binom{n}{\leq d} + 1$.

It is worth pointing out that the proof of Theorem 1.2 only requires the case $r = 2$ of Theorem 1.3. However, the general case $r \geq 2$ was essential for various applications to dynamics, such as [16, Theorem 4.5] (of which Theorem 1.2 is a special case). For some other applications, see [21, 16].

²This definition is independent of a choice of metric.

In the present paper, we provide a shorter and more conceptual proof of Theorem 1.3. In fact, we establish the following slightly stronger³ version of Theorem 1.3; its proof is given in Section 2.

Theorem 1.4. *Let $r \geq 2$ and $n \geq d \geq 1$. If $\mathcal{H} \subseteq \{1, \dots, r, \star\}^{[n]}$ is a partial class with $\text{VC}(\mathcal{H}) \leq d$, then*

$$(1) \quad C(\mathcal{H}) \leq \binom{n}{\leq \log \frac{n}{r-1} \binom{n}{\leq d}}.$$

The case $r = 2$ of Theorem 1.4 is a result of Alon, Hanneke, Holzman and Moran [2, Theorem 12] in their study of PAC learning theory. Our proof extends their ideas.

When $r = 2$ and $d = 1$, (1) gives $C(\mathcal{H}) \leq \binom{n}{\leq \log(n+1)} \leq n^{\log(n+1)}$. More generally, one can show that the RHS of (1) is at most $n^{(r-1)d \log(n+1)}$. Note also that for $r = O(1)$ and $d = o(n)$, we have $\log \frac{n}{r-1} \binom{n}{\leq d} = o(n)$ since $\binom{n}{\leq o(n)} = 2^{o(n)}$, which implies $C(\mathcal{H}) \leq \binom{n}{\leq o(n)} = 2^{o(n)}$.

Another result of Alon, Hanneke, Holzman and Moran [2, Theorem 11] shows that (1) is nearly tight when $r = 2$ and $d = O(1)$. Interestingly, its proof hinges on a recent breakthrough in communication complexity and its implications in graph theory by Balodis, Ben-David, Göös, Jain and Kothari [3].

Theorem 1.5 ([2]). *There is a partial class $\mathcal{H} \subseteq \{1, 2, \star\}^{[n]}$ with $\text{VC}(\mathcal{H}) \leq 1$ and $C(\mathcal{H}) \geq n^{(\log n)^{1-o(1)}}$.*

To us, Theorem 1.5 strongly suggests that Conjecture 1.1 is not true in general. Since Theorem 1.5 was stated in a different language in [2], for the sake of completeness we include its proof in Section 2.2.

1.2. Zero-dimensional systems. We now revisit the zero-dimensional case of Conjecture 1.1 and link it to a recently introduced concept in theoretical computer science. Let X be a topological space. A refinement of an open cover \mathcal{U} of X is a new open cover \mathcal{V} of X such that every set in \mathcal{V} is contained in some set in \mathcal{U} . We say X is **zero-dimensional** if every finite open cover of X has a clopen (closed and open) refinement. Using the Karpovskiy–Milman Theorem [19], which is a generalisation of the Sauer–Shelah Lemma, Huang and Ye [16, Theorem 5.4] verified Conjecture 1.1 for such spaces X .

Theorem 1.6 ([16]). *Let (X, T) be a TDS and let $\mathcal{U} = \{U_1, \dots, U_r\}$ be a clopen partition of X . Then $h_{\text{top}}^*(T, \mathcal{U}) = \log \ell$ for some $\ell \in \{1, \dots, r\}$. Moreover, one of the following alternatives holds.*

- (a) *If $\ell = r$, then $p_{X, \mathcal{U}}^*(n) = r^n$ for all $n \in \mathbb{N}$.*
- (b) *If $\ell = r - 1$, then there exists $c > 0$ such that $\ell^n \leq p_{X, \mathcal{U}}^*(n) \leq n^c \ell^n$ for all $n \geq 2$.*
- (c) *If $2 \leq \ell \leq r - 2$, then there exists $c > 0$ such that $\ell^n \leq p_{X, \mathcal{U}}^*(n) \leq n^{c \log n} \ell^n$ for all $n \geq 2$.*
- (d) *If $\ell = 1$, then there exists $c > 0$ such that $p_{X, \mathcal{U}}^*(n) \leq n^c$ for all $n \geq 2$.*

Alternative (d) tells us that Conjecture 1.1 does hold when X is zero-dimensional. Indeed, let \mathcal{U} be any finite open cover of X . Then \mathcal{U} has a clopen refinement $\mathcal{V} = \{V_1, \dots, V_r\}$ which forms a partition of X . Since \mathcal{V} is a refinement of \mathcal{U} , $p_{X, \mathcal{U}}^*(n) \leq p_{X, \mathcal{V}}^*(n)$ for all n . Moreover, as (X, T) is a null TDS, $h_{\text{top}}^*(T, \mathcal{U}) = 0$, and so it follows from Theorem 1.6 (d) that $p_{X, \mathcal{V}}^*(n) \leq n^c$ for $n \geq 2$. Therefore, $p_{X, \mathcal{U}}^*(n) \leq n^c$ for all $n \geq 2$, as desired.

Huang and Ye [16] handled alternatives (b)–(d) separately. In the present paper, we improve the multiplicative factor in (c) from quasipolynomial to polynomial in n , and provide a unified treatment of (b)–(d) in the following.

Theorem 1.7. *Let (X, T) be a TDS and let $\mathcal{U} = \{U_1, \dots, U_r\}$ be a clopen partition of X . Then $h_{\text{top}}^*(T, \mathcal{U}) = \log \ell$ for some $\ell \in \{1, \dots, r\}$. Moreover, one of the following alternatives holds.*

- (a) *If $\ell = r$, then $p_{X, \mathcal{U}}^*(n) = r^n$ for all $n \in \mathbb{N}$.*
- (b) *If $\ell \leq r - 1$, then there exists $c > 0$ such that $\ell^n \leq p_{X, \mathcal{U}}^*(n) \leq n^c \ell^n$ for all $n \geq 2$.*

The key step in proving Theorem 1.7 is to find a suitable generalisation of the classical VC-dimension, and a corresponding Sauer–Shelah type lemma. As it turns out, one such generalisation

³For instance, Theorem 1.4 does not require n to be large with respect to r .

is k -Natarajan dimension, which encompasses the Natarajan dimension [25], and the Steele dimension [28]. Following Daniely, Schapira and Shahaf [8], we define the k -**Natarajan dimension** of a total class $\mathcal{H} \subseteq Y^Z$, denoted by $\dim_k(\mathcal{H})$, to be the maximum size of a subset $S \subseteq Z$ for which $\mathcal{H}|_S$ contains a subclass of the form $\prod_{i \in S} Y_i$, where Y_i is a k -element subset of Y for each $i \in S$.⁴ This recent concept has found applications in computer science and game theory [8], and in machine learning [7].

The corresponding generalisation of the Sauer–Shelah Lemma reads as follows.

Theorem 1.8. *Let $r \geq k \geq 2$ and $n \geq d \geq 1$. For any total class $\mathcal{H} \subseteq \{1, \dots, r\}^{[n]}$ with $\dim_k(\mathcal{H}) \leq d$,*

$$|\mathcal{H}| \leq (k-1)^{n-d} \sum_{i=0}^d \binom{n-i-1}{d-i} \binom{r}{k}^{d-i} r^i.$$

Remarks.

- (i) Theorem 1.8 gives $|\mathcal{H}| = O_{r,k,d}(n^d(k-1)^n)$. This bound is asymptotically tight, as shown by the total class consisting of all vectors in which at most d coordinates are greater than $k-1$. Moreover, the bound is sharp whenever $r = k \geq 2$ (see (iii) and (v) below).
- (ii) Weaker bounds were obtained by Daniely, Schapira and Shahaf [8, Theorem 1.5] and Charikar and Pabbaraju [7, Theorem 7] using a different method.
- (iii) Theorem 1.8 recovers the classic Sauer–Shelah Lemma [29, 26, 27], by taking $r = k = 2$ and noting that $\sum_{i=0}^d \binom{n-i-1}{d-i} 2^i = \sum_{i=0}^d \binom{n}{i}$.
- (iv) Specialising Theorem 1.8 to $k = 2$ yields an improved Natarajan theorem [25].
- (v) For $r = k$, $\dim_k(\mathcal{H})$ coincides with the Steele dimension [28] and Theorem 1.8 gives the same bound as in [28].

We give the proof of Theorem 1.8 in Section 3.3. We then derive Theorem 1.7 from Theorem 1.8 in Section 3.1. To the best of our knowledge, this is the first application of the k -Natarajan dimension to topological dynamics. In Section 3.2 we use Theorem 1.7 to study the complexity of symbolic dynamics (see Theorem 3.2). Finally, we close the paper with some concluding remarks in Section 4.

Notation. We use standard notation throughout. In particular, $[n]$ denotes the set $\{1, 2, \dots, n\}$. Given two nonnegative functions f and g of some underlying parameter n , we write $f = O(g)$ to mean that there are positive constants C and n_0 such that $f(n) \leq Cg(n)$ for all $n \geq n_0$, and $f = o(g)$ means that $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$. For the sake of clarity of presentation, we systematically omit floor and ceiling signs. Unless stated otherwise, all logarithms are base 2.

Acknowledgement. J. M. was supported in part by the National Key R and D Program of China 2020YFA0713100, National Natural Science Foundation of China grants 12125106, Innovation Program for Quantum Science and Technology 2021ZD0302902, and Anhui Initiative in Quantum Information Technologies grant AHY150200. T. T. was supported by the Excellent Young Talents Program (Overseas) of the National Natural Science Foundation of China.

2. A SAUER–SHELAH LEMMA FOR PARTIAL CLASSES

In Section 2.1 we give a short proof of Theorem 1.4. We then show in Section 2.2 that the bound given by Theorem 1.4 is essentially tight.

2.1. Proof of Theorem 1.4. In this section, we provide a proof of Theorem 1.4, extending ideas from [2]. We need to show that any partial class with bounded VC-dimension admits a small net. We shall construct the net via an algorithm. We first fix some notations and make some innocuous observations.

Let $\mathcal{H} \subseteq \{1, \dots, r, \star\}^Z$ be a partial class. The **shattering strength** of \mathcal{H} , denoted by $s(\mathcal{H})$, is the number of subsets $S \subseteq Z$ that are shattered by \mathcal{H} . By convention, the shattering strength of the empty

⁴This definition does not depend on a choice of Y .

class is 0, and the empty set is shattered by all nonempty classes (and so the shattering strength of any nonempty class is at least 1). It is easy to see that $s(\mathcal{H}) \leq \binom{|Z|}{\leq_{\text{VC}(\mathcal{H})}}$. For $(i, j) \in Z \times [r]$, we denote

$$\mathcal{H}_{i \rightarrow j} = \{h \in \mathcal{H} : h(i) = j\}.$$

Define the VC-minority function $M_{\mathcal{H}}: Z \rightarrow [r]$ of \mathcal{H} by letting $M_{\mathcal{H}}(i)$ be the value $j \in [r]$ which minimises $s(\mathcal{H}_{i \rightarrow j})$, with an arbitrary tie-breaking rule. Observe that for any $i \in Z$,

$$(2) \quad (r-1) \cdot s(\mathcal{H}) \geq s(\mathcal{H}_{i \rightarrow 1}) + \dots + s(\mathcal{H}_{i \rightarrow r}).$$

In particular,

$$(3) \quad s(\mathcal{H}_{i \rightarrow j}) \leq \frac{r-1}{r} \cdot s(\mathcal{H}), \text{ where } j = M_{\mathcal{H}}(i).$$

To see (2), for any subset $S \subseteq Z$ with $i \notin S$, we consider the contribution of the pair $S, S \cup \{i\}$ to both sides of the inequality. We note that every set S that is shattered by one of the classes $\mathcal{H}_{i \rightarrow 1}, \dots, \mathcal{H}_{i \rightarrow r}$ is also shattered by \mathcal{H} , and if S is shattered by all of the $\mathcal{H}_{i \rightarrow j}$ then both S and $S \cup \{i\}$ are shattered by \mathcal{H} .

We shall use the following algorithm to construct a small net of a given partial class.

The algorithm. Fix a partial class $\mathcal{H} \subseteq \{1, \dots, r, \star\}^{[n]}$. For any partial function $h \in \mathcal{H}$, the algorithm will output an index set $A^n \subseteq [n]$ and a total function $f \in \{1, \dots, r\}^{[n]}$. Set $A^0 = \emptyset$ and $\mathcal{H}^0 = \mathcal{H}$. For $i = 1, \dots, n$, do the following:

- (1) Compute the value of the VC-minority function of \mathcal{H}^{i-1} at i . Denote this value by j .
- (2) If $h(i) \neq j$, then set $f(i) = j$, $A^i = A^{i-1}$ and $\mathcal{H}^i = \mathcal{H}^{i-1}$.
- (3) If $h(i) = j$, then set $f(i) = j + 1 \pmod{r}$, $A^i = A^{i-1} \cup \{i\}$ and $\mathcal{H}^i = (\mathcal{H}^{i-1})_{i \rightarrow j}$.

The outputs of the algorithm satisfy the following properties.

Lemma 2.1.

- (a) $h \in \mathcal{H}^i$ for every $0 \leq i \leq n$. In particular, $s(\mathcal{H}^i) \geq 1$ for every $0 \leq i \leq n$.
- (b) $h(i) \neq f(i)$ for every $i \in [n]$.
- (c) f is determined by A^n .
- (d) A^n is a subset of $[n]$ of size at most $\log_{\frac{r}{r-1}} s(\mathcal{H})$.

Proof. Properties (a) and (b) are easy to verify. For the others, write $A^n = \{a_1 < a_2 < \dots < a_k\}$ and let $a_0 = 0$, $a_{k+1} = n + 1$. Given $i \in [n]$, there must exist $\ell = \ell(i) \in [k + 1]$ with $a_{\ell-1} < i \leq a_\ell$. Denote by $j = j(i)$ the value of the VC-minority function of $\mathcal{H}^{a_{\ell-1}}$ at i . A simple induction on ℓ shows that $A^i = \{a_1, \dots, a_{\ell-1}\}$, $\mathcal{H}^i = \mathcal{H}^{a_{\ell-1}}$ and $f(i) = j$ when $a_{\ell-1} < i < a_\ell$, while $A^i = \{a_1, \dots, a_\ell\}$, $\mathcal{H}^i = (\mathcal{H}^{a_{\ell-1}})_{i \rightarrow j}$ and $f(i) = j + 1 \pmod{r}$ when $i = a_\ell$. From this we see that for every $i \in [n]$, A^i , \mathcal{H}^i and $f(i)$ are uniquely determined by A^n and i . This implies (c).

From (3) and the above discussion, we get $s(\mathcal{H}^{a_\ell}) \leq \frac{r-1}{r} \cdot s(\mathcal{H}^{a_{\ell-1}})$ for every $1 \leq \ell \leq k$. Together with (a), this implies $1 \leq s(\mathcal{H}^{a_k}) \leq \left(\frac{r-1}{r}\right)^k \cdot s(\mathcal{H}^{a_0}) = \left(\frac{r-1}{r}\right)^k \cdot s(\mathcal{H})$, which results in $k \leq \log_{\frac{r}{r-1}} s(\mathcal{H})$, as desired. \square

From Lemma 2.1, we quickly deduce Theorem 1.4.

Proof of Theorem 1.4. Let $\mathcal{H} \subseteq \{1, \dots, r, \star\}^{[n]}$ be a partial class with $\text{VC}(\mathcal{H}) \leq d$. Run the algorithm for each $h \in \mathcal{H}$, and let \mathcal{F} be the total class consisting of all outputs f . By property (b) in Lemma 2.1, \mathcal{F} is a net of \mathcal{H} . Finally, from properties (c) and (d) in Lemma 2.1, we find

$$|\mathcal{F}| \leq \binom{n}{\leq \log_{\frac{r}{r-1}} s(\mathcal{H})} \leq \binom{n}{\leq \log_{\frac{r}{r-1}} \binom{n}{\leq d}},$$

where the second inequality holds since $s(\mathcal{H}) \leq \binom{n}{\leq_{\text{VC}(\mathcal{H})}} \leq \binom{n}{\leq d}$. This completes our proof. \square

2.2. Proof of Theorem 1.5. In this section we present a proof of Theorem 1.5, due to Alon, Hanneke, Holzman and Moran [2]. The proof exploits a recent breakthrough in communication complexity

and graph theory, namely Theorem 2.2 below, which provides a near-optimal solution to the Alon–Saks–Seymour problem in graph theory (for background on this problem, see the survey by Bousquet, Lagoutte and Thomassé [6]). Let $G = (V, E)$ be a simple graph. Recall that the chromatic number of G , denoted by $\chi(G)$, is the minimum k for which there exists a labelling $c: V \rightarrow [k]$ such that every edge $\{u, v\} \in E$ satisfies $c(u) \neq c(v)$. The **biclique partition number** of G , denoted $\text{bp}(G)$, is the minimum number of bicliques (i.e. complete bipartite graphs) needed to partition the edges of G . The following result follows from a recent line of breakthroughs by Göös [12]; Göös, Lovett, Meka, Watson and Zuckerman [13]; Balodis, Ben-David, Göös, Jain and Kothari [3]:

Theorem 2.2 ([3]). *For every positive integer n there exists a graph G with $\text{bp}(G) = n$ and*

$$\chi(G) \geq n^{(\log n)^{1-o(1)}},$$

where the term $o(1)$ tends to zero as n goes to infinity.

The following result allows us to use the graph G promised by Theorem 2.2 to construct a partial class \mathcal{H} with small VC-dimension and large covering number.

Lemma 2.3. *For every graph G with $\text{bp}(G) = n$, there exists a partial class $\mathcal{H} \subseteq \{1, 2, \star\}^{[n]}$ with $\text{VC}(\mathcal{H}) \leq 1$ and $C(\mathcal{H}) \geq \chi(G)$.*

We remark that our proof also gives that $|\mathcal{H}| = |V(G)|$. Before proving Lemma 2.3, let us deduce Theorem 1.5 from it.

Proof of Theorem 1.5 assuming Lemma 2.3. Let G be the graph given by Theorem 2.2. Then we have $\text{bp}(G) = n$ and $\chi(G) \geq n^{(\log n)^{1-o(1)}}$. By Lemma 2.3, there is a partial class $\mathcal{H} \subseteq \{1, 2, \star\}^{[n]}$ with $\text{VC}(\mathcal{H}) \leq 1$ and $C(\mathcal{H}) \geq \chi(G) \geq n^{(\log n)^{1-o(1)}}$, proving Theorem 1.5. \square

To complete the proof of Theorem 1.5, it remains to prove Lemma 2.3, the task we now pursue.

Proof of Lemma 2.3. Suppose we have a partition of $E(G)$ as disjoint union of $\mathcal{B}(L_i, R_i)$ for $i \in [n]$, where $\mathcal{B}(L_i, R_i)$ denotes the edge set of the complete bipartite graph with parts L_i and R_i . For each $v \in V(G)$, let h_v be a partial function in $\{1, 2, \star\}^{[n]}$ given by

$$h_v(i) = \begin{cases} 1 & \text{if } v \in L_i \\ 2 & \text{if } v \in R_i \\ \star & \text{otherwise.} \end{cases}$$

Set $\mathcal{H} = \{h_v : v \in V(G)\}$. We have to show that the partial class $\mathcal{H} \subseteq \{1, 2, \star\}^{[n]}$ satisfies $\text{VC}(\mathcal{H}) \leq 1$ and $C(\mathcal{H}) \geq \chi(G)$.

Suppose for the contrary that $\text{VC}(\mathcal{H}) \geq 2$. Then there must exist two distinct coordinates $i, j \in [n]$ such that $\{i, j\}$ is shattered by \mathcal{H} . In particular, we can find two vertices $u, v \in V(G)$ with $(h_u(i), h_u(j)) = (1, 1)$ and $(h_v(i), h_v(j)) = (2, 2)$. From the definitions of h_u and h_v , we get $u \in L_i \cap L_j$ and $v \in R_i \cap R_j$. Hence $\{u, v\}$ is covered by both $\mathcal{B}(L_i, R_i)$ and $\mathcal{B}(L_j, R_j)$, which contradicts the assumption that $\cup_{i=1}^n \mathcal{B}(L_i, R_i)$ is an edge partition of G .

It remains to show that $C(\mathcal{H}) \geq \chi(G)$. Indeed, from the definition of $C(\mathcal{H})$, there exists a total class $\mathcal{C} \subseteq \{1, 2\}^{[n]}$ that satisfies

- (i) $|\mathcal{C}| = C(\mathcal{H})$;
- (ii) for each vertex $v \in V$ there is a total function $c_v \in \mathcal{C}$ such that $h_v(i) \neq c_v(i)$ for all $i \in [n]$.

Assign to each vertex $v \in V$ the colour $c_v \in \mathcal{C}$. We claim that this is a proper colouring, and so $C(\mathcal{H}) = |\mathcal{C}| \geq \chi(G)$, as desired. Indeed, let $\{u, v\}$ be any edge in G . Since $\cup_{i=1}^n \mathcal{B}(L_i, R_i)$ is an edge partition of G , $\{u, v\} \in \mathcal{B}(L_i, R_i)$ for some $i \in [n]$. Let $u \in L_i$ and $v \in R_i$. By the definitions of h_u and h_v , we thus obtain $h_u(i) = 1$ and $h_v(i) = 2$. It then follows from (ii) that $c_u(i) = 2$ and $c_v(i) = 1$. We conclude that u and v are assigned different colours, completing our proof. \square

3. k -NATARAJAN DIMENSION AND ITS APPLICATIONS

As mentioned earlier, in this section we study the complexity function $p_{X,\mathcal{U}}^*(n)$ when \mathcal{U} is a clopen partition of X . We derive the main result of this section, namely Theorem 1.7, from Theorem 1.8. The proof of Theorem 1.8 itself is deferred to Section 3.3. In Section 3.2, we use Theorem 1.7 to study the complexity of symbolic dynamics, improving another result of Huang and Ye [16].

We begin with an auxiliary lemma that is used to prove the statements given later in the subsections.

Lemma 3.1. *Let (X, T) be a TDS and let $\mathcal{U} = \{U_1, \dots, U_r\}$ be a clopen partition of X . Then, for any subset $S \subseteq \mathbb{N} \cup \{0\}$,*

$$N\left(\bigvee_{i \in S} T^{-i}\mathcal{U}\right) = \#\left\{f \in [r]^S : \bigcap_{i \in S} T^{-i}U_{f(i)} \neq \emptyset\right\}.$$

Proof. Since \mathcal{U} is a partition of X , the sets in the open cover $\bigvee_{i \in S} T^{-i}\mathcal{U}$ are pairwise disjoint. Hence the minimum size of a subcover of $\bigvee_{i \in S} T^{-i}\mathcal{U}$ is exactly the number of nonempty sets in $\bigvee_{i \in S} T^{-i}\mathcal{U}$. Moreover, every set in $\bigvee_{i \in S} T^{-i}\mathcal{U}$ is of the form $\bigcap_{i \in S} T^{-i}U_{f(i)}$ for some function $f \in \{1, \dots, r\}^S$. Therefore, $N\left(\bigvee_{i \in S} T^{-i}\mathcal{U}\right) = \#\left\{f \in [r]^S : \bigcap_{i \in S} T^{-i}U_{f(i)} \neq \emptyset\right\}$. \square

3.1. An application in topological dynamical systems. Given a total class $\mathcal{H} \subseteq \{1, \dots, r\}^Z$ and a positive integer k , we say a subset $S \subseteq Z$ is k -Natarajan shattered by \mathcal{H} if $\mathcal{H}|_S$ contains a subclass of the form $\prod_{i \in S} Y_i$, where Y_i is a k -element subset of $\{1, \dots, r\}$ for each $i \in S$. Then, $\dim_k(\mathcal{H})$ equals the maximum size of a k -Natarajan shattered set.

Proof of Theorem 1.7 assuming Theorem 1.8. Let $\mathcal{U} = \{U_1, \dots, U_r\}$ be a clopen partition of X . For each $n \in \mathbb{N}$, let V_n be a set of n nonnegative integers such that $N\left(\bigvee_{i \in V_n} T^{-i}\mathcal{U}\right) = p_{X,\mathcal{U}}^*(n)$. Define

$$\mathcal{H}_n = \left\{h \in [r]^{V_n} : \bigcap_{i \in V_n} T^{-i}U_{h(i)} \neq \emptyset\right\}.$$

Then, by Lemma 3.1, $N\left(\bigvee_{i \in V_n} T^{-i}\mathcal{U}\right) = |\mathcal{H}_n|$, resulting in

$$p_{X,\mathcal{U}}^*(n) = |\mathcal{H}_n|.$$

Let ℓ be the maximum integer such that $\limsup_{n \rightarrow \infty} \dim_\ell(\mathcal{H}_n) = \infty$. Then we have $\ell \in \{1, \dots, r\}$. We consider the upper bound on $p_{X,\mathcal{U}}^*(n)$ first. If $\ell = r$, then clearly $p_{X,\mathcal{U}}^*(n) \leq |\mathcal{U}|^n = r^n = \ell^n$. Now suppose $\ell \leq r - 1$, then we have $\dim_{\ell+1}(\mathcal{H}_n) = O(1)$. Since $\mathcal{H}_n \subseteq \{1, \dots, r\}^{V_n}$, $|V_n| = n$ and $\dim_{\ell+1}(\mathcal{H}_n) = O(1)$, we derive from Theorem 1.8 that $|\mathcal{H}_n| \leq n^{O(1)}\ell^n$, and so

$$p_{X,\mathcal{U}}^*(n) = |\mathcal{H}_n| \leq n^{O(1)}\ell^n \text{ when } \ell \leq r - 1.$$

We proceed to lower bound $p_{X,\mathcal{U}}^*(n)$. Because $\limsup_{n \rightarrow \infty} \dim_\ell(\mathcal{H}_n) = \infty$, there exist a sequence $\{n(k)\}_{k \in \mathbb{N}}$ of positive integers and a sequence $\{W_k\}_{k \in \mathbb{N}}$ of sets satisfying

- (i) W_k is ℓ -Natarajan shattered by $\mathcal{H}_{n(k)}$.
- (ii) W_k is a subset of $V_{n(k)}$ of size k ;

For every $k \in \mathbb{N}$, we have

$$\begin{aligned} p_{X,\mathcal{U}}^*(k) &\stackrel{(ii)}{\geq} N\left(\bigvee_{i \in W_k} T^{-i}\mathcal{U}\right) = \#\left\{f \in [r]^{W_k} : \bigcap_{i \in W_k} T^{-i}U_{f(i)} \neq \emptyset\right\} \\ &\geq |\mathcal{H}_{n(k)}|_{W_k} \stackrel{(i)}{\geq} \ell^{|W_k|} \stackrel{(ii)}{=} \ell^k, \end{aligned}$$

where the first equality follows from Lemma 3.1, and in the second inequality we used the definition of $\mathcal{H}_{n(k)}$. We therefore get $p_{X,\mathcal{U}}^*(n) = \ell^n$ when $\ell = r$ and

$$\ell^n \leq p_{X,\mathcal{U}}^*(n) \leq n^{O(1)}\ell^n \text{ when } \ell \leq r - 1.$$

From this we find $h_{\text{top}}^*(T, \mathcal{U}) = \limsup_{n \rightarrow +\infty} \frac{1}{n} \log p_{X,\mathcal{U}}^*(n) = \log \ell$, where $\ell \in \{1, \dots, r\}$. This completes our proof. \square

3.2. An application in symbolic dynamics. Given an integer $r \geq 2$, we consider the product set $\Omega_r = \{1, \dots, r\}^{\mathbb{N}}$. Topology on $\{1, \dots, r\}$ is discrete, and Ω_r is endowed with the product topology. Since $\{1, \dots, r\}$ is compact and metrisable, so is the product space Ω_r . Concretely, one can equip Ω_r with the metric $d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \sum_{n \geq 1} 2^{-n} \mathbf{1}_{x_n \neq y_n}$. Let $T: \Omega_r \rightarrow \Omega_r$ be the shift $T(x_n)_{n \in \mathbb{N}} := (x_{n+1})_{n \in \mathbb{N}}$. A **subshift** is a closed T -invariant subset of Ω_r .

Consider a subshift $X \subseteq \Omega_r$. We see that (X, T) is a TDS. For $1 \leq i \leq r$, U_i denotes the clopen set $X \cap \{x \in \Omega_r: x_1 = i\}$. Then $\mathcal{U}_0 := \{U_1, \dots, U_r\}$ is a clopen partition of X . Hence we can define

$$p_X^*(n) := p_{X, \mathcal{U}_0}^*(n) \text{ for } n \in \mathbb{N}, \text{ and } h^*(X) := h_{\text{top}}^*(T, \mathcal{U}_0).$$

As a direct application of Theorem 1.7, we get the following result.

Theorem 3.2. *For any subshift (X, T) on r letters, one has $h^*(X) = \log \ell$ for some $\ell \in \{1, \dots, r\}$. Moreover, one of the following alternatives holds.*

- (a) *If $\ell = r$, then $p_X^*(n) = r^n$ for all $n \in \mathbb{N}$.*
- (b) *If $\ell \leq r - 1$, then there exists a constant $c > 0$ such that $\ell^n \leq p_X^*(n) \leq n^c \ell^n$ for all $n \geq 2$.*

We remark that for $\ell \leq r - 1$, a much weaker bound of the form $p_X^*(n) \leq n^{O(\log n)} \ell^n$ was obtained by Huang and Ye [16, Theorem 5.5].

For the rest of this section, we give a more explicit expression for $p_X^*(n)$. Huang and Ye [16] claimed, without a proof, that

$$(4) \quad p_X^*(n) = p_{X, \mathcal{U}_0}^*(n) = \max_{0 \leq s_1 < \dots < s_n} \#\{x_{1+s_1} \dots x_{1+s_n} : x \in X\}.$$

For the reader's convenience, we provide a proof of this simple fact. Let $s_1 < \dots < s_n$ be any sequence of n nonnegative integers. Since \mathcal{U}_0 is a clopen partition of X , it follows from Lemma 3.1 that

$$\begin{aligned} N\left(\bigvee_{i \in [n]} T^{-s_i} \mathcal{U}\right) &= \#\left\{f \in [r]^{[n]} : \bigcap_{i \in [n]} T^{-s_i} U_{f(i)} \neq \emptyset\right\} \\ &= \#\left\{f \in [r]^{[n]} : \text{there is } x \in X \text{ with } x \in \bigcap_{i \in [n]} T^{-s_i} U_{f(i)}\right\} \\ &= \#\left\{f \in [r]^{[n]} : \text{there is } x \in X \text{ with } T^{s_i} x \in U_{f(i)} \text{ for every } 1 \leq i \leq n\right\} \\ &= \#\left\{f \in [r]^{[n]} : \text{there is } x \in X \text{ with } x_{1+s_1} = f(1), \dots, x_{1+s_n} = f(n)\right\} \\ &= \#\{x_{1+s_1} \dots x_{1+s_n} : x \in X\}. \end{aligned}$$

Taking the maximum over all sequences $s_1 < \dots < s_n$ yields (4).

In a special case when X is the closure of the orbit of a word $a \in \Omega_r$ under the shift map T , one can further simplify (4). Indeed, for every $x \in X = \overline{\{a, Ta, T^2a, \dots\}}$ and for every $n_0 \in \mathbb{N}$, there exists $y \in \{a, Ta, T^2a, \dots\}$ such that $d(x, y) < 2^{-n_0}$. Suppose $y = T^{m-1}a$ for some $m \in \mathbb{N}$. Then the condition $d(x, y) < 2^{-n_0}$ forces $x_i = y_i = a_{m+i}$ for all $i \leq n_0$. Together with (4), this implies

$$p_X^*(n) = \max_{0 \leq s_1 < \dots < s_n} \#\{a_{m+s_1} \dots a_{m+s_n} : m \in \mathbb{N}\}.$$

The right-hand side of the equation is also called the **maximal pattern complexity** of a . Actually, this concept was introduced by Kamae and Zamboni [17, 18], and was the inspiration behind the work of Huang and Ye [16] on the maximal pattern entropy.

3.3. A lemma on k -Natarajan dimension. In this section we provide a proof of Theorem 1.8, which is inspired by [9]. Throughout the section, we view elements of product spaces as vectors. For a vector x , we denote by x_i the value of the i -th coordinate. For integers a and b , we employ the interval notation

$$[a, b] := \{x \in \mathbb{Z} : a \leq x \leq b\}.$$

The following lemma makes up the bulk of the proof of Theorem 1.8.

Lemma 3.3. Given integers r and k with $r \geq k \geq 2$, let Ω denote the alphabet

$$\{b_1, b_2, \dots, b_{k-1}\} \cup \left\{c_A : A \in \binom{[r]}{k}\right\}$$

of size $k - 1 + \binom{r}{k}$. Then for every class $\mathcal{H} \subseteq [r]^{[n]}$, there exist n maps $\varphi_1, \dots, \varphi_n$ and $n + 1$ classes $\mathcal{H}_0 := \mathcal{H}, \mathcal{H}_1, \dots, \mathcal{H}_n$ (see the diagram below) with the following properties.

(P1) For $0 \leq i \leq n$, \mathcal{H}_i is a subclass of $\Omega^{[i]} \times [r]^{[i+1, n]}$.

(P2) For $1 \leq i \leq n$, $\varphi_i : \mathcal{H}_{i-1} \rightarrow \mathcal{H}_i$ is a bijection that preserves all but the i -th coordinate.

(P3) Every element in \mathcal{H}_n has at most $\dim_k(\mathcal{H})$ coordinates in $\{c_A : A \in \binom{[r]}{k}\}$.

$$\begin{array}{ccccccc} [r]^{[n]} & & \Omega^{[1]} \times [r]^{[2, n]} & & \Omega^{[2]} \times [r]^{[3, n]} & & \dots & & \Omega^{[n]} \\ \cup & & \cup & & \cup & & & & \cup \\ \mathcal{H}_0 = \mathcal{H} & \xrightarrow{\varphi_1} & \mathcal{H}_1 & \xrightarrow{\varphi_2} & \mathcal{H}_2 & \longrightarrow & \dots & \xrightarrow{\varphi_n} & \mathcal{H}_n \end{array}$$

Assuming Lemma 3.3, we now give a proof of Theorem 1.8.

Proof of Theorem 1.8 assuming Lemma 3.3. Consider a class $\mathcal{H} \subseteq [r]^{[n]}$ with $\dim_k(\mathcal{H}) \leq d$. Applying Lemma 3.3 to \mathcal{H} , we get $\Omega, \varphi_1, \dots, \varphi_n, \mathcal{H}_1, \dots, \mathcal{H}_n$. Set $\Omega' = \{b_1, \dots, b_{k-1}\}, \Omega'' = \{c_A : A \in \binom{[r]}{k}\}$ and $\varphi = \varphi_n \circ \dots \circ \varphi_1$. By (P3), for every $x \in \mathcal{H}$, $\varphi(x)$ has at least $n - d$ coordinates in Ω' . Hence \mathcal{H} is partitioned into $d + 1$ subclasses $\mathcal{G}_0, \dots, \mathcal{G}_d$, where

$$\mathcal{G}_i = \{x \in \mathcal{H} : \varphi(x)_{n-i} \in \Omega' \text{ and } \varphi(x)|_{[n-i]} \text{ has exactly } n - d \text{ coordinates in } \Omega'\}.$$

Denote

$$\mathcal{F}_i = \{(\varphi_{n-i} \circ \dots \circ \varphi_1)(x) : x \in \mathcal{G}_i\}.$$

From (P1) and (P2), we get $\mathcal{F}_i \subseteq \Omega^{[n-i]} \times [r]^{[n-i+1, n]}$ and $|\mathcal{F}_i| = |\mathcal{G}_i|$. To bound the size of \mathcal{F}_i , let y be any vector of \mathcal{F}_i . Then $y = (\varphi_{n-i} \circ \dots \circ \varphi_1)(x)$ for some $x \in \mathcal{G}_i$. By (P2), $\varphi(x) = (\varphi_n \circ \dots \circ \varphi_{n-i+1})(y)$ and y agree in the first $n - i$ coordinates. On the other hand, as $x \in \mathcal{G}_i$, we know that $\varphi(x)_{n-i} \in \Omega'$, and that $\varphi(x)|_{[n-i]}$ has exactly $n - d$ (respectively $d - i$) coordinates in Ω' (respectively Ω''). Therefore, $y_{n-i} \in \Omega'$, and $y|_{[n-i]}$ has exactly $n - d$ (respectively $d - i$) coordinates in Ω' (respectively Ω''). From this, we obtain

$$|\mathcal{F}_i|_{[n-i]} \leq \binom{n-i-1}{n-d-1} (k-1)^{n-d} \binom{r}{k}^{d-i} = (k-1)^{n-d} \binom{n-i-1}{d-i} \binom{r}{k}^{d-i}.$$

Combining this with the trivial bound $|\mathcal{F}_i|_{[n-i+1, n]} \leq r^i$ yields

$$|\mathcal{F}_i| \leq |\mathcal{F}_i|_{[n-i]} \cdot |\mathcal{F}_i|_{[n-i+1, n]} \leq (k-1)^{n-d} \binom{n-i-1}{d-i} \binom{r}{k}^{d-i} \cdot r^i.$$

Therefore, we have

$$|\mathcal{H}| = \sum_{i=0}^d |\mathcal{G}_i| = \sum_{i=0}^d |\mathcal{F}_i| \leq (k-1)^{n-d} \sum_{i=0}^d \binom{n-i-1}{d-i} \binom{r}{k}^{d-i} r^i,$$

finishing the proof. □

Construction. We recursively define $\varphi_1, \mathcal{H}_1, \dots, \varphi_n, \mathcal{H}_n$. Suppose we have defined

$$\mathcal{H}_0, \varphi_1, \mathcal{H}_1, \dots, \varphi_{i-1}, \mathcal{H}_{i-1}$$

for some $1 \leq i \leq n$. We shall construct a map $\varphi_i : \mathcal{H}_{i-1} \rightarrow \Omega^{[i]} \times [r]^{[i+1, n]}$ and let $\mathcal{H}_i = \varphi_i(\mathcal{H}_{i-1})$. For $x \in \mathcal{H}_{i-1}$, the i -th block of \mathcal{H}_{i-1} containing x is the class

$$\partial^{(i)}(x) = \{y \in \mathcal{H}_{i-1} : x \text{ and } y \text{ agree in all but possibly the } i\text{-th coordinate}\}.$$

Note that $x \in \partial^{(i)}(x)$. Since $y \in \partial^{(i)}(x)$ if and only if $\partial^{(i)}(x) = \partial^{(i)}(y)$, \mathcal{H}_{i-1} is decomposed into i -th blocks. We thus only need to define φ_i on each i -th block. Given $x = (x_1, \dots, x_n) \in \mathcal{H}_{i-1}$, order the elements of $\partial^{(i)}(x)$ as

$$(x_1, \dots, x_{i-1}, s_1, x_{i+1}, \dots, x_n), \dots, (x_1, \dots, x_{i-1}, s_t, x_{i+1}, \dots, x_n), \text{ with } 1 \leq s_1 < \dots < s_t \leq r.$$

For $1 \leq j \leq t$, define $\varphi_i(x_1, \dots, x_{i-1}, s_j, x_{i+1}, \dots, x_n)$ to be the vector

- Except the i -th coordinate, the other coordinates of $\varphi_i(x_1, \dots, x_{i-1}, s_j, x_{i+1}, \dots, x_n)$ and x are the same;
- The i -th coordinate of $\varphi_i(x_1, \dots, x_{i-1}, s_j, x_{i+1}, \dots, x_n)$ is b_j if $1 \leq j \leq k-1$, and $c_{\{s_1, \dots, s_{k-1}, s_j\}}$ if $j \geq k$.

As these vectors $\varphi_i(x_1, \dots, x_{i-1}, s_j, x_{i+1}, \dots, x_n)$ lie in $\Omega^{[i]} \times [r]^{[i+1, n]}$, $\mathcal{H}_i = \varphi_i(\mathcal{H}_{i-1})$ is a subclass of $\Omega^{[i]} \times [r]^{[i+1, n]}$, thereby verifying (P1).

To verify the other properties, we first make some simple observations that follow readily from the definition of φ_i .

Observation 3.4. *The following hold for every $i \in [n]$ and $x \in \mathcal{H}_{i-1}$.*

- (a) $\varphi_i(x)$ preserves all but the i -th coordinate of x .
- (b) The restriction of φ_i to $\partial^{(i)}(x)$ is injective.
- (c) Suppose $x = (x_1, \dots, x_n) \in \mathcal{H}$ and the i -th coordinate of $\varphi_i(x)$ is c_A for some $A \in \binom{[r]}{k}$. Then $\mathcal{H}_{i-1} \supseteq \partial^{(i)}(x)$ contains $\{x_1\} \times \dots \times \{x_{i-1}\} \times A \times \{x_{i+1}\} \times \dots \times \{x_n\}$.

Proof of Lemma 3.3 (continuation). (P2) As $\mathcal{H}_i = \varphi_i(\mathcal{H}_{i-1})$, evidently φ_i is surjective. So what's left is to show that φ_i is injective. Suppose $\varphi_i(x) = \varphi_i(y)$. By Observation 3.4 (a), φ_i preserves all but the i -th coordinate, so x and y agree in all but possibly the i -th coordinate. Thus, x and y are two vectors of $\partial^{(i)}(x)$ with $\varphi_i(x) = \varphi_i(y)$. But by Observation 3.4 (b), the map φ_i restricted to $\partial^{(i)}(x)$ is injective, so one must have $x = y$. This proves (P2).

(P3) Let $d = \dim_k(\mathcal{H})$. Suppose for the contrary that there is a vector $y \in \mathcal{H}_n$ together with a $(d+1)$ -element subset $I \subseteq [n]$ such that for every $i \in I$, $y_i = c_{A_i}$ for some $A_i \in \binom{[r]}{k}$. By repeatedly applying Observation 3.4 (a) and (c) in a reversed ordering with respect to the coordinates, we find that $\mathcal{H}|_I$ contains $\prod_{i \in I} A_i$, implying $\dim_k(\mathcal{H}) \geq d+1$, a contradiction. This finishes the proof. \square

4. CONCLUDING REMARKS

Let (X, T) be a null topological dynamical system. In this paper we studied the maximal pattern complexity of (X, T) by making connections to extremal combinatorics and theoretical computer science. A very interesting problem which remains widely open is to find the optimal upper bound for the complexity function $p_{X, \mathcal{U}}^*: \mathbb{N} \rightarrow \mathbb{N}$. Huang and Ye conjectured that $p_{X, \mathcal{U}}^*$ is of polynomial order. In light of Theorem 1.5, we believe that one cannot improve the bound $p_{X, \mathcal{U}}^*(n) \leq n^{O(\log n)}$, and in particular, Huang and Ye's conjecture does not hold.

Question 4.1. *Does there exist a null topological dynamical system (X, T) together with a finite open cover \mathcal{U} of X such that $p_{X, \mathcal{U}}^*(n) \geq n^{c \log n}$ for every n , for some fixed constant $c > 0$?*

A positive answer to Question 4.1 would imply the existence of a sequence of graphs $\{G_n\}$ with $\text{bp}(G_n) = n$ and $\chi(G_n) \geq n^{c \log n}$ (see the proofs of Theorems 1.2 and 1.5). So a natural approach is to build such (X, T) from the infinite collection of graphs G provided by Theorem 2.2. One of the obstacles we faced is that the proof of Theorem 2.2 provides little insight on the structure of the concluded graphs G . In part, this is due to the fact that the graphs are obtained by a series of reductions, some of which are unintuitive (see [12, 13, 3]).

In all known examples of null systems, the complexity function $p_{X, \mathcal{U}}^*(n)$ grows linear with n . As a first step towards Question 4.1, the following question asks whether it is possible to construct a null system (X, T) such that for some open cover \mathcal{U} of X , the complexity function $p_{X, \mathcal{U}}^*(n)$ grows faster than linear.

Question 4.2. *Is there a null topological dynamical system (X, T) together with a finite open cover \mathcal{U} of X such that $p_{X, \mathcal{U}}^*(n) \geq n^{1+c}$ for every n , for some fixed constant $c > 0$?*

Relevant to the question is a result due to Huang and Sudakov [15, Theorem 1.2] which asserts that there exists an infinite collection of graphs G such that $\chi(G) \geq \text{bp}(G)^{6/5+o(1)}$. In contrast to the proof of Theorem 2.2, Huang and Sudakov's construction is explicit.

Another intriguing question that deserves further investigation is the tightness of the bounds in Theorems 1.3 and 1.4 for $r \geq 3$. Theorem 1.5 tells us that these bounds are essentially tight for $r = 2$. Some of the arguments in the proof of Theorem 1.5 do generalise to larger r . Let us recall some notions from hypergraph theory. Let $G = (V, E)$ be an r -graph. The r -partition number of G , denoted by $f_r(G)$, is the minimum number of complete r -partite r -graphs needed to partition the edge set of G . The chromatic number $\chi(G)$ of G is the minimum k for which there exists a colouring $c: V \rightarrow [k]$ such that every edge $e \in E$ contains two vertices u, v with $c(u) \neq c(v)$. One can easily extend Lemma 2.3 to r -graphs as follows.

Proposition 4.3. *For every r -graph G with $f_r(G) = n$, there exists a partial class $\mathcal{H} \subseteq \{1, \dots, r, \star\}^{[n]}$ with $\text{VC}(\mathcal{H}) \leq 1$ and $C(\mathcal{H}) \geq \chi(G)$.*

What is missing is an analogue of Theorem 2.2 for r -graphs.

Question 4.4 (A hypergraph Alon–Saks–Seymour problem). *Let $r \geq 3$. For every n , is there an r -graph G such that $f_r(G) = n$ and $\chi(G) \geq n^{c_r \log n}$, where $c_r > 0$ is a constant depending only on r ?*

APPENDIX A. PROOF OF THEOREM 1.2

In this section we present Huang and Ye's proof [16] of Theorem 1.2, with several simplifications. Let (X, T) be a TDS. For a finite open cover \mathcal{U} of X , define $L(\mathcal{U}) = \limsup_{n \rightarrow +\infty} \frac{\log p_{X, \mathcal{U}}^*(n)}{\log^2(n+1)}$. Then one can restate Theorem 1.2 as follows.

Theorem A.1. *If (X, T) is a null TDS, then for every finite open cover \mathcal{U}*

$$L(\mathcal{U}) = O(1).$$

Throughout the section, A^c denotes the complement $X \setminus A$ of A .

Proof of Theorem A.1. Suppose for the contrary that $L(\mathcal{U}) = \infty$. By Lemma A.3 below, there is an open cover $\mathcal{V} = \{V_1, V_2\}$ of size 2 such that $L(\mathcal{V}) = \infty$. Then, for each $d \in \mathbb{N}$, there exists a finite set $S \subset \mathbb{N} \cup \{0\}$ with $N(\bigvee_{i \in S} T^{-i}\mathcal{V}) > (|S| + 1)^{4d \log(|S|+1)}$. For each $x \in X$, let h_x be a partial function in $\{1, 2, \star\}^S$ defined as

$$h_x(i) = \begin{cases} 1 & \text{if } x \in T^{-i}(V_1^c) \\ 2 & \text{if } x \in T^{-i}(V_2^c) \\ \star & \text{otherwise.} \end{cases}$$

Since $\{V_1, V_2\}$ is a cover of X , V_1^c and V_2^c are disjoint, and hence h_x is well-defined. Consider the partial class $\mathcal{H} := \{h_x : x \in X\}$.

Claim A.2. $C(\mathcal{H}) = N(\bigvee_{i \in S} T^{-i}\mathcal{V})$.

Proof. Let $\mathcal{F} \subseteq \{1, 2\}^S$ be a total class. We can infer from the definitions of \mathcal{H} and h_x that

$$\begin{aligned} \mathcal{F} \text{ is a net of } \mathcal{H} &\iff \text{for each } h_x \in \mathcal{H} \text{ there is } f \in \mathcal{F} \text{ such that } h_x(i) \neq f(i) \text{ for all } i \in S \\ &\iff \text{for each } x \in X \text{ there is } f \in \mathcal{F} \text{ such that } x \notin T^{-i}(V_{f(i)}^c) \text{ for all } i \in S \\ &\iff \text{for each } x \in X \text{ there is } f \in \mathcal{F} \text{ such that } x \in T^{-i}V_{f(i)} \text{ for all } i \in S \\ &\iff \text{for each } x \in X \text{ there is } f \in \mathcal{F} \text{ such that } x \in \bigcap_{i \in S} T^{-i}V_{f(i)} \\ &\iff \left\{ \bigcap_{i \in S} T^{-i}V_{f(i)} : f \in \mathcal{F} \right\} \text{ is a cover of } X. \end{aligned}$$

Letting \mathcal{F} be a net of minimum size of \mathcal{H} , this yields the lower bound

$$C(\mathcal{H}) = |\mathcal{F}| \geq \left| \left\{ \bigcap_{i \in S} T^{-i} V_{f(i)} : f \in \mathcal{F} \right\} \right| \geq N \left(\bigvee_{i \in S} T^{-i} \mathcal{V} \right).$$

For the upper bound, let \mathcal{W} be a minimum subcover of $\bigvee_{i \in S} T^{-i} \mathcal{V}$. Then \mathcal{W} can be written as $\mathcal{W} = \left\{ \bigcap_{i \in S} T^{-i} V_{f(i)} : f \in \mathcal{F} \right\}$, where $\mathcal{F} \subseteq \{1, 2\}^S$ is a total class with $|\mathcal{F}| = |\mathcal{W}|$. As \mathcal{W} is a cover of X , we find that \mathcal{F} is a net of \mathcal{H} . Thus $C(\mathcal{H}) \leq |\mathcal{F}| = |\mathcal{W}| = N \left(\bigvee_{i \in S} T^{-i} \mathcal{V} \right)$. We are done. \square

From Claim A.2 we obtain $C(\mathcal{H}) = N \left(\bigvee_{i \in S} T^{-i} \mathcal{V} \right) \geq (|S| + 1)^{4d \log(|S| + 1)}$. Thus, by the remark after Theorem 1.4, there exists a size- d subset $W \subseteq S$ with $\mathcal{H}|_W \supseteq \{1, 2\}^W$. Let $x \in X$ be an element such that $h_x|_W$ lies in $\{1, 2\}^W$. Suppose x is contained in $\bigcap_{i \in W} T^{-i} V_{f(i)}$ for some $f \in \{1, 2\}^W$, then we must have $h_x(i) \neq f(i)$ for all $i \in W$, and so f is uniquely determined by $h_x|_W$. Therefore, we have

$$N \left(\bigvee_{i \in W} T^{-i} \mathcal{V} \right) \geq |\mathcal{H}|_W \cap \{1, 2\}^W = 2^{|W|}.$$

Letting $|W| = d \rightarrow \infty$ yields $h_{\text{top}}^*(T, \mathcal{V}) \geq 1 > 0$, a contradiction. \square

The rest of this section is devoted to establish the following lemma that was used in the proof Theorem A.1.

Lemma A.3. *Let (X, T) be a TDS. Suppose that \mathcal{U} is a finite open cover of X with $L(\mathcal{U}) = +\infty$. Then there is an open cover $\mathcal{V} = \{V_1, V_2\}$ of size two with $L(\mathcal{V}) = +\infty$.*

As in [16], we follow the arguments of Blanchard [4]. We shall make use of basic properties of the function L .

Proposition A.4. *Let (X, T) be a TDS. Then the following properties hold.*

- (i) **(Monotone)** *If \mathcal{V} is a refinement of \mathcal{U} , then $L(\mathcal{U}) \leq L(\mathcal{V})$.*
- (ii) **(Subadditive)** *If \mathcal{U}, \mathcal{V} are open covers of X , then $L(\mathcal{U} \vee \mathcal{V}) \leq L(\mathcal{U}) + L(\mathcal{V})$.*

Proof. For (i), consider any set S of nonnegative integers. Since \mathcal{V} is a refinement of \mathcal{U} , $\bigvee_{i \in S} T^{-i} \mathcal{V}$ is a refinement of $\bigvee_{i \in S} T^{-i} \mathcal{U}$, and so $N \left(\bigvee_{i \in S} T^{-i} \mathcal{U} \right) \leq N \left(\bigvee_{i \in S} T^{-i} \mathcal{V} \right)$. It follows that $p_{X, \mathcal{U}}^*(n) \leq p_{X, \mathcal{V}}^*(n)$ for every $n \in \mathbb{N}$, which implies $L(\mathcal{U}) \leq L(\mathcal{V})$.

For every $n \in \mathbb{N}$ we have

$$\begin{aligned} p_{X, \mathcal{U} \vee \mathcal{V}}^*(n) &= \max_S N \left(\bigvee_{i \in S} T^{-i} (\mathcal{U} \vee \mathcal{V}) \right) = \max_S N \left(\bigvee_{i \in S} (T^{-i} \mathcal{U} \vee T^{-i} \mathcal{V}) \right) \\ &\leq \max_S N \left(\bigvee_{i \in S} T^{-i} \mathcal{U} \right) \cdot N \left(\bigvee_{i \in S} T^{-i} \mathcal{V} \right) \\ &\leq \max_S N \left(\bigvee_{i \in S} T^{-i} \mathcal{U} \right) \cdot \max_T N \left(\bigvee_{i \in T} T^{-i} \mathcal{V} \right) \\ &= p_{X, \mathcal{U}}^*(n) \cdot p_{X, \mathcal{V}}^*(n), \end{aligned}$$

where the maximums are taken over all size- n subsets $S, T \subset \mathbb{N} \cup \{0\}$. Taking the logarithm and then dividing by $\log^2(n + 1)$, we obtain $L(\mathcal{U} \vee \mathcal{V}) \leq L(\mathcal{U}) + L(\mathcal{V})$. \square

Proof of Lemma A.3. Let $\mathcal{U} = \{U_1, \dots, U_k\}$. We first observe that $U_1 \neq X$. Suppose otherwise that $U_1 = X$, then for every subset $S \subseteq \mathbb{N} \cup \{0\}$ we have $X \in \bigvee_{i \in S} T^{-i} \mathcal{U}$, and so the open cover $\bigvee_{i \in S} T^{-i} \mathcal{U}$ has a subcover of size 1, which implies $L(\mathcal{U}) = 0$, a contradiction.

Given a subset $A \subseteq X$, we shall use $\text{diam}(A)$ to denote the diameter of A . We now inductively construct a sequence of closed sets $(A_n)_{n \geq 0}$ with the following three properties

- (P1) $U_1^c = A_0 \supseteq A_1 \cdots$;
- (P2) $\text{diam}(A_n) \leq 2^{-n}$ for every $n \geq 1$;
- (P3) $L(\mathcal{U}_n) = +\infty$ for every $n \geq 0$, where $\mathcal{U}_n := \{A_n^c, U_2, \dots, U_k\}$.

Clearly, $A_0 = U_1^c$ is a closed set that satisfies (P1)–(P3). Suppose that we have already constructed A_n . Since A_n is a closed subset of the compact set X , A_n is also compact, and so we can cover A_n by a finite number of closed balls B_1, \dots, B_ℓ of radius 2^{-n-2} . For $1 \leq i \leq \ell$, let $A_{n+1,i} = A_n \cap B_i$ and $\mathcal{U}_{n+1,i} = \{A_{n+1,i}^c, U_2, \dots, U_k\}$. Since $\mathcal{U}_n = \{A_n^c, U_2, \dots, U_k\}$ is an open cover of X and $A_{n+1,i}^c$ is an open set containing A_n^c , we find that $\mathcal{U}_{n+1,i}$ is also an open cover of X . The definition of $A_{n+1,i}$ gives $A_{n+1,i}^c = (A_n \cap B_i)^c = A_n^c \cup B_i^c$. Hence

$$\bigcap_{i=1}^{\ell} A_{n+1,i}^c = \bigcap_{i=1}^{\ell} (A_n^c \cup B_i^c) = A_n^c \cup (A_n \cap \bigcap_{i=1}^{\ell} B_i^c) = A_n^c \cup \emptyset = A_n^c,$$

where the third identity holds since A_n is covered by B_1, \dots, B_ℓ . It follows that every set in $\bigvee_{i=1}^{\ell} \mathcal{U}_{n+1,i}$ is contained in some set in $\mathcal{U}_n = \{A_n^c, U_2, \dots, U_k\}$. In other words, the open cover $\bigvee_{i=1}^{\ell} \mathcal{U}_{n+1,i}$ is a refinement of \mathcal{U}_n . Using Proposition A.4, we thus obtain

$$+\infty = L(\mathcal{U}_n) \leq L\left(\bigvee_{i=1}^{\ell} \mathcal{U}_{n+1,i}\right) \leq \sum_{i=1}^{\ell} L(\mathcal{U}_{n+1,i}).$$

So $L(\mathcal{U}_{n+1,i}) = +\infty$ for some $i \in [\ell]$. Set $A_{n+1} = A_{n+1,i}$. As $A_{n+1} = A_n \cap B_i$, A_{n+1} is a closed subset of A_n of diameter $\text{diam}(A_{n+1}) \leq \text{diam}(B_i) \leq 2 \cdot 2^{-n-2} = 2^{-n-1}$. Therefore, A_{n+1} has the desired properties.

From (P3) and the observation at the beginning of the proof, we see that $A_n^c \neq X$, and so $A_n \neq \emptyset$. This, together with (P1), (P3) and the compactness of X , yields $\bigcap_{n \geq 0} A_n = \{x\}$ for some $x \in X$. Because $x \in A_0 = U_1^c$ and $\mathcal{U} = \{U_1, \dots, U_k\}$ is a cover of X , there is $\ell \in \{2, \dots, k\}$ with $x \in U_\ell$. Since U_ℓ is open, there exists $\varepsilon > 0$ such that U_ℓ contains an open ball $B(x, \varepsilon)$ centered at x of radius $\varepsilon > 0$. On the other hand, since $\bigcap_{n \geq 0} A_n = \{x\}$ and $\lim_{n \rightarrow \infty} \text{diam}(A_n) = 0$, one has $A_n \subseteq B(x, \varepsilon)$ for n sufficient large. For such an n , let $V_1 = A_n^c$ and $V_2 = U_\ell$. Since $V_1 \cup V_2 \supseteq B(x, \varepsilon)^c \cup B(x, \varepsilon) = X$, $\mathcal{V} = \{V_1, V_2\}$ is a subcover of \mathcal{U}_n . Finally, we have

$$L(\mathcal{V}) \geq L(\mathcal{U}_n) = +\infty.$$

This completes our proof. □

REFERENCES

- [1] Roy L. Adler, Alan G. Konheim, M. Harry McAndrew. Topological entropy. *Transactions of the American Mathematical Society* 114.2 (1965), 309–319.
- [2] Noga Alon, Steve Hanneke, Ron Holzman, Shay Moran. A Theory of PAC Learnability of Partial Concept classes. *Proc. FOCS* (2021), 658–671.
- [3] Kaspars Balodis, Shalev Ben-David, Mika Göös, Siddhartha Jain, Robin Kothari. Unambiguous DNFs and Alon–Saks–Seymour. *Proc. FOCS* (2021), 116–124.
- [4] François Blanchard. A disjointness theorem involving topological entropy. *Bulletin de la Société Mathématique de France* 121.4 (1993), 565–578.
- [5] François Blanchard, Bernard Host, Alejandro Maass. Topological complexity. *Ergodic Theory and Dynamical Systems* 20.3 (2000), 641–662.
- [6] Nicolas Bousquet, Aurélie Lagoutte, Stéphan Thomassé. Clique versus independent set. *European Journal of Combinatorics* 40 (2014), 73–92.
- [7] Moses Charikar, Chirag Pabbaraju. A characterization of list learnability. *Proc. STOC* (2023), 1713–1726.
- [8] Amit Daniely, Michael Schapira, Gal Shahaf. Inapproximability of truthful mechanisms via generalizations of the Vapnik–Chervonenkis dimension. *SIAM Journal on Computing* 47.1 (2018), 96–120.
- [9] Zoltán Füredi, Attila Sali. Optimal multivalued shattering. *SIAM Journal on Discrete Mathematics* 26.2 (2012), 737–744.
- [10] Eli Glasner, Xiangdong Ye. Local entropy theory. *Ergodic Theory and Dynamical Systems* 29.2 (2009), 321–356.
- [11] Tim N. T. Goodman. Topological sequence entropy. *Proceedings of the London Mathematical Society* 3.2 (1974), 331–350.
- [12] Mika Göös. Lower bounds for clique vs. independent set. In *Proceedings of the 56th Symposium on Foundations of Computer Science (FOCS)*, pages 1066–1076. IEEE, 2015.

- [13] Mika Göös, Shachar Lovett, Raghu Meka, Thomas Watson, David Zuckerman. Rectangles are nonnegative juntas. *SIAM Journal on Computing* 45.5 (2016), 1835–1869.
- [14] Boris Hasselblatt, Anatole Katok. Principal structures. In *Handbook of Dynamical Systems*, North-Holland, Amsterdam, 1A (2002), 1–203.
- [15] Hao Huang, Benny Sudakov. A counterexample to the Alon-Saks-Seymour conjecture and related problems. *Combinatorica* 32 (2012), 205–219.
- [16] Wen Huang, Xiangdong Ye. Combinatorial lemmas and applications to dynamics. *Advances in Mathematics* 220.6 (2009), 1689–1716.
- [17] Teturo Kamae, Luca Zamboni. Sequence entropy and the maximal pattern complexity of infinite words. *Ergodic Theory and Dynamical Systems* 22 (2002), 1191–1199.
- [18] Teturo Kamae, Luca Zamboni. Maximal pattern complexity for discrete systems. *Ergodic Theory and Dynamical Systems* 22 (2002), 1201–1214.
- [19] Mark G. Karpovsky, Vitali D. Milman. Coordinate density of sets of vectors. *Discrete Mathematics* 24 (1978), 177–184.
- [20] Anatole Katok. Fifty years of entropy in dynamics: 1958–2007. In *Frontiers in Entropy Across the Disciplines: Panorama of Entropy: Theory, Computation, and Applications* (2023), 353–404.
- [21] David Kerr, Hanfeng Li. Independence in topological and C^* -dynamics. *Mathematische Annalen* 338.4 (2007), 869–926.
- [22] David Kerr, Hanfeng Li. *Ergodic Theory. Independence and Dichotomies*. Springer, Cham, 2016.
- [23] Andrey N. Kolmogorov. A new metric invariant of transitive dynamical systems and endomorphisms of Lebesgue spaces. *Doklady of Russian Academy of Sciences* 119 (1958), 861–864.
- [24] Jian Li. Chaos and entropy for interval maps. *Journal of Dynamics and Differential Equations* 23 (2011), 333–52.
- [25] Balas K. Natarajan. On learning sets and functions. *Machine Learning* 4 (1989), 67–97.
- [26] Norbert Sauer. On the density of families of sets. *Journal of Combinatory Theory, Series A* 13 (1972), 145–147.
- [27] Saharon Shelah. A combinatorial problem: Stability and order for models and theories in infinitary language. *Pacific Journal of Mathematics* 41 (1972), 247–261.
- [28] J. Michael Steele. Existence of submatrices with all possible columns. *Journal of Combinatorial Theory, Series A* 24 (1978), 84–88.
- [29] Vladimir Vapnik, Alexey Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Probability Theory and Its Applications* 16 (1971), 264–280.
- [30] Yini Yang. Some properties of circle maps with zero topological entropy. *Nonlinearity* 34 (2021), 2781–2799.

SCHOOL OF MATHEMATICS AND STATISTICS, FUZHOU UNIVERSITY, FUZHOU, CHINA
Email address: `grgao@fzu.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCES AND TECHNOLOGY OF CHINA, HEFEI, CHINA
Email address: `jiema@ustc.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCES AND TECHNOLOGY OF CHINA, HEFEI, CHINA
Email address: `rong_ming_yuan@mail.ustc.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCES AND TECHNOLOGY OF CHINA, HEFEI, CHINA.
Email address: `trantuan@ustc.edu.cn`