

# Counting triangles in regular graphs

Jialin He<sup>1</sup>

Xinmin Hou<sup>2,4</sup>

Jie Ma<sup>3,4</sup>

Tianying Xie<sup>5</sup>

## Abstract

In this paper, we investigate the minimum number of triangles, denoted by  $t(n, k)$ , in  $n$ -vertex  $k$ -regular graphs, where  $n$  is an odd integer and  $k$  is an even integer. The well-known Andrásfai-Erdős-Sós Theorem has established that  $t(n, k) > 0$  if  $k > \frac{2n}{5}$ . In a striking work, Lo has provided the exact value of  $t(n, k)$  for sufficiently large  $n$ , given that  $\frac{2n}{5} + \frac{12\sqrt{n}}{5} < k < \frac{n}{2}$ . Here, we bridge the gap between the aforementioned results by determining the precise value of  $t(n, k)$  in the entire range  $\frac{2n}{5} < k < \frac{n}{2}$ . This confirms a conjecture of Cambie, de Verclos, and Kang.

## 1 Introduction

We investigate the number of triangles guaranteed in regular graphs with edge density blow one half. To be precise, for positive integers  $n$  and  $k$ , we define  $t(n, k)$  to be the minimum number of triangles over all  $k$ -regular graphs on  $n$  vertices. In the case when  $n$  is even, it is easy to see that  $t(n, k) = 0$  if and only if  $k \leq \frac{n}{2}$ , which can be derived from the classic Mantel's Theorem [6] and the existence of  $n$ -vertex  $k$ -regular bipartite graphs for any  $k \leq \frac{n}{2}$ . In this paper, we mainly focus on the case when  $n$  is odd.<sup>1</sup> A special case of a celebrated theorem of Andrásfai, Erdős and Sós [1] states that every  $n$ -vertex graph with minimum degree greater than  $\frac{2n}{5}$  either is bipartite or contains some triangles. Since there exist no bipartite regular graphs with an odd number of vertices, this result implies that in the case when  $n$  is odd, we have  $t(n, k) > 0$  if  $k > \frac{2n}{5}$ .<sup>2</sup> It is natural to ask for the exact value of  $t(n, k)$  in the range  $\frac{2n}{5} < k \leq \frac{n}{2}$ . The following theorem proved by Lo [3] offers a comprehensive understanding for nearly all integers  $k$  within this range when the odd integer  $n$  is sufficiently large.

---

<sup>1</sup>Department of Mathematics, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon 999077, Hong Kong. Partially supported by Hong Kong RGC grant GRF 16308219 and Hong Kong RGC grant ECS 26304920. Email: majlhe@ust.hk

<sup>2</sup>CAS Key Laboratory of Wu Wen-Tsun Mathematics, University of Science and Technology of China, Hefei, Anhui 230026, PR China. Partially supported by the National Natural Science Foundation of China grant 12071453. Email: xmhou@ustc.edu.cn

<sup>3</sup>School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, China. Research supported by National Key Research and Development Program of China 2020YFA0713100, National Natural Science Foundation of China grant 12125106, and Anhui Initiative in Quantum Information Technologies grant AHY150200. Email: jiema@ustc.edu.cn.

<sup>4</sup>Hefei National Laboratory, University of Science and Technology of China, Hefei 230088, China. Research supported by Innovation Program for Quantum Science and Technology 2021ZD0302902.

<sup>5</sup>School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, China. Email: xiety@ustc.edu.cn

<sup>1</sup>If  $n$  is odd, it only makes sense to consider when  $k$  is even. This is because there exist no regular graphs with odd number of vertices and odd degree.

<sup>2</sup>In fact, for infinitely many odd integers  $n$ , we can state this as  $t(n, k) > 0$  if and only if  $k > \frac{2n}{5}$ .

**Theorem 1.1** (Lo, [3]). *For every odd integer  $n \geq 10^7$  and even integer  $k$  with  $\frac{2n}{5} + \frac{12\sqrt{n}}{5} < k < \frac{n}{2}$ ,*

$$t(n, k) = \frac{k}{4}(3k - n - 1).$$

*Moreover, the extremal graphs for  $t(n, k)$  must belong to the family  $\mathcal{G}(n, k)$  given by Definition 2.1.*

Very recently, Cambie, de Verclos, and Kang [2] have revisited this problem by investigating the concept of regular Turán numbers. One of their main results states that  $t(n, k) \geq \frac{n^2}{300}$  for all odd  $n$  and even  $k$  with  $k > \frac{2n}{5}$ . They also highlighted the following conjecture (see Conjecture 20 in [2]).

**Conjecture 1.2** (Cambie, de Verclos and Kang, [2]). *For every odd integer  $n$  and even integer  $k$  with  $\frac{2n}{5} < k < \frac{n}{2}$ ,  $t(n, k) = \frac{k}{4}(3k - n - 1)$ . Moreover, the extremal graphs must belong to  $\mathcal{G}(n, k)$ .*

The main contribution of this paper is to establish this conjecture for sufficiently large integers  $n$ .

**Theorem 1.3.** *For every odd integer  $n \geq 10^9$  and even integer  $k$  with  $\frac{2n}{5} < k < \frac{n}{2}$ ,  $t(n, k) = \frac{k}{4}(3k - n - 1)$ . Moreover, the extremal graphs must belong to  $\mathcal{G}(n, k)$ .*


Our proof of Theorem 1.3 employs the approach utilized in [3], incorporating several innovative technical ideas. For a brief overview of the proof, we direct readers to the initial portions of both Section 3 and Section 4. As a direct corollary of Theorem 1.3, we are able to improve the aforementioned lower bound of  $t(n, k)$  by Cambie et al. in the following.

**Corollary 1.4.** *For every odd integer  $n \geq 10^9$  and even integer  $k > \frac{2n}{5}$ , we have  $t(n, k) \geq \frac{(n+1)^2}{50}$ , where the equality holds if and only if  $k = \frac{2(n+1)}{5}$  and  $n+1$  is divisible by 5.*

The rest of the paper is organized as follows. Section 2 provides some preliminaries and introduces the graph family  $\mathcal{G}(n, k)$ . Section 3 focuses on reducing the proof of Theorem 1.3 to the key intermediate result, namely Theorem 3.1. The proof of Corollary 1.4 will also be presented at the end of Section 3. Section 4 presents the complete proof of Theorem 3.1. The concluding section offers some related remarks.

## 2 Preliminaries

We use standard notations on graphs throughout the paper. Let  $G$  be a graph. We use  $\Delta(G)$  to denote the maximum degree of  $G$ . For a subset  $U \subseteq V(G)$ , the subgraph of  $G$  induced by the vertex set  $U$  is denoted by  $G[U]$ , while the subgraph obtained from  $G$  by deleting all vertices in  $U$  is expressed by  $G \setminus U$ . Let  $N_G(U) = \bigcap_{v \in U} N_G(v)$  be the set consisting of common neighbors of all vertices of  $U$  in  $G$ . For a subset of edges  $A \subseteq E(G)$ , we denote  $G \setminus A$  the subgraph of  $G$  obtained by deleting all edges in  $A$ . Suppose that  $U, W \subset V(G)$  are disjoint. We denote  $E_G(U, W)$  to be the set of edges of  $G$  between  $U$  and  $W$  and let  $e_G(U, W) = |E_G(U, W)|$ . For a vertex  $v \in V(G)$  and a subset  $X \subseteq V(G)$ , we define  $d_X(v) = |N_G(v) \cap X|$ . For an integer  $k \geq 1$ , a  $k$ -factor of  $G$  is a spanning  $k$ -regular subgraph of  $G$ . We use  $T(G)$  to denote the number of triangles in  $G$ . For a vertex  $v \in V(G)$  and an edge  $e \in E(G)$ , we use  $T(v)$  and  $T(e)$  to denote the number of triangles containing the vertex  $v$  and the edge  $e$  respectively. For positive integers  $m$ , we write  $[m]$  for the set  $\{1, 2, \dots, m\}$ . We often drop above subscripts when they are clear from context, and omit floors and ceilings whenever they are not critical.



**Figure 1.** The construction of extremal graphs  $\mathcal{G}(n, k)$ .

Let  $n$  be an odd integer  $n$  and  $k$  be an even integer  $k$  satisfying  $\frac{2n}{5} < k < \frac{n}{2}$ . We now define an important family of  $n$ -vertex  $k$ -regular graphs  $\mathcal{G}(n, k)$  as follows (see Figure 1).

**Definition 2.1.** Let  $G$  be a complete bipartite graph with parts  $X, Y$ , each of size  $\frac{n-1}{2}$ . Let  $A \subseteq X$ ,  $B \subseteq Y$  be two subsets of size  $\frac{k}{2}$ . The family  $\mathcal{G}(n, k)$  consists of all  $n$ -vertex graphs obtained by the following procedures: adding a new vertex  $v$  to  $G$ , joining  $v$  to all vertices of  $A$  and  $B$ , removing the edges of a  $\frac{n-2k+1}{2}$ -factor from the induced subgraph  $G[A \cup B]$ , and removing the edges of a  $\frac{n-2k-1}{2}$ -factor from the induced subgraph  $G[(X \setminus A) \cup (Y \setminus B)]$ .

It is easy to see that any graph  $G \in \mathcal{G}(n, k)$  is  $k$ -regular. Moreover, all triangles in  $G$  contain the vertex  $v$ , and every neighbor of  $v$  is contained in exactly  $\frac{k}{2} - \frac{n-2k+1}{2} = \frac{3k-n-1}{2}$  triangles. Thus, the number of triangles in any graph  $G \in \mathcal{G}(n, k)$  is equal to  $\frac{k}{4}(3k - n - 1)$ .

### 3 Proofs of Theorem 1.3 and Corollary 1.4

In this section, we prove our main result – Theorem 1.3. We make use of Lo’s approach [3], where he proved that for any odd integer  $n \geq 10^7$  and even  $k$  satisfying  $\frac{2n}{5} + \frac{12\sqrt{n}}{5} < k < \frac{n}{2}$ , let  $G$  be an  $n$ -vertex  $k$ -regular graph with the minimum number of triangles, then  $G \in \mathcal{G}(n, k)$ . A key intermediate step in Lo’s proof is to show, by using Andrásfai-Erdős-Sós Theorem [1], that such  $G$  can be made bipartite by deleting  $O(k^{\frac{3}{2}})$  edges. In our proof, we present a novel insight wherein, by employing a more thorough structural analysis instead of Andrásfai-Erdős-Sós Theorem, we establish that, when the broader condition  $\frac{2n}{5} < k < \frac{n}{2}$  holds, one can also make the graph  $G$  bipartite by removing  $O(k^{\frac{3}{2}})$  edges. To be formal, we prove the following.

**Theorem 3.1.** Let  $G$  be an  $n$ -vertex  $k$ -regular graph, where  $n \geq 10^9$  is odd and  $\frac{2n}{5} < k < \frac{n}{2}$  is even. If  $T(G) \leq \frac{k}{4}(3k - n - 1)$ , then  $G$  can be made bipartite by deleting at most  $\frac{15}{2}k^{\frac{3}{2}}$  edges.

In the remaining of this section, we will demonstrate the proof of Theorem 1.3 while assuming the validity of Theorem 3.1. We also need a useful lemma proved by Lo (see Lemma 3.2 in [3]). For a graph  $H$ , let  $\phi(H)$  be the sum of squares of degrees of all vertices of  $H$ .

**Lemma 3.2** (Lo [3]). *Let  $r > 4500$  be a positive integer, and let  $H$  be an  $n$ -vertex graph with  $\Delta(H) \leq r$  and  $e(H) = \beta r$  for  $1 \leq \beta < \frac{6}{5}$ . Then  $\phi(H) \leq (\beta^2 - 2\beta + 2)r^2 + (5\beta - 4)r$ . Furthermore, equality holds if and only if  $H$  has a vertex  $u$  of degree  $r$ , a vertex  $v \in N(u)$  with degree  $(\beta - 1)r + 1$  and  $n - r - 1$  isolated vertices.*

To ensure completeness, we provide a comprehensive proof, adapted from [3], illustrating the reduction of Theorem 1.3 to Theorem 3.1, as follows.

**Proof of Theorem 1.3 (Assuming Theorem 3.1).** The proof presented here has been adjusted from Lo's work [3]. Let  $G$  be an  $n$ -vertex  $k$ -regular graph, where  $n \geq 10^9$  is odd and  $\frac{2n}{5} < k < \frac{n}{2}$  is even, such that  $T(G)$  is minimum. By the construction of  $\mathcal{G}(n, k)$ , we see that  $T(G) \leq \frac{k}{4}(3k - n - 1)$ .

First partition  $V(G) = X \cup Y$  with  $e(X, Y)$  maximal. Without loss of generality, suppose that  $|X| > |Y|$ . Define  $e(X) = \frac{1}{2}\beta k$ . By Theorem 3.1,  $G$  can be made bipartite by deleting at most  $\frac{15}{2}k^{\frac{3}{2}}$  edges. Since  $e(X) + e(Y)$  is minimum, we have that  $e(X) \leq \frac{15}{2}k^{\frac{3}{2}}$ , which implies that  $\beta \leq 15\sqrt{k}$ . Since  $2e(X) = k|X| - e(X, Y) \geq k(|X| - |Y|)$ , we get that  $\beta \geq |X| - |Y| \geq 1$ .

For any edge  $uv \in E(G[X])$ ,  $T(uv) \geq d_Y(u) + d_Y(v) - |Y| = 2k - |Y| - (d_X(u) + d_X(v))$ , where the right hand side only counts the number of triangles  $uvy$  with  $y \in Y$ . Adding up, we have

$$T(G) \geq e(X)(2k - |Y|) - \phi(G[X]) = \frac{1}{2}\beta k(2k - |Y|) - \phi(G[X]). \quad (1)$$

Using  $T(G) \leq \frac{k}{4}(3k - n - 1)$ ,  $k > \frac{2n}{5}$ ,  $|Y| \leq \frac{n-1}{2}$  and  $\beta \geq 1$ , we can get a lower bound of  $\phi(G[X])$  that (see the equation (4.3) in [3])

$$\phi(G[X]) \geq \frac{1}{8}(3\beta - 1)k^2. \quad (2)$$

In order to apply Lemma 3.2 with  $H := \phi(G[X])$ , we now proceed to show that  $\beta < \frac{6}{5}$ . Let  $S$  be the set of vertices  $x \in X$  with  $d_X(x) \geq \alpha\sqrt{\beta k}$ , where  $\alpha \leq \frac{1}{2}\sqrt{k/\beta}$  is some real to be decided later. Then,  $|S| \leq 2e(X)/\alpha\sqrt{\beta k} = \sqrt{\beta k}/\alpha$ . Also  $\Sigma := \sum_{u \in S} d_X(u) \leq e(X) + \binom{|S|}{2} \leq (\frac{1}{2} + \frac{1}{2\alpha^2})\beta k$ . By the maximality of  $e(X, Y)$ , we have  $d_X(x) \leq \frac{k}{2}$  and  $d_Y(y) \leq \frac{k}{2}$  for all  $x \in X$  and  $y \in Y$ . Hence, we have

$$\begin{aligned} \phi(G[X]) &= \sum_{u \in S} d_X(u)^2 + \sum_{v \in X \setminus S} d_X(v)^2 \leq \frac{k}{2} \cdot \Sigma + \alpha\sqrt{\beta k} \cdot (2e(X) - \Sigma) \\ &\leq \frac{1}{2} \left( \frac{1}{2} + \frac{1}{2\alpha^2} \right) \beta k^2 + \alpha \left( \frac{1}{2} - \frac{1}{2\alpha^2} \right) (\beta k)^{\frac{3}{2}} \leq \left( \frac{1}{4} + \frac{1}{4\alpha^2} \right) \beta k^2 + \frac{\alpha}{2} (\beta k)^{\frac{3}{2}}, \end{aligned}$$

where the second-to-last inequality holds since  $\alpha \leq \frac{1}{2}\sqrt{k/\beta}$  and  $\Sigma \leq (\frac{1}{2} + \frac{1}{2\alpha^2})\beta k$ . Recall  $\beta \leq 15\sqrt{k}$  and  $k > \frac{2n}{5} \geq 2^{24}$ . So we have  $(k/\beta)^{1/6} < \frac{1}{2}\sqrt{k/\beta}$ . Thus, we can set  $\alpha = (k/\beta)^{1/6}$  to get that

$$\phi(G[X]) \leq \frac{1}{4}\beta k^2 + \frac{1}{4}\beta^{\frac{4}{3}}k^{\frac{5}{3}} + \frac{1}{2}\beta^{\frac{4}{3}}k^{\frac{5}{3}} = \frac{1}{4}\beta k^2 + \frac{3}{4}\beta^{\frac{4}{3}}k^{\frac{5}{3}}.$$

Combining with the inequality (2), we get that

$$\beta - 1 - 6\beta^{\frac{4}{3}}k^{-\frac{1}{3}} \leq 0.$$

Let  $f(\beta) = \beta - 1 - 6\beta^{\frac{4}{3}}k^{-\frac{1}{3}}$ . To prove  $\beta < \frac{6}{5}$ , it is enough to show that  $f(\beta) > 0$  when  $\beta \in [\frac{6}{5}, 15\sqrt{k}]$ . Since  $f''(\beta) = -\frac{8}{3}\beta^{-\frac{2}{3}}k^{-\frac{1}{3}} < 0$ , by convexity, we only need to check the cases when  $\beta = \frac{6}{5}$  and

$\beta = 15\sqrt{k}$ . If  $\beta = \frac{6}{5}$ , we have  $f(\beta) = \frac{1}{5} - 6 \cdot (\frac{6}{5})^{\frac{4}{3}} \cdot k^{-1/3} > 0$ , as  $k > 2^{24}$ . If  $\beta = 15\sqrt{k}$ , we have  $f(\beta) = 15\sqrt{k} - 1 - 6 \cdot 15^{\frac{4}{3}} \cdot k^{\frac{1}{3}} > 0$ , as  $k > 2^{24}$ . This completes the proof of that  $\beta < \frac{6}{5}$ .

Note that  $\frac{6}{5} > \beta \geq |X| - |Y| \geq 1$ . So  $|X| - |Y|$  is an integer which must be 1. By applying Lemma 3.2 with  $H := G[X]$  and  $r := \frac{k}{2}$ , we conclude that  $\phi(G[X]) \leq \frac{1}{4}(\beta^2 - 2\beta + 2)k^2 + \frac{1}{2}(5\beta - 4)k$ . Then, by (1), we have

$$\begin{aligned} T(G) &\geq \frac{1}{2}\beta k(2k - |Y|) - \phi(G[X]) \geq \frac{1}{2}\beta k \left(2k - \frac{n-1}{2}\right) - \frac{1}{4}(\beta^2 - 2\beta + 2)k^2 - \frac{1}{2}(5\beta - 4)k \\ &= \frac{k}{4} \left(4\beta k - (\beta^2 - 2\beta + 2)k - \beta(n-1) - 2(5\beta - 4)\right) \geq \frac{k}{4}(3k - n - 1), \end{aligned} \quad (3)$$

where the last inequality holds because  $g(\beta) := 4\beta k - (\beta^2 - 2\beta + 2)k - \beta(n-1) - 2(5\beta - 4) \geq g(1) = 3k - n - 1$ , which follows from the fact  $g'(\beta) = 6k - 2\beta k - (n-1) - 10 > 0$  (as  $\beta < \frac{6}{5}$ ,  $n < \frac{5k}{2}$  and  $k > 2^{24}$ ). Thus, we get the equality in (3), from which we can easily drive that  $\beta = 1$ ,  $|Y| = \frac{n-1}{2}$ ,  $|X| = \frac{n+1}{2}$ , and  $e(G[Y]) = 0$ . Moreover, by Lemma 3.2,  $G[X]$  consists of a star centered at  $v$  with  $\frac{k}{2}$  edges and  $\frac{n-k-1}{2}$  isolated vertices. By (1), any edge  $vw$  in  $G[X]$  is contained in exactly  $2k - |Y| - (d_X(v) + d_X(w)) = 2k - \frac{n-1}{2} - (\frac{k}{2} + 1) = \frac{3k-n-1}{2}$  triangles. Taking into account all of this information, it becomes evident that  $G \in \mathcal{G}(n, k)$ , which completes the proof of Theorem 1.3. ■

To conclude this section, we give the proof of Corollary 1.4.

**Proof of Corollary 1.4.** Let  $n \geq 10^9$  be odd and  $k > \frac{2n}{5}$  be even. Note that in fact,  $k \geq \frac{2(n+1)}{5}$ . By Theorem 1.3, when  $\frac{2n}{5} < k < \frac{n}{2}$ , we have  $t(n, k) = \frac{k}{4}(3k - n - 1)$ , which is an increasing function with variable  $k$ . Thus in this range, we can get that  $t(n, k) \geq \frac{n+1}{10}(3 \cdot \frac{2n+2}{5} - n - 1) = \frac{(n+1)^2}{50}$ , where the equality holds if and only if  $k = \frac{2(n+1)}{5}$  and  $n+1$  is divisible by 5. Now consider  $k \geq \frac{n}{2}$ . Since  $n$  is odd, we see  $k \geq \frac{n+1}{2}$ . Let  $G$  be any  $n$ -vertex  $k$ -regular graph. Using the classic Moon-Moser inequality [7], we have  $T(G) \geq \frac{4e(G)}{3} \left(\frac{e(G)}{n} - \frac{n}{4}\right)$ . Since  $e(G) = \frac{kn}{2}$ , it gives that  $T(G) \geq \frac{2kn}{3} \left(\frac{k}{2} - \frac{n}{4}\right) > \frac{n^2}{3} \left(\frac{n+1}{4} - \frac{n}{4}\right) = \frac{n^2}{12} > \frac{(n+1)^2}{50}$ , which completes the proof of Corollary 1.4. ■

## 4 Proof of Theorem 3.1

This section is devoted to the proof of Theorem 3.1. Throughout this section, we assume that  $n \geq 10^9$  is odd and  $k$  is even satisfying  $\frac{2n}{5} < k < \frac{n}{2}$ , and  $G$  is an  $n$ -vertex  $k$ -regular graph with

$$T(G) \leq \frac{k}{4}(3k - n - 1). \quad (4)$$

Our goal is to show that  $G$  can be made bipartite by deleting at most  $\frac{15}{2}k^{\frac{3}{2}}$  edges. Call an edge  $e \in E(G)$  **heavy** if  $T(e) \geq \frac{3k-n-1}{3}$ . Let  $H$  be the spanning subgraph of  $G$  whose edge set consists of all heavy edges of  $G$ . Let

$$U = \{u \in V(H) : d_H(u) \geq \frac{3}{2}\sqrt{k}\} \quad \text{and} \quad G' = G \setminus E(H).$$

Building upon the approach used in [3], it is crucial to show that  $G'$  is “close” to be a bipartite graph. To make the first step, we have the following proposition proved in [3].

**Proposition 4.1.**  *$G'$  is triangle-free.*

*Proof.* It is enough to show that any triangle in  $G$  contains at least one heavy edge. Let  $T$  be any triangle in  $G$  with edges  $e_1, e_2, e_3$ . For  $i \in \{0, 1, 2, 3\}$ , let  $m_i$  denote the number of vertices in  $G$  with exact  $i$  neighbors in  $T$ . By double-counting, we have  $\sum_{i=0}^3 m_i = n$ , and  $\sum_{i=0}^3 im_i = 3k$ , which implies that  $T(e_1) + T(e_2) + T(e_3) = m_2 + 3m_3 \geq m_2 + 2m_3 = 3k - n + m_0 \geq 3k - n$ . Hence at least one of the edges of  $T$  is heavy. This proves that  $G' = G \setminus E(H)$  is triangle-free.  $\square$

The next proposition is important in our proof, which says that  $G' \setminus U$  does not contain five-cycles.

**Proposition 4.2.**  $G' \setminus U$  is  $C_5$ -free.

The remaining proof is structured into two subsections. In Subsection 4.1, we demonstrate how the proof of Proposition 4.2 leads to the derivation of Theorem 3.1. In Subsection 4.2, we complete the proof of Proposition 4.2.

#### 4.1 Reducing to Proposition 4.2

**Proof of Theorem 3.1 (Assuming Proposition 4.2).** First, by double-counting the number of pairs of edges contained in triangles, and the definition of  $H$ , we can get that

$$T(G) = \frac{1}{3} \sum_{e \in E(G)} T(e) \geq \frac{1}{3} \sum_{e \in E(H)} T(e) \geq \frac{1}{3} \cdot \frac{3k - n - 1}{3} \cdot |E(H)|.$$

Thus, by (4), we have  $|E(H)| \leq \frac{9}{4}k$ . Then, by the definition of  $U$ , we have that

$$|U| \leq \frac{2|E(H)|}{3\sqrt{k}/2} \leq \frac{9k/2}{3\sqrt{k}/2} = 3\sqrt{k}.$$

Next, we claim that  $G' \setminus U$  is bipartite. Suppose not. Then there is an odd cycle  $C_{2\ell+1}$  with minimum length  $2\ell + 1$  in  $G' \setminus U$ . By Propositions 4.1 and 4.2, we have  $\ell \geq 3$ . Let  $V(C_{2\ell+1}) = \{v_1, v_2, \dots, v_{2\ell+1}\}$ . By Proposition 4.1, we have  $N_{G'}(v_1) \cap N_{G'}(v_2) = \emptyset$ . By the minimality of  $C_{2\ell+1}$ , we can conclude that  $N_{G'}(v_1) \cap N_{G'}(v_{\ell+2}) \subseteq U$  and  $N_{G'}(v_2) \cap N_{G'}(v_{\ell+2}) \subseteq U$ . Otherwise, one can find a cycle in  $G' \setminus U$  with odd length  $t \in \{\ell + 2, \ell + 3\}$ ; since  $\ell + 3 < 2\ell + 1$  when  $\ell \geq 3$ , this contradicts our choice of  $C_{2\ell+1}$ . For any vertex  $v \in V(G') \setminus U$ ,  $d_H(v) \leq \frac{3}{2}\sqrt{k}$ , which also says that  $|N_{G'}(v)| \geq k - \frac{3}{2}\sqrt{k}$ . Since  $|U| \leq 3\sqrt{k}$ , we can get that

$$n \geq |N_{G'}(v_1) \cup N_{G'}(v_2) \cup N_{G'}(v_{\ell+2})| \geq |N_{G'}(v_1)| + |N_{G'}(v_2)| + |N_{G'}(v_{\ell+2})| - |U| \geq 3(k - \frac{3}{2}\sqrt{k}) - 3\sqrt{k},$$

which implies that  $n \geq 3k - \frac{15}{2}\sqrt{k} > \frac{5}{2}k > n$  (as  $k > \frac{2}{5}n \geq 10^8$ ). This contradiction proves the claim.

Therefore,  $G'$  can be made bipartite by deleting all edges with at least one endpoint in  $U$ , which is at most  $\binom{|U|}{2} + |U|(n - |U|) \leq n|U| - \frac{1}{2}|U|^2 \leq \frac{5}{2}k|U| - \frac{1}{2}|U|^2$ . Since  $\frac{5}{2}k|U| - \frac{1}{2}|U|^2$  is monotonically increasing with respect to  $|U|$  when  $|U| \leq 3\sqrt{k}$ ,  $G'$  can be made bipartite by deleting at most  $\frac{5}{2}k|U| - \frac{1}{2}|U|^2 \leq \frac{15}{2}k^{\frac{3}{2}} - \frac{9}{2}k$  edges. Note that  $G' = G \setminus E(H)$  and  $|E(H)| \leq \frac{9}{4}k$ . So  $G$  can be made bipartite by deleting at most  $(\frac{15}{2}k^{\frac{3}{2}} - \frac{9}{2}k) + |E(H)| < \frac{15}{2}k^{\frac{3}{2}}$  edges, completing the proof of Theorem 3.1.  $\blacksquare$

## 4.2 Proof of Proposition 4.2

We begin with an outline of the proof for Proposition 4.2. Our approach primarily relies on stability-based reasoning. To begin, we assume, for the sake of contradiction, that  $G' \setminus U$  contains a  $C_5$ . We then establish that  $G'$  closely resembles a balanced blow-up of  $C_5$ . Furthermore, by analyzing the addition of edges from  $H$  back into  $G$ , we observe an excess of triangles in  $G$ . This contradicts our initial assumption that  $T(G) \leq \frac{k}{4}(3k - n - 1)$ . Throughout the proof, all indices will be modulo 5.

Suppose on the contrary that there is a  $C_5$ , say  $u_1u_2u_3u_4u_5u_1$ , in  $G' \setminus U$ . For  $i \in [5]$ , we define  $N_i^1 = N_{G'}(u_{i-1}) \cap N_{G'}(u_{i+1})$ . By Proposition 4.1,  $G'$  is triangle-free, so  $N_{G'}(u_i) \cap N_{G'}(u_{i+1}) = \emptyset$ . Then, for  $i \in [5]$ , we can get that

$$n \geq |N_{G'}(u_{i-1}) \cup N_{G'}(u_i) \cup N_{G'}(u_{i+1})| \geq d_{G'}(u_{i-1}) + d_{G'}(u_i) + d_{G'}(u_{i+1}) - |N_i^1|,$$

which implies that

$$|N_i^1| \geq \sum_{j=i-1}^{i+1} d_{G'}(u_j) - n. \quad (5)$$

Since  $G'$  is triangle-free, for different  $i, j \in [5]$ , we also have  $N_i^1 \cap N_j^1 = \emptyset$ . Thus, by (5), we have

$$n \geq \left| \bigcup_{i=1}^5 N_i^1 \right| = \sum_{i=1}^5 |N_i^1| \geq 3 \sum_{i=1}^5 d_{G'}(u_i) - 5n, \quad (6)$$

By (5) again (summing over indices  $[5] \setminus \{i\}$ ), we get  $n - |N_i^1| \geq \left( 3 \sum_{i=1}^5 d_{G'}(u_i) - \sum_{j=i-1}^{i+1} d_{G'}(u_j) \right) - 4n$ , which implies that  $|N_i^1| \leq 5n - 3 \sum_{j=1}^5 d_{G'}(u_j) + \sum_{j=i-1}^{i+1} d_{G'}(u_j)$ . Therefore, for any  $i \in [5]$ , we have

$$\sum_{j=i-1}^{i+1} d_{G'}(u_j) - n \leq |N_i^1| \leq 5n - 3 \sum_{j=1}^5 d_{G'}(u_j) + \sum_{j=i-1}^{i+1} d_{G'}(u_j) \quad (7)$$

Note that  $d_{G'}(u) \geq k - \frac{3}{2}\sqrt{k}$  holds for any  $u \in V(G') \setminus U$ . Hence we have  $\sum_{j=1}^5 d_{G'}(u_j) \geq 5(k - \frac{3}{2}\sqrt{k})$ .

Now we choose  $v_1v_2v_3v_4v_5v_1$  to be a  $C_5$  in  $G'$  such that

$$\sum_{i=1}^5 d_{G'}(v_i) = \max \left\{ \sum_{j=1}^5 d_{G'}(w_j) : G'[\{w_1, w_2, w_3, w_4, w_5\}] \text{ contains a } C_5 \right\}.$$


Clearly the above cycle  $u_1u_2u_3u_4u_5u_1$  is also a  $C_5$  in  $G'$ . So we have

$$\sum_{i=1}^5 d_{G'}(v_i) \geq \sum_{i=1}^5 d_{G'}(u_i) \geq 5 \left( k - \frac{3}{2}\sqrt{k} \right). \quad (8)$$

We partition the vertex set  $V(G)$  using the information from the cycle  $v_1v_2v_3v_4v_5v_1$  in the following (see Figure 2). For  $i \in [5]$ , let  $a_i := d_H(v_i) = k - d_{G'}(v_i)$ ,

$$a := \sum_{i=1}^5 a_i = 5k - \sum_{i=1}^5 d_{G'}(v_i), \quad N_i := N_{G'}(v_{i-1}) \cap N_{G'}(v_{i+1}) \quad \text{and} \quad Z := V(G) \setminus (\cup_{i=1}^5 N_i).$$

Note that (5), (6) and (7) also hold for  $N_i$  and  $v_i$ . Since  $\sum_{i=1}^5 d_{G'}(v_i) = 5k - a$ , by (6), we have that



**Figure 2.** The structure of  $G'$  if  $G' \setminus U$  contains a  $C_5$ .

$$n \geq \sum_{i=1}^5 |N_i| \geq 3(5k - a) - 5n = 15k - 5n - 3a, \quad (9)$$

which implies that  $\frac{2}{5}n < k \leq \frac{2}{5}n + \frac{a}{5}$ . From now on, we define  $b := \frac{5k}{2} - n$ . Since  $k$  is even,  $b$  is an integer satisfying that  $1 \leq b \leq \frac{a}{2}$ . By (8), we also see that  $2 \leq a \leq \frac{15\sqrt{k}}{2}$ .

Next, we show that  $a$  can be bounded from above by an absolute constant. We first point out that for any  $i \in [5]$  and any vertex  $w \in N_i$ , it holds  $d_{G'}(w) \leq d_{G'}(v_i)$  (equivalently,  $d_H(w) \geq d_H(v_i) = a_i$ ). Otherwise,  $G'[\{v_1, v_2, v_3, v_4, v_5, w\} - \{v_i\}]$  also contains a  $C_5$  with  $\sum_{i=1}^5 d_{G'}(v_i) + d_{G'}(w) - d_{G'}(v_i) > \sum_{i=1}^5 d_{G'}(v_i)$ , a contradiction to the definition of  $v_1v_2v_3v_4v_5v_1$ . Also note that  $\sum_{i=1}^5 \sum_{w \in N_i} d_H(w) \leq 2|E(H)| \leq \frac{9}{2}k$ . Using (7) (for  $N_i$ ) and the facts that  $n < \frac{5}{2}k$  and  $a \leq \frac{15\sqrt{k}}{2}$ , we have that

$$\begin{aligned} \frac{9}{2}k &\geq \sum_{i=1}^5 \sum_{w \in N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i| \geq \sum_{i=1}^5 a_i \left( 3k - \sum_{j=i-1}^{i+1} a_j - n \right) \\ &\geq \frac{k}{2} \sum_{i=1}^5 a_i - \sum_{i=1}^5 \sum_{j=i-1}^{i+1} a_i a_j \geq \frac{ak}{2} - a^2 \geq \frac{ak}{2} - \frac{225}{4}k, \end{aligned}$$

which implies that  $a \leq 121$ . Therefore in the rest of the proof, it suffices to consider the cases

$$2 \leq a \leq 121. \quad (10)$$

#### 4.2.1 Refining the structure

To deal with the remaining cases, we establish more inequalities and accurate estimations to enhance the graph structure. Let  $i \in [5]$ . First, by substituting  $a$  and  $b$  and using the fact  $3k - a \leq \sum_{j=i-1}^{i+1} d_{G'}(v_j) \leq 3k$ , we can refine the inequality (7) and obtain the following new inequality

$$\frac{k}{2} - a + b = 3k - a - n \leq |N_i| \leq 5n - 3(5k - a) + 3k = \frac{k}{2} + 3a - 5b. \quad (11)$$

Next, we bound  $|Z|$  in terms of  $a$  and  $b$ . Note that  $G'$  is triangle-free. So any vertex of  $Z$  is adjacent



to at most one vertex of  $\{v_1, v_2, v_3, v_4, v_5\}$  in  $G'$ . Thus,  $|Z| \geq \sum_{i=1}^5 (d_{G'}(v_i) - |N_{i-1}| - |N_{i+1}|) = 5k - a - 2 \sum_{i=1}^5 |N_i| = 5k - a - 2(n - |Z|)$ , which implies that  $|Z| \leq a - 2b$ . On the other hand, for any  $i \in [5]$ ,  $|N_{i-1}| + |N_{i+1}| \leq d_{G'}(v_i)$ . Thus,  $\sum_{i=1}^5 |N_i| \leq (\sum_{i=1}^5 d_{G'}(v_i))/2 = (5k - a)/2$ , and  $|Z| = n - \sum_{i=1}^5 |N_i| \geq (\frac{5}{2}k - b) - \frac{5k-a}{2} = \frac{a}{2} - b$ . Summarizing, we have

$$\frac{a}{2} - b \leq |Z| \leq a - 2b. \quad (12)$$

As we discussed at the beginning of this proof, we want to demonstrate that when adding the edges of  $H$  back into  $G$ , there will be many triangles in  $G$ , thus contradicting (4). For this purpose, it is important to classify the edges  $e \in E(H)$  and estimate the value of  $T(e)$ . Set  $\alpha := \frac{1}{14}$  and let

$$A = \{v \in V(G) : d_H(v) \leq \alpha k\} \quad \text{and} \quad B = V(G) \setminus A.$$

Since  $|E(H)| \leq \frac{9}{4}k$ , we have that  $|B| \leq \frac{9k/2}{\alpha k} \leq \frac{9}{2\alpha} = 63$ . Let  $E^0 := E(H) \setminus E(H[Z])$  and it is straightforward to see that  $E^0 = E_p \cup E_q \cup E_r \cup E_s$ , where  $E_p = \{xy \in E(H) : x, y \in N_i \text{ for some } i \in [5]\}$ ,  $E_q = \{xy \in E(H) : x \in N_i \text{ and } y \in N_{i+1} \text{ for some } i \in [5]\}$ ,  $E_r = \{xy \in E(H) : x \in N_i, y \in N_j \text{ and } |i - j| = 2 \text{ for some } i, j \in [5]\}$  and  $E_s = \{xy \in E(H) : x \in Z \text{ and } y \in N_i \text{ for some } i \in [5]\}$ .

Let  $E^0 = E^1 \cup E^2 \cup E^3$ , where  $E^1 = E^0 \cap E(H[A])$ ,  $E^2 = E^0 \cap E(H[A, B])$  and  $E^3 = E^0 \cap E(H[B])$ . The next claim shows that most edges of  $E^1$  are contained in many different triangles.

**Claim 4.1.** *For  $\lambda \in \{p, q, r, s\}$ , let  $E_\lambda^1 = E_\lambda \cap E^1$ . Then, we have*

$$(1) \text{ for any edge } x_1y_1 \in E_p^1, |N_{G'}(x_1) \cap N_{G'}(y_1)| \geq \frac{6}{7}k - 840;$$

$$(2) E_q^1 = \emptyset;$$

$$(3) \text{ for any edge } x_3y_3 \in E_r^1, |N_{G'}(x_3) \cap N_{G'}(y_3)| \geq \frac{5}{14}k - 1400;$$

$$(4) \text{ for any edge } x_4y_4 \in E_s^1, |N_{G'}(x_4) \cap N_{G'}(y_4)| \geq \frac{5}{14}k - 1400.$$

*Proof.* For the first statement, let  $x_1y_1 \in E_p^1$ . We may assume that  $x_1, y_1 \in N_1$ , then the common neighbors of  $x_1$  and  $y_1$  in  $G'$  must belong to  $N_2 \cup N_5 \cup Z$ , otherwise there is a triangle in  $G'$  containing  $v_2$  or  $v_5$ . Thus, by (11), (12) and the definition of  $A$ , we have that  $|N_{G'}(x_1) \cap N_{G'}(y_1)| \geq d_{G'}(x_1) + d_{G'}(y_1) - |N_2| - |N_5| - |Z| \geq 2(1 - \alpha)k - 2(\frac{k}{2} + 3a - 5b) - (a - 2b) = (1 - 2\alpha)k - 7a + 12b \geq \frac{6}{7}k - 840$ , the last inequality holds since  $a \leq 121$  and  $b \geq 1$ .

For the second statement, suppose on the contrary that there is an edge  $x_2y_2 \in E_q^1$ . If  $x_2y_2z_2$  is a triangle in  $G$  and  $z_2 \in \bigcup_{i=1}^5 N_i$ , then  $z_2$  must belong to  $N_H(x_2) \cup N_H(y_2)$ , otherwise there exist  $w \in \{x_2, y_2\}$  and some  $i \in [5]$  such that  $v_iwz_2$  is a triangle in  $G'$ . Thus, by (12) and the definition of  $A$ , the number of triangles in  $G$  containing  $x_2y_2$  is at most  $d_H(x_2) + d_H(y_2) + |Z| \leq 2\alpha k + a - 2b \leq \frac{k}{7} + 121 < \frac{3k-n-1}{3}$ , where the last inequality holds since  $n = \frac{5}{2}k - b$  and  $k \geq 10^8$ , which contradicts that  $x_2y_2 \in E(H)$  is heavy. Hence,  $E_q^1 = \emptyset$ .

For the third statement, let  $x_3y_3 \in E_r^1$ . We may assume that  $x_3 \in N_1$  and  $y_3 \in N_3$ , then the common neighbors of  $x_3$  and  $y_3$  in  $G'$  must belong to  $N_2 \cup Z$ . Note  $N_{G'}(x_3) \subseteq N_2 \cup N_5 \cup Z$ , which implies that  $N_{G'}(x_3) \setminus N_5 \subseteq N_2 \cup Z$ . Similarly,  $N_{G'}(y_3) \setminus N_4 \subseteq N_2 \cup Z$ . Thus, by (11), (12) and the definition of  $A$ , we have that  $|N_{G'}(x_3) \cap N_{G'}(y_3)| \geq (d_{G'}(x_3) - |N_5|) + (d_{G'}(y_3) - |N_4|) - |N_2| - |Z| \geq 2(1 - \alpha)k - 3(\frac{k}{2} + 3a - 5b) - (a - 2b) = \frac{k}{2} - 2\alpha k - 10a + 17b \geq \frac{5}{14}k - 1400$ .

For the last statement, let  $x_4y_4 \in E_s^1$  with  $x_4 \in Z$ . We first claim that there is an integer  $i \in [5]$  such that  $|N_{G'}(x_4) \cap N_i|, |N_{G'}(x_4) \cap N_{i+2}| \geq \frac{3}{7}k - 410$  and  $|N_{G'}(x_4) \cap N_j| \leq 27$  for  $j \in [5] \setminus \{i, i+2\}$ . Indeed, for any vertex  $z \in Z$  and any integer  $i \in [5]$ , since  $G'$  is triangle free, there is no edge between  $N_{G'}(z) \cap N_i$  and  $N_{G'}(z) \cap N_{i+1}$  in  $G'$ . Thus, any vertex  $v \in N_{G'}(z) \cap N_i$  has degree at most  $(|N_{i-1}| - |N_{G'}(z) \cap N_{i-1}|) + (|N_{i+1}| - |N_{G'}(z) \cap N_{i+1}|) + |Z|$  in  $G'$ , which is equivalent to  $d_H(v) \geq k - (|N_{i-1}| + |N_{i+1}| + |Z| - |N_{G'}(z) \cap N_{i+1}| - |N_{G'}(z) \cap N_{i-1}|)$ . Therefore, we can conclude that for any  $z \in Z$ ,

$$|N_{G'}(z) \cap N_i| \leq \frac{2|E(H)|}{k - (|N_{i-1}| + |N_{i+1}| + |Z| - |N_{G'}(z) \cap N_{i+1}| - |N_{G'}(z) \cap N_{i-1}|)}. \quad (13)$$

Since  $d_{G'}(x_4) \geq k - \alpha k$ , without loss of generality, we may assume that  $|N_{G'}(x_4) \cap N_1| \geq (k - \alpha k - |Z|)/5 \geq (k - \alpha k - a + 2b)/5$ . By (11) and (12), since  $k \geq 10^8$  and  $a \leq 121$ , we have

$$|N_{G'}(x_4) \cap N_2| \leq \frac{9k/2}{k - 2(\frac{k}{2} + 3a - 5b) - (a - 2b) + (k - \alpha k - a + 2b)/5} = \frac{45k}{\frac{13}{7}k - 72a + 124b} \leq 27.$$

Similarly, we have  $|N_{G'}(x_4) \cap N_5| \leq 27$ . Furthermore, without loss of generality, we may assume that  $|N_{G'}(x_4) \cap N_3| \geq (k - \alpha k - |N_1| - |Z| - 54)/2 \geq (\frac{k}{2} - \alpha k - 4a + 7b - 54)/2$ . Then, by using (13) again, we have  $|N_{G'}(x_4) \cap N_4| \leq 27$ . Therefore, by (11) and (12), since  $a \leq 121$  and  $b \geq 1$ , we can get that

$$|N_{G'}(x_4) \cap N_1| \geq k - \alpha k - |N_3| - |Z| - 81 \geq \frac{k}{2} - \alpha k - 4a + 7b - 81 \geq \frac{3}{7}k - 558.$$

Similarly, we can also get that  $|N_{G'}(x_4) \cap N_3| \geq \frac{3}{7}k - 558$ . Thus, we have proved that most of the vertices of  $N_{G'}(x_4)$  are contained in  $N_1$  and  $N_3$ .

Next, we show that  $x_4$  and  $y_4$  have many common neighbors in  $G'$ . First, we claim that  $y_4 \notin (N_1 \cup N_3)$ . By symmetry, we only need to show that  $y_4 \notin N_1$ . Suppose on the contrary that  $y_4 \in N_1$ , then the third vertex of any triangle in  $G$  containing  $x_4y_4$  must belong to  $(N_G(x_4) \cap (N_2 \cup N_5 \cup Z)) \cup (N_G(y_4) \cap (N_1 \cup N_3 \cup N_4))$ . It is easy to see that  $|N_G(x_4) \cap (N_2 \cup N_5 \cup Z)| \leq d_H(x_4) + |N_{G'}(x_4) \cap N_2| + |N_{G'}(x_4) \cap N_5| + |Z| \leq \alpha k + a - 2b + 54$  and  $|N_G(y_4) \cap (N_1 \cup N_3 \cup N_4)| \leq d_H(y_4) \leq \alpha k$ . Thus the number of triangles containing edge  $x_4y_4$  is at most  $2\alpha k + a - 2b + 54 \leq \frac{k}{7} + 175 < \frac{3k-n-1}{3}$ , where the last inequality holds since  $n = \frac{5}{2}k - b$  and  $k \geq 10^8$ , which contradicts that  $x_4y_4 \in E(H)$  is heavy. If  $y_4 \in N_2$ , then  $N_{G'}(y_4) \subseteq N_1 \cup N_3 \cup Z$ . By (11) and (12),

$$\begin{aligned} |N_{G'}(x_4) \cap N_{G'}(y_4)| &\geq |N_{G'}(x_4) \cap (N_1 \cup N_3 \cup Z)| + d_{G'}(y_4) - |N_1| - |N_3| - |Z| \\ &\geq 2(\frac{3}{7}k - 558) + \frac{13}{14}k - 2(\frac{k}{2} + 3a - 5b) - (a - 2b) \geq \frac{11}{14}k - 1951 \geq \frac{5}{14}k - 1400. \end{aligned}$$

If  $y_4 \in N_4$ , then  $N_{G'}(y_4) \subseteq N_3 \cup N_5 \cup Z$ . By (11) and (12),

$$\begin{aligned} |N_{G'}(x_4) \cap N_{G'}(y_4)| &\geq |N_{G'}(x_4) \cap (N_3 \cup N_5 \cup Z)| + d_{G'}(y_4) - |N_3| - |N_5| - |Z| \\ &\geq \frac{3}{7}k - 558 + \frac{13}{14}k - 2(\frac{k}{2} + 3a - 5b) - (a - 2b) \geq \frac{5}{14}k - 1400. \end{aligned}$$

Similarly, if  $y_4 \in N_5$ , then  $N_{G'}(y_4) \subseteq N_1 \cup N_4 \cup Z$  and we can show that  $|N_{G'}(x_4) \cap N_{G'}(y_4)| \geq \frac{5}{14}k - 1400$ . This completes the proof of Claim 4.1.  $\square$

### 4.2.2 Reducing to the cases $a \in \{2, 3\}$

Having the refined structure, we now proceed to complete the proof of Proposition 4.2. By double-counting the edges of  $E(H[B]) \setminus E(H[Z])$ , we have that  $|E^2| + 2|E^3| = \sum_{i=1}^5 \sum_{w \in B \cap N_i} d_H(w) + \sum_{z \in Z \cap B} |N_H(z) \cap (\bigcup_{i=1}^5 N_i)|$ . Since  $E^0 = E(H) \setminus E(H[Z])$ , we have that

$$\begin{aligned} 2(|E^1| + |E^2| + |E^3|) &= 2|E^0| = \sum_{i=1}^5 \sum_{w \in N_i} d_H(w) + \sum_{z \in Z} \left| N_H(z) \cap \left( \bigcup_{i=1}^5 N_i \right) \right| = \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \\ &+ \sum_{i=1}^5 \sum_{w \in B \cap N_i} d_H(w) + \sum_{z \in Z} \left| N_H(z) \cap \left( \bigcup_{i=1}^5 N_i \right) \right| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) + |E^2| + 2|E^3|. \end{aligned} \quad (14)$$

Recall that any vertex  $w \in N_i$  satisfies that  $d_H(w) \geq d_H(v_i) = a_i$ . By (11), we have  $|N_i| \geq \frac{k}{2} - a$ . Combining with (14) and  $|B| \leq 63$ , we have

$$2|E^1| + |E^2| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i \cap A| \geq \sum_{i=1}^5 a_i (|N_i| - |B|) \geq \frac{ak}{2} - a^2 - 63a. \quad (15)$$

Let  $T_2$  be set of triangles containing at least one edge in  $E^2$ . Let  $M$  be the number of pairs  $(e, T)$  satisfying that  $T \in T_2$  contains  $e \in E^2$ . By the definition of  $H$ , we get  $M \geq \frac{3k-n-1}{3}|E^2|$ . Since any triangle in  $G$  contains at most two edges in  $E(H[A, B])$ , we get  $M \leq 2|T_2|$ . Hence,

$$|T_2| \geq \frac{M}{2} \geq \frac{3k-n-1}{6}|E^2| \geq \frac{k}{12}|E^2|. \quad (16)$$

Since  $G' = G \setminus E(H)$  is triangle-free, all the triangles we have considered in Claim 4.1 are disjoint from each other and disjoint from the triangles in  $T_2$ . Then, using Claim 4.1 (note that  $|E_q^1| = 0$ ), (15) and (16), we can derive that

$$\begin{aligned} T(G) &\geq \left( \frac{6}{7}k - 840 \right) |E_p^1| + \left( \frac{5}{14}k - 1400 \right) |E_r^1| + \left( \frac{5}{14}k - 1400 \right) |E_s^1| + |T_2| \\ &\geq \frac{k}{6} \left( |E_p^1| + |E_r^1| + |E_s^1| \right) + \frac{k}{12} |E^2| = \frac{k}{12} \left( 2|E^1| + |E^2| \right) \geq \frac{a}{24}k^2 - \frac{a^2 + 63a}{12}k. \end{aligned} \quad (17)$$

Suppose that  $a \geq 4$ . Then by (10), we have  $4 \leq a \leq 121$ . Since  $k \geq 10^8$ ,  $n = \frac{5}{2}k - b$  and  $b \leq \frac{a}{2}$ , it is easy to see that using (17), we have  $T(G) \geq \frac{1}{6}k^2 - \frac{67}{3}k > \frac{1}{8}k^2 + \frac{b-1}{4}k = \frac{k}{4}(3k - n - 1)$ , a contradiction to our assumption (4). Hence, we only need to examine the remaining two cases of  $a = 2$  and  $a = 3$ , which will be addressed in the subsequent subsections.

### 4.2.3 The case when $a = 3$

In this case, since  $1 \leq b \leq \frac{a}{2}$ , we get that  $b = 1$  and thus  $n = \frac{5}{2}k - 1$ . Then, by (12),  $\frac{1}{2} \leq |Z| \leq 1$ , which implies that  $|Z| = 1$ . Let  $Z = \{z\}$ . By (11), we have that  $\frac{k}{2} - 2 \leq |N_i| \leq \frac{k}{2} + 4$  for  $i \in [5]$ . We first claim that  $|B| \geq 2$ . Indeed, if  $|B| \leq 1$ , then  $|E^2| \leq k$ . By (15), we have  $2|E^1| + |E^2| \geq \frac{ak}{2} - a^2 - 63a \geq \frac{3}{2}k - 200$ ,

which implies that  $|E^1| \geq \frac{k}{4} - 100$ . Then, since  $k \geq 10^8$ , Claim 4.1 implies that

$$\begin{aligned} T(G) &\geq \left(\frac{6}{7}k - 840\right) |E_p^1| + \left(\frac{5}{14}k - 1400\right) |E_r^1| + \left(\frac{5}{14}k - 1400\right) |E_s^1| + \frac{k}{12} |E^2| \\ &\geq \left(\frac{5}{14}k - 1400\right) |E^1| + \frac{k}{12} |E^2| \geq \left(\frac{5}{14}k - 1400\right) \left(\frac{k}{4} - 100\right) + \frac{k^2}{12} > \frac{1}{8}k^2 = \frac{k(3k - n - 1)}{4}, \end{aligned}$$

which is a contradiction to our assumption (4). Thus, we may assume that  $|B| \geq 2$ . Then, since  $|Z| = 1$ , we can get that  $|B \cap N_i| \geq 1$  for some integer  $i \in [5]$ . Without loss of generality, we may assume that there is a vertex  $w_1 \in B \cap N_1$ .

Next, we show that  $|N_H(w_1) \cap (N_1 \cup N_3 \cup N_4)| \geq \frac{k}{28}$ . Indeed, if  $|N_H(w_1) \cap (N_1 \cup N_3 \cup N_4)| < \frac{k}{28}$ , then  $|N_H(w_1) \cap (N_2 \cup N_5)| \geq d_H(w_1) - \frac{k}{28} - 1 \geq \frac{k}{28} - 1 > |B|$ . We may choose a vertex  $w_2 \in N_H(w_1) \cap N_2 \cap A$ . Then, since  $k \geq 10^8$ , the number of triangles in  $G$  containing  $w_1 w_2$  is at most

$$|N_H(w_1) \cap (N_1 \cup N_3 \cup N_4)| + d_H(w_2) + |Z| < \frac{k}{28} + \frac{k}{14} + 1 < \frac{k}{6} = \frac{3k - n - 1}{3},$$

which is a contradiction to  $w_1 w_2 \in E(H)$  is heavy.

We denote the set of vertex in  $(N_2 \cup N_5) \cap A$  that are not adjacent to  $w_1$  in  $G$  by  $S$ . In the following, we prove that  $|S|$  is big and the degree of most vertices of  $S$  in  $H$  is larger than  $v_2$  or  $v_5$ . In this way, we can improve the bound of (15) and get a contradiction.

Since  $|N_H(w_1) \cap (N_1 \cup N_3 \cup N_4)| \geq \frac{k}{28}$ ,  $|N_G(w_1) \cap (N_2 \cup N_5)| = |N_{G'}(w_1) \cap (N_2 \cup N_5)| + |N_H(w_1) \cap (N_2 \cup N_5)| \leq k - |N_H(w_1) \cap (N_1 \cup N_3 \cup N_4)| \leq k - \frac{k}{28}$ . Thus, we get that

$$|S| \geq |N_2| + |N_5| - |N_G(w_1) \cap (N_2 \cup N_5)| - |B| \geq 2\left(\frac{k}{2} - 2\right) - k + \frac{k}{28} - 63 \geq \frac{k}{28} - 67. \quad (18)$$

Next, we analysis the connection between these vertices and  $z$ . For  $i \in [5]$ , if all the vertices  $v_i$  are not contained in  $N_{G'}(z)$ , then similar to (5), we can get  $|N_i| \geq d_{G'}(v_{i-1}) + d_{G'}(v_i) + d_{G'}(v_{i+1}) - n + 1$ , which implies that  $n - 1 = \sum_{i=1}^5 |N_i| \geq 3 \sum_{i=1}^5 d_{G'}(v_i) - 5n + 5 = 3(5k - 3) - 5n + 5$ , and thus we have  $n \geq \frac{5}{2}k - \frac{1}{2}$ , which is a contradiction to  $n = \frac{5}{2}k - 1$ . Thus, we may assume that  $z v_j \in E(G')$  for some  $j \in [5]$ . According to our definition of  $N_i$ , for  $i \neq j$ ,  $z v_i \notin E(G')$ . Then, similar to (6), we can get that  $n - 1 = \sum_{i=1}^5 |N_i| \geq 3 \sum_{i=1}^5 d_{G'}(v_i) - 5n + 2 = 3(5k - 3) - 5n + 2$ , and thus we have  $n \geq \frac{5}{2}k - 1$ , which implies that inequality holds. Therefore, we can get  $N_{G'}(v_j) = N_{j-1} \cup N_{j+1} \cup \{z\}$  and for any  $w \in N_j$ ,  $N_{G'}(w) \subseteq N_{j-1} \cup N_{j+1} \cup \{z\}$ . For  $i \in [5]$ ,  $i \neq j$ ,  $N_{G'}(v_i) = N_{i-1} \cup N_{i+1}$  and for any  $w \in N_i$ ,  $N_{G'}(w) \subseteq N_{i-1} \cup N_{i+1} \cup \{z\}$ .

First note that  $N_{G'}(z) \cap (N_{j-1} \cup N_{j+1}) = \emptyset$ , or we will get a triangle in  $G'$ . Let  $Z_j = N_j \setminus (N_{G'}(z) \cup B)$ ,  $Z_{j-2} = (N_{G'}(z) \cap N_{j-2}) \setminus B$  and  $Z_{j+2} = (N_{G'}(z) \cap N_{j+2}) \setminus B$ . Now, we claim that  $|Z_j|, |Z_{j-2}|, |Z_{j+2}| \leq \frac{k}{112}$ . Indeed, if  $|Z_j| > \frac{k}{112}$ , then, since  $z v_j \in E(G')$ , comparing the vertex of  $Z_j$  with  $v_j$ , we can get  $\sum_{i=1}^5 \sum_{w \in A \cap N_i} d_{G'}(w) \leq \sum_{i=1}^5 (k - a_i) |N_i \cap A| - \frac{k}{112}$ . By (15), this implies that

$$2|E^1| + |E^2| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i \cap A| + \frac{k}{112} \geq 3\left(\frac{k}{2} - 65\right) + \frac{k}{112} = \frac{169}{112}k - 200 > \frac{3}{2}k,$$

and by (17), since  $k \geq 10^8$ , we can get the number of triangles in  $G$  is at least,

$$T(G) \geq \frac{k}{12} (2|E^1| + |E^2|) \geq \frac{k}{12} \cdot \frac{3}{2}k = \frac{1}{8}k^2 = \frac{k(3k - n - 1)}{4},$$

which is a contradiction to our assumption (4) shows that  $|Z_j| \leq \frac{k}{112}$ . We denote the property that  $2|E^1| + |E^2| > \frac{3}{2}k$  by  $(\star)$ . By the analysis above, when we get  $(\star)$ , we are done. The proof of the remaining two cases are similar to the above. If  $|Z_{j-2}| > \frac{k}{112}$  and  $|Z_{j+2}| = 0$ , then we can get that  $\sum_{i=1}^5 \sum_{w \in A \cap N_i} d_{G'}(w) \leq \sum_{i=1}^5 (k - a_i) |N_i \cap A| + |Z_{j-2}| - 2|Z_{j-2}| \leq \sum_{i=1}^5 (k - a_i) |N_i \cap A| - \frac{k}{112}$ . If  $|Z_{j-2}| > \frac{k}{112}$  and  $|Z_{j+2}| \neq 0$ , then we can get that  $\sum_{i=1}^5 \sum_{w \in A \cap N_i} d_{G'}(w) \leq \sum_{i=1}^5 (k - a_i) |N_i \cap A| + |Z_{j-2}| + |Z_{j+2}| - 2|Z_{j-2}||Z_{j+2}| \leq \sum_{i=1}^5 (k - a_i) |N_i \cap A| - \frac{k}{112} + 1$ . Doing the above process again, we will get a contradiction. Therefore, we have  $|Z_{j-2}|, |Z_{j+2}| \leq \frac{k}{112}$ .

Now, recall that  $w_1 \in B \cap N_1$ ,  $S$  is the set of vertex in  $(N_2 \cup N_5) \cap A$  that are not adjacent to  $w_1$  in  $G$  and  $zv_j \in E(G')$  for some  $j \in [5]$ . By symmetry, it suffices to consider  $j \in \{1, 2, 3\}$ . If  $j = 1$ , since all the vertices of  $S$  are not adjacent to  $w_1$  in  $G$ , comparing these vertices with  $v_2$  or  $v_5$ , by (18), we can get that

$$2|E^1| + |E^2| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i \cap A| + |S| > 3\left(\frac{k}{2} - 65\right) + \frac{k}{112} > \frac{3}{2}k,$$

again we get  $(\star)$ . If  $j = 2$ , let  $S' = S \setminus (Z_2 \cup Z_5)$ . Since  $|Z_2|, |Z_5| \leq \frac{k}{112}$ , we have  $|S'| \geq |S| - 2 \cdot \frac{k}{112} \geq \frac{k}{56} - 67$ . And for  $v \in S' \cap N_2$ ,  $v$  is not adjacent to  $z$ . For  $v \in S' \cap N_5$ ,  $v$  is not adjacent to  $z$  and  $w_1$ . Comparing these vertices with  $v_2$  or  $v_5$ , we can get that

$$2|E^1| + |E^2| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i \cap A| + |S'| > 3\left(\frac{k}{2} - 65\right) + \frac{k}{112} > \frac{3}{2}k,$$

again we get  $(\star)$ . If  $j = 3$ , since  $G'$  is triangle-free and  $N_{G'}(v_3) = N_2 \cup N_4 \cup \{z\}$ ,  $N_{G'}(z) \cap N_2 = N_{G'}(z) \cap N_4 = \emptyset$ . Let  $S'' = S \setminus Z_5$ , since  $|Z_5| \leq \frac{k}{112}$ , we have  $|S''| \geq |S| - \frac{k}{112} \geq \frac{3}{112}k - 67$ . And for any vertex  $v \in S''$ ,  $v$  is not adjacent to  $z$  and  $w_1$ . Comparing these vertices with  $v_2$  or  $v_5$ , we can get that

$$2|E^1| + |E^2| \geq \sum_{i=1}^5 \sum_{w \in A \cap N_i} d_H(w) \geq \sum_{i=1}^5 a_i |N_i \cap A| + |S''| > 3\left(\frac{k}{2} - 65\right) + \frac{k}{112} > \frac{3}{2}k,$$

again we get  $(\star)$ . Therefore, we have completed the proof when  $a = 3$ .

#### 4.2.4 The case when $a = 2$

In this case, since  $1 \leq b \leq \frac{a}{2}$ , we get that  $b = 1$  and thus  $n = \frac{5}{2}k - 1$ . By (12), we have  $|Z| = 0$ . Then,  $V(G) = \bigcup_{i=1}^5 N_i$ . By (11), we have that  $\frac{k}{2} - 1 \leq |N_i| \leq \frac{k}{2} + 1$  for  $i \in [5]$ . Note that the equality holds in (6), which implies that  $N_{G'}(v_i) = N_{i-1} \cup N_{i+1}$  holds for  $i \in [5]$ . Moreover, for any  $w \in N_i$ ,  $N_{G'}(w) \subseteq N_{i-1} \cup N_{i+1}$ . Let  $\mathcal{E} = \bigcup_{i=1}^5 E(H[N_i, N_{i+1}])$ . We define  $H_1 = H \setminus \mathcal{E}$  and  $G_1 = G' \cup \mathcal{E}$ . By our definition,  $G_1$  is also triangle-free. Similarly, we classify the edges of  $H_1$  (in fact, here we can think of  $H_1$  and  $G_1$  as the previous  $H$  and  $G'$  respectively). Let  $A_1 = \{v \in V(G) : d_{H_1}(v) \leq \frac{k}{14}\}$  and  $B_1 = V(G) - A_1$ . Since  $|E(H_1)| \leq |E(H)| \leq \frac{9}{4}k$ , we have  $|B_1| \leq 63$ .

Let  $F^0 = F_p \cup F_r$ , where  $F_p = \{xy \in E(H_1) : x, y \in N_i \text{ for some } i \in [5]\}$  and  $F_r = \{xy \in E(H_1) :$

$x \in N_i$ ,  $y \in N_j$  and  $|i-j| = 2$  for some  $i, j \in [5]$ . Let  $F^1 = F^0 \cap E(H_1[A_1])$ ,  $F^2 = F^0 \cap E(H_1[A_1, B_1])$  and  $F^3 = F^0 \cap E(H_1[B_1])$ . Note that  $F^0 = E(H_1)$ . Thus, we have

$$2(|F^1| + |F^2| + |F^3|) = 2|F^0| = \sum_{i=1}^5 \sum_{w \in N_i} d_{H_1}(w) = \sum_{i=1}^5 \left( \sum_{w \in A_1 \cap N_i} d_{H_1}(w) + \sum_{w \in B_1 \cap N_i} d_{H_1}(w) \right),$$

and  $|F^2| + 2|F^3| = \sum_{i=1}^5 \sum_{w \in B_1 \cap N_i} d_{H_1}(w)$ , which implies that

$$2|F^1| + |F^2| = \sum_{i=1}^5 \sum_{w \in A_1 \cap N_i} d_{H_1}(w). \quad (19)$$

For any integer  $i \in [5]$  and any vertex of  $w \in N_i \cap B_1$ , since  $|N_{i-1}| + |N_{i+1}| = k - a_i$ , there are at least  $|N_{i-1}| + |N_{i+1}| - (k - d_{H_1}(w)) - |B_1| = d_{H_1}(w) - a_i - |B_1|$  vertices of  $(N_{i-1} \cup N_{i+1}) \cap A_1$  are not adjacent to  $w$  in  $G$ . Thus, for  $i \in [5]$ , summing up the number of vertices of  $(N_{i-1} \cup N_{i+1}) \cap A_1$  that are not adjacent to  $w$  with  $w \in N_i \cap B_1$ , we can get that

$$\sum_{i=1}^5 \sum_{w \in A_1 \cap N_i} (d_{H_1}(w) - a_i) \geq \sum_{i=1}^5 \sum_{w \in B_1 \cap N_i} (d_{H_1}(w) - a_i - |B_1|). \quad (20)$$

By (19) and (20), since  $|N_i| \geq \frac{k}{2} - 1$  and  $|B_1| \leq 63$ , we can get that

$$\begin{aligned} 2|F^1| &= \sum_{i=1}^5 \sum_{w \in A_1 \cap N_i} d_{H_1}(w) - |F^2| \geq \sum_{i=1}^5 a_i |N_i \cap A_1| + \sum_{i=1}^5 \sum_{w \in A_1 \cap N_i} (d_{H_1}(w) - a_i) - \sum_{w \in B_1} d_{H_1}(w) \\ &\geq 2 \left( \frac{k}{2} - 1 - |B_1| \right) - \sum_{i=1}^5 \sum_{w \in B_1 \cap N_i} (a_i + |B_1|) \geq k - 128 - |B_1|(2 + |B_1|) \geq k - 4400. \end{aligned}$$

Let  $F_p^1 = F_p \cap F^1$  and  $F_r^1 = F_r \cap F^1$ . We can obtain the same conclusion as for the Claim 4.1 (replace  $G'$  and  $H$  with  $G_1$  and  $H_1$  respectively). Since  $G_1$  is triangle-free and  $E(G_1) = (E(G) \setminus E(H)) \cup \mathcal{E}$ , all the triangles we considered are disjoint from each other. Using Claim 4.1 and  $k > 10^8$ , we have

$$T(G) \geq \left( \frac{5}{14}k - 1400 \right) \left( \frac{k}{2} - 2200 \right) > \frac{k^2}{8} = \frac{k(3k - n - 1)}{4},$$

which is a contradiction to our assumption (4). We have completed the proof of Proposition 4.2.  $\blacksquare$

## 5 Concluding remarks

Our main result establishes a tight lower bound for the number of triangles in  $n$ -vertex  $k$ -regular graphs under certain conditions (specifically, when  $n$  is odd and sufficiently large, and  $k$  is an even integer satisfying  $\frac{2n}{5} < k < \frac{n}{2}$ ). The same problem was also investigated by Lo in [4, 5] for the case of  $k \geq \frac{n}{2}$ . For related problems concerning the count of cliques  $K_r$  in regular graphs or graphs with a bounded minimum degree, we recommend referring to [5].

It is also interesting to explore the minimum number  $t_H(n, k)$  of copies of a general graph  $H$  in an  $n$ -vertex  $k$ -regular graph. Results of Cambie, de Verclos, and Kang [2] indicate that there is a

distinction based on whether the chromatic number  $\chi(H) = 3$  or not (see Theorems 2, 13, and 14 in [2]). For all odd cycles  $C_{2\ell-1}$  where  $\ell \geq 2$ , Cambie et al. [2] obtained the following tight result:

If  $n$  is even and  $k > \frac{n}{2}$  or  $n$  is odd and  $k > \frac{2n}{2\ell+1}$ , then  $t_{C_{2\ell-1}}(n, k) > 0$ .

Assuming  $n$  is odd and  $k$  is even, satisfying  $\frac{2n}{2\ell+1} < k < \frac{n}{2}$ , it would be of great interest to determine the exact value of  $t_{C_{2\ell-1}}(n, k)$ . This endeavor to generalize Theorem 1.3 appears to be challenging.

## References

- [1] B. Andrásfai, P. Erdős and V. T. Sós, On the connection between chromatic number, maximal clique and minimal degree of a graph, *Discrete Math.*, **8**, (1974), 205–218.
- [2] S. Cambie, R. de Joannis de Verclos and R. J. Kang, Regular Turán numbers and some Gan-Loh-Sudakov-type problems, *J. Graph Theory*, **16**, (2022), 1–19.
- [3] A. S. L. Lo, Triangles in regular graphs with density below one half, *Combin. Probab. Comput.*, **18**, (2009), 435–440.
- [4] A. S. L. Lo, Cliques in graphs, PhD thesis, University of Cambridge, (2010).
- [5] A. S. L. Lo, Cliques in graphs with bounded minimum degree, *Combin. Probab. Comput.*, **21**, (2012), 457–482.
- [6] W. Mantel, Problem 28, *Winkundige Opgaven* 10, (1907), 60–61.
- [7] J. W. Moon and L. Moser, On cliques in graphs, *Israel J. of Math.*, **3**, (1965), 23–28.