

In situ molecular level studies on membrane related peptides and proteins in real time using sum frequency generation vibrational spectroscopy

Shuji Ye, Khoi Tan Nguyen, Stéphanie V. Le Clair, Zhan Chen *

Department of Chemistry, University of Michigan, 930 North University Ave., Ann Arbor, MI 48109, USA

ARTICLE INFO

Article history:

Received 4 November 2008

Received in revised form 11 March 2009

Accepted 13 March 2009

Available online 21 March 2009

Keywords:

Sum frequency generation

SFG

Cell membrane

Antimicrobial peptide

Membrane protein

Lipid bilayer

Melittin

G proteins

Alamethicin

Tachypletin I

ABSTRACT

Sum frequency generation (SFG) vibrational spectroscopy has been demonstrated to be a powerful technique to study the molecular structures of surfaces and interfaces in different chemical environments. This review summarizes recent SFG studies on hybrid bilayer membranes and substrate-supported lipid monolayers and bilayers, the interaction between peptides/proteins and lipid monolayers/bilayers, and bilayer perturbation induced by peptides/proteins. To demonstrate the ability of SFG to determine the orientations of various secondary structures, studies on the interactions between different peptides/proteins (melittin, G proteins, alamethicin, and tachypletin I) and lipid bilayers are discussed. Molecular level details revealed by SFG in these studies show that SFG can provide a unique understanding on the interactions between a lipid monolayer/bilayer and peptides/proteins in real time, *in situ* and without any exogenous labeling.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Cell membranes play a crucial role in many biological functions of cells. They govern all interactions between cells and their environments, such as the exchange of information and ions/molecules between the inside and outside of the cells. The cell membrane can be quite complex and dynamic, with a thickness of several nanometers. A cell membrane mainly consists of a lipid bilayer, which is an arrangement of lipids in which the acyl chains of each layer interact through hydrophobic interactions and the hydrophilic head groups face the inside and outside of the cell. A variety of peptides and proteins are also embedded inside or associated with the cell membranes, helping to fulfill various cellular functions (Katsaras and Gutberlet, 2001; Mateo et al., 2006; Yeagle, 2005). It is important to study the structures and kinetics of membrane embedded/associated peptides/proteins in order to understand their functions. Results from these types of studies can help in the rational design of molecules that can more effectively mediate or interfere with various cellular events in the desired manner.

Different experimental tools have been used to study peptides/proteins in the membrane environment, and excellent results have been obtained. For example, surface plasmon resonance spectroscopy

measurements allow for the determination of the peptide/protein coverage on membranes (Beseničar et al., 2006; Devanathan et al., 2006; Salamon et al., 1997). Neutron reflection studies have also permitted the determination of the amount of adsorbed peptides/proteins (Fragneto-Cusani, 2001; Haas et al., 2007; Kučerka et al., 2007). Atomic force microscopy techniques, on the other hand, are very useful in studying the packing and surface ordering of membrane-bound proteins (Alessandrini and Facci, 2005; Brassieur et al., 2008; Engel and Gaub, 2008; Johnston, 2007; Laflamme et al., 2008; Richter et al., 2006). Ellipsometry, in contrast, can be utilized to follow proteins' adsorption kinetics onto membranes and determine the adsorbed protein film thickness (Faiss et al., 2008). Excellent review articles have been published that summarize the applications of these techniques (and others) to the study of the interactions between peptides/proteins and lipid membranes (Cabiaux, 2004; Lee, 2005; McIntosh and Simon, 2006). Even though some molecular level information can be acquired by probing the interactions between cell membranes and proteins/peptides using these analytical tools, further details regarding such molecular interactions need to be elucidated.

Several spectroscopic techniques that can probe detailed structural information have been applied to study membrane-related peptides and proteins, leading to in-depth understanding of such molecular interactions. For example, Attenuated Total Reflectance Fourier Transform Infrared (ATR-FTIR) measurements can provide

* Corresponding author. Fax: +1 734 647 4865.

E-mail address: zhanc@umich.edu (Z. Chen).

vibrational spectra (or fingerprints) of membrane proteins and peptides. However, the technique may suffer from a lack of intrinsic surface sensitivity and therefore the signals from the molecules in the cell membrane environment might be confused with those in the bulk environment, e.g., in the bulk solution. Also, ATR-FTIR only gives one vibrational measurable of the tilt angle $\langle \cos^2\theta \rangle$ for orientation determination of a membrane protein or peptide (Tamm and Tatulian, 1997). Nuclear magnetic resonance (NMR) spectroscopy is another powerful spectroscopic technique that is widely used to characterize peptide/protein-lipid interactions (Bader et al., 2003; Bechinger, 1999, 2004; Lindblom and Grobner, 2006; Naito and Kawamura, 2007; Wang, 2008). Solution and solid-state NMR have been successfully used to determine the site-specific secondary structures and dynamics of membrane-bound proteins (Andronesi et al., 2005; Dürr et al., 2007; Fernández et al., 2001). Both solution and solid-state NMR can also be used to study several aspects of the lipid-protein interactions, such as looking at the proteins' effects on the lipid dynamics and determining which parts of the protein are interacting with the head group or tail region of the lipids (Dvinskikh et al., 2007; Lee et al., 2008). However, due to NMR's low sensitivity, large amounts of protein/peptide are required (milligram quantities), sometimes isotope-labeled proteins are needed, and experimental time can be very long. Difficulties can also be encountered in sample preparation (i.e., high concentrations of sample can cause membrane proteins to aggregate). The different model membranes used in NMR also make it complicated to create an asymmetric lipid bilayer (i.e., controlling the composition of each leaflet).

Over the last two decades, sum frequency generation (SFG) vibrational spectroscopy has been developed into a very powerful and highly versatile spectroscopic tool for surface and interfacial studies (Anglin and Conboy, 2008; Bain, 1995; Baldelli, 2008; Belkin and Shen, 2005; Chen, 2007; Dreesen et al., 2004a,b; Fourkas et al., 2007; Gautam and Dhinojwala, 2002; Gopalakrishnan et al., 2006; Gracias et al., 1999; Holman et al., 2004; Hopkins et al., 2005; Iwahashi et al., 2008; Kim et al., 2008; Koffas et al., 2004; Li et al., 2008; Ma and Allen, 2006; Moore and Richmond, 2008; Opdahl et al., 2004; Richmond, 2002; Rupprechter and Weilach, 2008; Shen and Ostroverkhov, 2006; Shultz et al., 2002; Stiopin et al., 2008; Voges et al., 2007; Yang et al., 2002; Ye et al., 2008). SFG is a vibrational technique that is intrinsically surface-sensitive, requires small amounts of sample, and with which the experiments can be done *in situ* and in real-time (Allen et al., 2000; Bain, 1995; Belkin and Shen, 2005; Buck and Himmelhaus, 2001; Chen, 2007; Chen and Chen, 2006; Chen et al., 2002b, 2005a; Eisenthal, 1992; Gopalakrishnan et al., 2006; Gracias et al., 1999; Hopkins et al., 2005; Koffas et al., 2004; Lambert et al., 2005; Miranda and Shen, 1999; Moore and Richmond, 2008; Opdahl et al., 2004; Perry et al., 2006; Richmond, 2001, 2002; Shen, 1989; Shen and Ostroverkhov, 2006; Shultz et al., 2000; Tadjeddine and Peremans, 1996; Wang et al., 2005a; Williams and Beattie, 2002; Zhuang and Shen, 1996). SFG permits the identification of interfacial molecular species (or chemical groups), and also provides information about the interfacial structure, such as the orientation and the orientation distribution of functional groups on the surface. SFG has been applied to study the structure and orientation of biomolecules, such as lipids (Anderson et al., 2006; Anglin et al., 2007; Anglin and Conboy, 2008; Chen et al., 2007b; Doyle et al., 2004; Harper and Allen, 2007; Kim et al., 2001; Levy and Briggman, 2007; Liu and Conboy, 2004a,b, 2005a,b, 2007; Lobau et al., 1999; Ma and Allen, 2006, 2007; Nickolov et al., 2006; Ohe et al., 2004; Petralli-Mallow et al., 1999; Sovago et al., 2007; Watry et al., 2003; White et al., 2006), and peptides/proteins (including membrane-related proteins/peptides) (Chen and Chen, 2006; Chen et al., 2005a,b; Clarke et al., 2005; Dreesen et al., 2004a,b; Evans-Nguyen et al., 2006; Humbert et al. 2006; Kim and Cremer, 2001; Kim et al.,

2002, 2003; Kim and Somorjai, 2003; Knoesen et al., 2004; Mermut et al., 2006; Rocha-Mendoza et al., 2007; Sartenaer et al., 2007; Wang et al., 2003a,b,c, 2005b, 2006; York et al., 2008).

Planar substrate-supported lipid bilayers have been widely used as a model to mimic cell membranes. Their suitability for biological studies has been extensively tested. These lipid bilayers are readily prepared by directly depositing lipid monolayers or bilayers onto the substrates such as glass, mica, quartz, and silicon surfaces using Langmuir-Blodgett method or vesicle fusion method (Kalb et al., 1992; Steinem et al., 2000; Tamm and McConnell, 1985; Tamm, 1988; Tamm and Tatulian, 1997; Thompson and Palmer, 1988). In addition, many research groups are also employing different strategies to improve the properties of supported lipid bilayers (i.e., using ultrathin polymer to support lipid bilayers) (Sackmann, 1996; Tanaka and Sackmann, 2005; Zhao and Tamm, 2003). Previous studies have indicated that planar substrate-supported lipid bilayers can offer several advantages over other model membranes, such as free-standing lipid bilayers, solvent-free lipid bilayers, or phospholipid vesicles. Planar substrate-supported lipid bilayers are unilamellar and geometrically well defined. They can maintain excellent mechanical stability without losing their fluid nature. These advantages of substrate-supported bilayers make it possible to carry out experiments that probe structural and dynamic properties of membranes and protein-lipid interactions, using the surface analytical techniques mentioned above (Castellana and Cremer, 2006; Kalb et al., 1992; McConnell et al., 1986; Sackmann, 1996; Tamm and McConnell, 1985; Tamm, 1988; Tamm and Tatulian, 1997; Tanaka and Sackmann, 2005).

In this paper, we will first present a brief introduction of the theoretical background needed to understand SFG, and then summarize recent studies on the interactions between lipid membranes (monolayers and bilayers, focusing especially on substrate-supported lipid bilayers) and biomolecules monitored by SFG in real time and *in situ*. The names of lipids mentioned in this paper and their respective abbreviations are listed in Table 1.

2. Theoretical background of SFG

Although it has only been about twenty years since the first SFG spectra were recorded by Shen (Guyotsonnest et al., 1987; Hunt et al., 1987; Zhu et al., 1987), SFG is now being used by a growing number of research groups for a variety of applications, including polymer and biological interface studies. Many excellent review

Table 1

The full names of lipids and their abbreviations that appear in this paper.

Abbr.	Full Name
POPC	1-Palmitoyl-2-Oleoyl-sn-Glycero-3-Phosphocholine
POPG	1-Palmitoyl-2-Oleoyl-sn-Glycero-3-[Phospho-rac-(1-glycerol)] (Sodium Salt)
DMPC	1,2-Dimyristoyl-sn-Glycero-3-Phosphocholine
d-DMPC	1,2-Dimyristoyl-D54-sn-Glycero-3-Phosphocholine-1,1,2,2-D4-N,N,N-trimethyl-D9
DMPC-d54	1,2-Dimyristoyl-D54-sn-Glycero-3-Phosphocholine
DPPC	1,2-Dipalmitoyl-sn-Glycero-3-Phosphocholine
d-DPPC	1,2-Dipalmitoyl-D62-sn-Glycero-3-Phosphocholine-1,1,2,2-D4-N,N,N-trimethyl-D9
DPPG	1,2-Dipalmitoyl-sn-Glycero-3-[Phospho-rac-(1-glycerol)] (Sodium Salt)
d-DPPG	1,2-Dipalmitoyl-D62-sn-Glycero-3-[Phospho-rac-(1-glycerol)] (Sodium Salt)
DSPC	1,2-Distearoyl-sn-Glycero-3-Phosphocholine
d-DSPC or DSPC-d83	1,2-Distearoyl-D70-sn-Glycero-3-Phosphocholine-1,1,2,2-D4-N,N,N-trimethyl-D9
DSPC-d70	1,2-Distearoyl-D70-sn-Glycero-3-Phosphocholine
DSPG	1,2-Distearoyl-sn-Glycero-3-[Phospho-rac-(1-glycerol)] (Sodium Salt)




Fig. 1. SFG experimental geometry and energy diagram. (A) Total reflection geometry employed for the experiments described in Sections 3.3 and 3.4. A bilayer is immersed in a small reservoir with an approximate volume of 1.6–2.0 mL; (B) Simplified energy level diagram of vibrational sum frequency generation process.

papers have summarized details about SFG theory and research (Allen et al., 2000; Bain, 1995; Belkin and Shen, 2005; Buck and Himmelhaus, 2001; Chen, 2007; Chen and Chen, 2006; Chen et al., 2002b, 2005a; Eienthal, 1992; Gopalakrishnan et al., 2006; Gracias et al., 1999; Hopkins et al., 2005; Koffas et al., 2004; Lambert et al., 2005; Miranda and Shen, 1999; Moore and Richmond, 2008; Opdahl et al., 2004; Perry et al., 2006; Richmond, 2001, 2002; Shen, 1989; Shen and Ostroverkhov, 2006; Shultz et al., 2000; Tadjeddine and Peremans, 1996; Wang et al., 2005a; Williams and Beattie, 2002; Zhuang and Shen, 1996), which will not be repeated here. In a typical SFG experiment, two pulsed laser beams, one with a fixed frequency in the visible frequency range (ω_{vis}) and one with a tunable frequency in the infrared frequency range (ω_{IR}), are overlapped spatially and temporally on the sample (Fig. 1A). The SFG signal is generated at the sum frequency of the two input beams by the nonlinear process, $\omega_{\text{SF}} = \omega_{\text{vis}} + \omega_{\text{IR}}$. Therefore, the SFG process can be simply viewed as a combination of infrared (IR) absorbance and Raman scattering, as shown in the energy diagram in Fig. 1B. The intensity of the sum frequency signal is proportional to the square of the vibration's second-order nonlinear susceptibility $\chi_{\text{eff}}^{(2)}$ (Bain, 1995; Buck and Himmelhaus, 2001; Chen and Chen, 2006; Chen et al., 2002b, 2005a; Eienthal, 1992; Lambert et al., 2005; Miranda and Shen, 1999; Richmond, 2001; Shen, 1984, 1989; Tadjeddine and Peremans, 1996; Wang et al., 2005a; Williams and Beattie, 2002; Zhuang et al., 1999). Different components of $\chi_{\text{eff}}^{(2)}$ can be probed using different polarization combinations of the input and output laser beams. From such measurements, orientation information of surface molecules and functional groups can be deduced (Gautam and Dhinojwala, 2001; Hirose et al., 1992a,b, 1993; Kim and Somorjai, 2003; Shen, 1984; Wang et al., 2001). More details about SFG theory and data analysis can be found in Appendix A.

The selection rules of SFG make the technique surface sensitive. As we stated above, the SFG signal intensity is proportional to the square of the vibration's second-order nonlinear susceptibility $\chi_{\text{eff}}^{(2)}$. $\chi_{\text{eff}}^{(2)}$ vanishes when a material has inversion symmetry under the dipole approximation (Bain, 1995; Buck and Himmelhaus, 2001; Chen and Chen, 2006; Chen et al., 2002b, 2005a; Eienthal, 1992; Gautam and Dhinojwala, 2001; Hirose et al., 1992a,b, 1993; Lambert et al., 2005; Miranda and Shen, 1999; Richmond, 2001; Shen, 1984, 1989; Tadjeddine and Peremans, 1996; Wang et al., 2001, 2005a; Williams and Beattie, 2002; Zhuang et al., 1999). The majority of bulk materials exhibits inversion symmetry, thus they do not generate SFG signals. However, the presence of an interface causes a break in the symmetry, and therefore, molecules on surfaces or at interfaces can generate SFG signal. Therefore, SFG is an intrinsically surface/interface specific technique, different from linear vibrational spectroscopic techniques such as ATR-FTIR and Raman scattering.

3. Recent SFG studies on membrane-related peptides/proteins

In this section, we will summarize recent SFG studies on membrane-related peptides and proteins. Such studies have demonstrated that SFG can provide a unique understanding of the interactions between a lipid monolayer/bilayer and peptides/proteins without any exogenous labeling, in real time and *in situ*.

3.1. SFG studies on lipid monolayers, substrate-supported lipid bilayers and hybrid bilayer membranes

As mentioned above, it has been extensively shown that substrate-supported lipid bilayers are valid models for cell membranes (Castellana and Cremer, 2006; Kalb et al., 1992; McConnell et al., 1986; Sackmann, 1996; Steinem et al., 2000; Tamm and McConnell, 1985; Tamm, 1988; Tamm and Tatulian, 1997; Tanaka and Sackmann, 2005; Thompson and Palmer, 1988; Zhao and Tamm, 2003). SFG has been applied to investigate the structures of these lipid bilayer systems (Anglin et al., 2007; Anglin and Conboy, 2008; Chen et al., 2007b; Doyle et al., 2004; Levy and Briggman, 2007; Liu and Conboy, 2004a,b, 2005a,b, 2007; Petrali-Mallow et al., 1999), with a recent review article (Chen and Chen, 2006) summarizing these SFG studies. Here, we do not attempt to reiterate the details, but instead will give some brief discussions on how some fundamental questions regarding the interactions between peptides/proteins and lipid bilayers can be answered using SFG. For example, knowledge of the conformations of lipids in the bilayer, both before and after their interactions with antimicrobial peptides, aids in understanding the peptide's mode of interaction. These modes of interaction can be generally separated into the carpet, barrel-stave and toroidal pore modes (Yang et al., 2001; Chen and Chen, 2006). Theoretically, each of these peptide/protein–lipid bilayer interaction modes should induce different conformational changes on the lipid bilayers, resulting in distinct SFG spectral changes. Therefore, using these kinds of SFG lipid studies, details regarding the interactions of antimicrobial peptides, and other peptides/proteins, with lipid systems may be elucidated.

Before we go further to discuss the interactions between peptides/proteins and lipid bilayers, knowledge regarding the membrane model system, the lipid bilayer, is needed. When studying such interactions with SFG, substrate-supported lipid bilayers are used as the typical model for a cell membrane. Optical techniques using fluorescence labeling methods have previously been used to study the physical properties of such bilayers. Although these techniques offer important insights (e.g., about the flip-flop of lipids in a membrane, the transition temperature of a lipid system, and lipid domain segregation) (Bagatolli, 2006; Heberle et al., 2005; Lee, 2005), the incorporation of a fluorescent label on the lipids, or of a fluorescent probe inside the bilayer, may affect the behavior of

the system and generate some artifacts in the experiments. An advantage of using SFG to study these types of supported bilayers is that it is not necessary to introduce any bulky labels. The only form of labeling required to study the bilayers is the introduction of isotope labeling. In order to monitor each leaflet of a lipid bilayer individually, it is necessary to have one leaflet deuterated, in order to break the inversion symmetry of the bilayer and obtain SFG signal.

One important aspect of the behavior of lipids in cell membranes is their ability to translocate both laterally and across the membrane (flip-flop). With this in mind, Conboy and his colleagues have investigated some important physical properties of substrate-supported lipid bilayers, such as the kinetics of the flip-flop and the transition temperatures of various lipid bilayers using SFG (Liu and Conboy, 2004a,b, 2005a,b). More recently, they investigated the asymmetric distribution of domains in lipid bilayers by carrying out SFG spectroscopic measurements of symmetric C–H stretching modes of the fatty acid methyl groups (Liu and Conboy, 2007). In this research, they correlated the intensity of the C–H symmetric stretch of the fatty acid methyl groups with the symmetry of the lipid bilayer, with the stronger intensity obtained when the bilayer became more asymmetric. The breakage in symmetry was caused by the dislocation of the gel and liquid-crystalline phase domains at the transition temperature. They have also done SFG studies on the lateral pressure dependence of the phospholipid transmembrane diffusion rate in supported lipid bilayers (Anglin and Conboy, 2008). Their results indicated that the kinetics of lipid flip-flop in these membranes show a strong lateral pressure dependence. Based on these data, they successfully determined the activation area for phospholipid flip-flop (Anglin and Conboy, 2008).

Briggman and coworkers have employed hybrid bilayer membranes (HBM) as an alternative for supported lipid bilayers in their research (Anderson et al., 2004, 2006, 2007; Levy and Briggman, 2007; Petralli-Mallow et al., 1999). The HBM system studied has one single lipid layer deposited on top of a hydrophobic self-assembled monolayer (SAM). Briggman's group has extensively studied the properties of HBMs and found that the transition temperature of the lipid layer depends greatly on the packing density and the crystallinity of the SAM layer underneath it (Anderson et al., 2007). They also carried out SFG studies on the effect of cholesterol on phospholipids (Levy and Briggman, 2007). Their research has shown that HBMs could be used as excellent model membranes for biological studies on peripherally-bound proteins. However, it is challenging to use HBMs as cell membrane models to study lipid flip-flop and some transmembrane proteins that extensively interact with the inner-leaflet.

Many research groups also model cell membranes by using lipid monolayers. The fundamental properties such as transition temperatures and effects of cholesterol on monolayers have been investigated using SFG (Bonn et al., 2004; Ohe et al., 2007a,b; Roke et al., 2003).

3.2. SFG studies on the interactions between peptides/proteins and lipid monolayers/bilayers: C–H stretching frequency region

Early SFG studies on the interactions between proteins/peptides and lipid monolayers, HBMs and supported lipid bilayers focused on the C–H stretching frequency region. Table 2 contains a list of some of those studies. One of the very first SFG studies on the interaction between proteins and lipid monolayers was done by Cremer and his colleagues (Kim et al., 2003). They investigated the orientation of gramicidin A in a DMPC monolayer using the SFG C–H stretching signals generated from the side chains of gramicidin A. Their results indicated that the orientation of gramicidin A was concentration dependent (Kim et al., 2003).

Conboy and coworkers studied the effect of gramicidin A on the flip-flop of DSPC lipids in a substrate-supported phospholipid bilayer (Anglin et al., 2007). Using SFG and ATR-FTIR, they showed that gramicidin A induced rapid flip-flop of the DSPC lipids. In this study, the C–H stretching signals of the lipid's hydrophobic chain were studied by SFG and the amide signal of gramicidin A was observed by ATR-FTIR. The detection of amide signal from the peptide using ATR-FTIR indicated that gramicidin A was bound to the bilayer. SFG signal of the lipid C–H stretches could be observed because an asymmetric bilayer (with one leaflet deuterated) was used in the study. The time-dependence of the SFG C–H signals can be used to monitor the flip-flop rate, because as the system becomes more symmetric (from interchange between outer- and inner-leaflet lipids), the signal should decrease. By studying the time-dependent C–H signals both with and without gramicidin A bound, they showed that gramicidin A induced a faster flip-flop rate than when no gramicidin A was added to the bilayer.

Neivandt and coworkers examined a protein in the fibroblast growth factor (FGF) family, FGF 1, and its interaction with an HBM (Doyle et al., 2004). Using SFG, they showed that this protein caused deformation of the DSPG lipid layer even at very low concentrations. They also found that this process was reversible to a certain extent when the protein was washed off the HBM. Itoh and his colleagues investigated the interactions between antibiotic polymyxin B (PMB) and monolayers composed of DPPG, as well as DPPC (Ohe et al., 2004). By monitoring the SFG C–H stretching signals of the lipids' side chains and the O–H stretching signals of water molecules, along with the measurements of the pressure-area isotherms, they found that this antibiotic peptide bound to the negatively charged lipids (DPPG), but not to the neutral lipids (DPPC). They also showed that the binding of PMB to the lipid monolayer affected the phase transition of the lipid monolayer.

Because SFG can provide structural information such as functional group composition, orientation, and ordering at a surface or interface with a submonolayer sensitivity, it has also been applied to determine the average minimal inhibitory concentration (MIC) of antimicrobial peptides (or analogue oligomers) in membranes. Chen et al. used SFG to investigate the molecular interactions between a small antimicrobial oligomer and a single substrate-supported lipid bilayer with a hydrogenated leaflet (DPPG) and a deuterated leaflet (d-DPPG) (Chen et al., 2006). It was observed from the C–H and C–D stretching signals that the distal leaflet was disrupted at a very low peptide/lipid ratio, while the proximal leaflet remained intact below a threshold concentration very close to the MIC (0.8 mg/mL) value. The orientation of this oligomer was deduced by SFG C–H signals and the results indicated that the small antimicrobial oligomer acted as a “molecular knife” by disrupting primarily the outer leaflet of the bilayer at lower concentrations and further inserting into the entire bilayer at a certain critical (MIC) concentration.

3.3. SFG studies on the interaction between peptides/proteins and lipid bilayers: amide I frequency region

3.3.1. Detection of SFG amide I signal

The SFG results presented in the previous section are obtained from SFG signals acquired mainly in the C–H stretching frequency region, which are comprised of signals from the lipids and protein side chains. The feasibility of detecting SFG amide I signals from interfacial proteins/peptides was demonstrated by Chen and his coworkers in 2003 (Wang et al., 2003b). Because the water bending mode does not contribute noticeable SFG signals, it is not necessary to perform a background subtraction to obtain the SFG amide I signals from interfacial proteins and peptides; instead, they can be detected directly (Clarke et al., 2005; Wang et al., 2003b, 2005c). In addition, proteins in the bulk solution do not generate SFG sig-

Table 2

SFG studies on the interactions between proteins/peptides and lipid monolayers, HBMs and supported lipid bilayers.

Peptides or Proteins	Membrane	Studied region	Reference
Gramicidin A	DMPC-d54 monolayer	CH	Kim et al. (2003)
Polymyxin B	DPPC monolayer, DPPG monolayer	CH	Ohe et al. (2004)
FGF-1	DSPG hybrid bilayer membranes	CH	Doyle et al. (2004)
Antimicrobial Oligomers	DPPG/d-DPPG bilayer	CH, CD, Amide I	Chen et al. (2006)
Antimicrobial peptides	DPPG/d-DPPG bilayer	CH, CD, Amide I	Chen and Chen (2006)
Gramicidin A	DSPC/DSPC-d70 bilayer, DSPC-d83/DSPC-d83 bilayer	CH, CD	Anglin et al. (2007)
Melittin	dDPPG/dDPPG bilayer, dDPPG/DPPG bilayer	CH, CD	Chen et al., 2007b
Melittin	DPPG/DPPG	Amide I	Chen et al. (2007c)
Heterotrimeric G Protein $\beta\gamma$ Subunit	POPC/POPC bilayer, POPC/POPG bilayer	Amide I	Chen et al., 2007a

nals, and thus SFG can selectively probe interfacial proteins/peptides. Moreover, SFG is able to provide more measurements than ATR-FTIR in studying the orientation of interfacial proteins/peptides. In SFG studies, our group has also adopted a near total reflection experimental geometry (Fig. 1A) that enables us to obtain very strong SFG amide I signals of interfacial proteins, which makes the data analysis easier and more accurate (Wang et al., 2003b).

SFG amide I signals of proteins can be affected by the surface coverage, orientation, and secondary structures of the adsorbed proteins (Clarke et al., 2005; Wang et al., 2003b, 2005c). The amide I mode contains predominately the peptide C=O stretching bands. These C=O groups are held together by hydrogen bonds within the secondary structures and the frequency of the C=O stretch depends heavily on its hydrogen-bonded environment. The peak center of the amide I band, therefore, depends on the secondary structure adopted by the peptide/protein. The Chen group has successfully demonstrated that SFG amide I signals can be used to distinguish α -helical and β -sheet structures of peptides and proteins on polymers and lipid bilayers (Chen and Chen, 2006; Chen et al., 2005b, 2007a,c; Clarke et al., 2005; Wang et al., 2003b, 2005b,c, 2007, 2008). Recently, Somorjai's group developed a new optical parametric amplifier (OPA) in the SFG spectrometer that can also create a tunable infrared light between 1500 and 2000 cm^{-1} ; this IR beam can be used to study the amide I signals of interfacial peptides (York et al., 2008).

Based on previous FTIR and Raman studies, SFG amide signals of different secondary structures can be assigned. Using antimicrobial peptides that are known to adopt α -helical structures in lipid bilayers, Chen et al. were able to detect SFG amide I signals of α -helical structures centered at around 1650 cm^{-1} (Chen et al., 2007c). Using model β -sheet peptide tachyplesin I, Chen et al. also showed that SFG amide I signals from a β -sheet structure has characteristic peaks at 1635 and 1685 cm^{-1} , corresponding to the B2 mode and B1/B3 modes, respectively (Chen and Chen, 2006). Recent results in our laboratory on alamethicin indicate that SFG amide I signals of 3_{10} helical structures have peaks at around 1635 and 1670 cm^{-1} . These results are indeed in good agreement with previous FTIR and Raman studies. More details regarding the interfacial structures of different secondary structure domains of proteins/peptides can be obtained by the careful data analysis of SFG amide I signals collected using different polarization combinations, which will be presented below.

3.3.2. SFG studies on α -helical proteins/peptides in lipid bilayers

3.3.2.1. Data analysis for orientation determination using amide I band. As mentioned above, it has been shown experimentally that amide I peak centers are different for each type of secondary structure. Therefore, analyses of amide I signals provide insights into the protein backbone secondary structures. This section will focus on how to deduce the orientation of α -helical structures. Conventional polarized ATR-FTIR has been widely applied to study the orientation of α -helical structures of membrane peptides/proteins (Tamm

and Tatulian, 1997). In ATR-FTIR studies, the tilt angle of the α -helices can be calculated from the order parameter (S_θ), which is defined as:

$$S_\theta = \frac{3 \langle \cos^2 \theta \rangle - 1}{2}, \quad (1)$$

with θ being the tilt angle between the helix's principal axis and the surface normal. The bracket denotes the time and ensemble average. Theoretically, $\langle \cos^2 \theta \rangle$ can be determined from the measured intensity ratio in ATR-FTIR using p- and s-polarized IR light (Tamm and Tatulian, 1997). If we assume θ to have the simplest delta distribution, the orientation of the helix can be determined from this intensity ratio. The orientation in reality can be more complicated and such a simple distribution may not be adequate enough to describe the orientation of the peptide/protein. For example, a helix may adopt two different orientations or have a broad orientation distribution. To characterize such a more complicated orientation distribution, more measured parameters are needed. For example, when the parameter S_θ approaches zero, $\langle \cos^2 \theta \rangle$ is equal to 1/3, and thus there is always ambiguity in whether all helices have the same tilt angle around 54.7°, or a completely random orientation, or other orientations/distributions in between. In order to determine which of the three cases is correct, more measurements using different spectroscopic methods would be required (Chen et al., 2007c).

It has been shown that it is possible to deduce the orientation of functional groups, such as methyl, methylene, phenyl, and carbonyl, by measuring different tensor elements using SFG spectra collected with different polarization combinations (Briggman et al., 2001; Gautam et al., 2000; Hirose et al., 1992a,b, 1993; Oh-e et al., 2002; Tyrode et al., 2005). By analyzing the SFG amide I signals using group theory and projection operators, the Chen group has been able to investigate interfacial protein structures and deduce the orientation of α -helical peptides (Lee et al., 2006; Wang et al., 2007, 2008). They showed that both the amide I A mode and amide I E_1 mode of an α -helix can contribute to SFG signals (Wang et al., 2008). Using the near total internal reflection geometry (Wang et al., 2003b), ssp and ppp amide I signals can be collected. These signals are mainly due to contributions from the χ_{yyz} and χ_{zzz} susceptibility components, respectively. The dependence of χ_{yyz} and χ_{zzz} susceptibility components on the molecular hyperpolarizability is described by the following equations:

For the A mode:

$$\chi_{A,xxz} = \chi_{A,yyz} = \frac{1}{2} N_s [(1+r) \langle \cos \theta \rangle - (1-r) \langle \cos^3 \theta \rangle] \beta_{ccc} \quad (2)$$

$$\chi_{A,zzz} = N_s [r \langle \cos \theta \rangle + (1-r) \langle \cos^3 \theta \rangle] \beta_{ccc} \quad (3)$$

For the E_1 mode:

$$\chi_{E1,xxz} = \chi_{E1,yyz} = -N_s (\langle \cos \theta \rangle - \langle \cos^3 \theta \rangle) \beta_{aca} \quad (4)$$

$$\chi_{E1,zzz} = 2N_s (\langle \cos \theta \rangle - \langle \cos^3 \theta \rangle) \beta_{aca} \quad (5)$$

where β_{aca} and β_{ccc} are the molecular hyperpolarizability elements. The hyperpolarizability elements of an α -helix can be obtained from the product of the components of the Raman polarizability and IR transition dipole moment. Chen et al. (Chen et al., 2007c) deduced the relations among different hyperpolarizability tensor elements to be $r = \beta_{aad}/\beta_{ccc} \approx 0.54$ and $\beta_{aca} \approx 0.32 \beta_{ccc}$ (Lee and Krimm, 1998a,b; Marsh et al., 2000; Rintoul et al., 2000). N_s is the number density of ideal α -helix units composed of 18 amino acid residues. Due to the limited resolution of many SFG spectrometers ($\sim 5 \text{ cm}^{-1}$ or more), the A mode and E_1 mode cannot be readily resolved in the frequency domain, and therefore, the total susceptibility is often assumed to be the sum of the susceptibilities from these two modes (Chen et al., 2007c):

$$\chi_{yyz} = \chi_{A,yyz} + \chi_{E_1,yyz} \quad (6)$$

$$\chi_{zzz} = \chi_{A,zzz} + \chi_{E_1,zzz} \quad (7)$$

As Eqs. (2)–(5) indicate, only two measurables related to the orientation angle are independent: $\langle \cos\theta \rangle$ and $\langle \cos^3\theta \rangle$. Using different polarization combinations of the input and output laser beams, $\langle \cos\theta \rangle$ and $\langle \cos^3\theta \rangle$ can be deduced. If all α -helical structures on the surface/interface adopt the same orientation, $\langle \cos\theta \rangle$ and $\langle \cos^3\theta \rangle$ can be replaced by $\cos\theta$ and $\cos^3\theta$. Subsequently, the relationship between an SFG measurable and the orientation angle of the α -helix can be depicted. However, for most cases, it may not be correct to assume that all the α -helical structures on a surface/interface have the same orientation. For example, one protein may have two α -helical segments pointing two different directions. The following section discusses the orientation analysis of α -helical structures in some of these complex cases.

3.3.2.2. Example: α -helical melittin in membrane. Recently, Chen et al. used melittin as a model peptide to study the orientation of α -helical peptides in substrate-supported DPPG bilayers (Chen et al., 2007c). Chen et al. found that the SFG measurements were not compatible to those of a δ -distribution or a Gaussian distribution. These two distributions were therefore not adequate to describe the melittin orientation distribution inside a DPPG bilayer and the orientation distribution had to be more complex. Chen et al. assumed two δ -distributions as the orientation function, meaning that melittin was assumed to adopt two distinct orientations in the lipid bilayer. A fraction (N) of melittin molecules may orient with an angle of θ_1 , and a fraction ($1-N$) of melittin molecules can orient with another angle of θ_2 . By combined ATR-FTIR and SFG studies, all of these parameters, θ_1 , θ_2 and N , were successfully deduced. The obtained results from SFG and ATR-FTIR experiments indicated that melittin helices existed in two main populations in the lipid bilayer. About three-fourths of melittin molecules oriented parallel to the bilayer surface with a slight tilt, while the rest oriented more or less parallel to the surface normal, as shown in Fig. 2A and B.

In addition, Chen et al. also introduced the maximum entropy function to deduce the orientation distribution of melittin in a single lipid bilayer based on the ATR-FTIR and SFG measurements. Such a treatment does not have any assumptions regarding the orientation distribution function, e.g., assuming two δ -distributions. The deduced orientation distribution using the maximum entropy function was very similar to that obtained from the two δ -distributions, as shown in Fig. 2C. This research demonstrated the power of combining ATR-FTIR measurements, SFG data and the maximum entropy function analysis for deducing complicated orientations of membrane-bound peptides. These kinds of orientation determination results can be correlated to different modes of action of peptides' interactions with bilayers, and ultimately lead to an understanding of the mechanism of antimicrobial activity, for

example. Such an analysis can also be applied to study interfacial proteins with two (or even three) α -helical segments that adopt different orientations. A similar method has also been used to study the orientation distribution of two α -helical coiled-coils of fibrinogen at the polystyrene/fibrinogen solution (phosphate buffered solution with a total ionic strength of 0.14 M and a pH value of 7.4) interface (Wang et al., 2008).

3.3.2.3. Example: α -helical structure in G protein in lipid bilayer. Heterotrimeric guanine nucleotide-binding proteins (G proteins) are a family of peripheral membrane proteins that transduce extracellular signals (e.g., hormones and neurotransmitters), as sensed by G protein-coupled-receptors (GPCRs), to intracellular effector systems (e.g., ion channels and cell transcription machinery) (Cabrera-Vera et al., 2003; Neves et al., 2002). Each G protein is comprised of $G\alpha$, $G\beta$, and $G\gamma$ subunits, with $G\beta$ and $G\gamma$ forming a tightly associated dimer. In the resting state, a G protein exists in the $G\alpha\beta\gamma$ form. Upon GPCR activation, $G\alpha\beta\gamma$ releases the GDP originally bound to the $G\alpha$ subunit and the binding of GTP allows dissociation of $G\alpha\beta\gamma$ into $G\alpha$ -GTP and $G\beta\gamma$ (Gaudet et al., 1999; Lodowski et al., 2003). $G\alpha$ and $G\beta\gamma$ can then associate with their own effectors and trigger downstream signaling cascades. The cycle returns to the resting state when $G\alpha$ hydrolyzes GTP back to GDP (Koch, 2004; Neves et al., 2002; Oldham and Hamm, 2006; Pitcher et al., 1992).

Recently, (Chen et al., 2007a) investigated how the $G\beta_1\gamma_2$ subunit binds to and orients on a substrate-supported lipid bilayer using SFG. Both wild-type $G\beta_1\gamma_2$ subunits, which contain a geranylgeranyl anchor group, and only the soluble domain of the $G\beta_1\gamma_2$ subunits were used in this research. SFG spectra were collected from both types of G protein subunits in a hydrated POPG/POPG bilayer. These two $G\beta_1\gamma_2$ subunits showed very different SFG spectral properties (Fig. 3C and D). Even at relatively high concentrations (125 $\mu\text{g}/\text{mL}$), soluble $G\beta_1\gamma_2$ generated weaker signals than geranylgeranylated $G\beta_1\gamma_2$, with a peak centered at 1630 cm^{-1} , indicative of β -sheet secondary structure. The presence of the geranylgeranyl anchor group resulted in a significant enhancement of SFG amide I signal that was dominated by a peak at around 1650 cm^{-1} , which is characteristic of contributions from an α -helical structure. (Chen et al., 2007a) suggested that without the geranylgeranyl group, $G\beta_1\gamma_2$ adsorbs onto the surface with the β -propeller domain facing the membrane surface and the helical domains orienting more or less parallel to the surface (Fig. 3B, Chen et al., 2007a). On the other hand, for the wild-type $G\beta_1\gamma_2$ subunit, the β -propeller more or less orients perpendicular to the bilayer surface and the helical domains are ordered and no longer parallel to the surface. This orientation allows the β -propeller to keep its native semi-centrosymmetry, resulting in very weak β -sheet signal and causing the amide I signal to be dominated by a peak at 1650 cm^{-1} (originating from the ordered helical domains), as shown in Fig. 3A. From the measured SFG ppp and ssp intensity ratio, Chen et al. deduced the orientation angle of the wild-type $G\beta_1\gamma_2$ to be -35° from a reference orientation in which the β -sheets within the β -propeller are parallel to the membrane surface (Tesmer et al., 2005; Wall et al., 1995). This research demonstrates that SFG can be used to deduce the membrane protein orientation *in situ* by studying the orientation of α -helical components of a protein.

3.3.3. SFG studies on 3_{10} helical peptides in membrane bilayer: alamethicin

Alamethicin is a 20-residue hydrophobic antibiotic peptide that can form voltage-gated ion channels in membranes. It has been used frequently as a model for larger channel proteins (Tamm and Tatulian, 1997). In addition to the regular amino acids, the peptide contains eight aminoisobutyric acid units. Its crystal struc-




Fig. 2. (A) Schematic of melittin's two orientations in the lipid bilayer. (B) Orientation distribution function derived based on a dual δ -distribution. (C) Orientation distribution function derived based on the maximum entropy theory. Reproduced with permission from *J. Am. Chem. Soc.* 2007, 129, 1420–1427. Copyright 2007, American Chemical Society.

ture contains an α -helical domain and a 3_{10} helical domain (Fox and Richards, 1982).

An extensive amount of research has been performed to examine the alamethicin action mechanism on membranes (Cafiso, 1994; Duchohier and Wroblewski, 2001; Hall et al., 1984; Leitgeb et al., 2007; Mathew and Balaram, 1983a,b; Nagaraj and Balaram, 1981; Sansom, 1993a,b; Woolley and Wallace, 1992). It is currently believed that alamethicin interacts with cell membranes through the barrel-stave mode (Duchohier, 2004; Fox and Richards, 1982; Laver, 1994; Mathew and Balaram, 1983a,b; Sansom, 1993a,b) with the resulting conducting pores in the membrane formed by

parallel bundles of 3–12 helical alamethicin monomers surrounding a central, water-filled pore. However, further details on the structural origin of some important properties of alamethicin channels in the membrane, such as the strong dependence of their conductivity on the transmembrane potential (Stella et al., 2007), are not known. In addition, contradicting orientations of alamethicin in the membrane in the absence of voltage have been reported. Alamethicin has been suggested to adopt a transmembrane orientation (Bak et al., 2001; Kessel et al., 2000; Marsh et al., 2007a; North et al., 1995), lie on the membrane surface (Banerjee et al., 1985; Ionov et al., 2000; Mottamal and Lazaridis, 2006), or both




Fig. 3. Schematics of $G\beta_1\gamma_2$ adsorbed onto a POPG/POPG bilayer deduced from SFG spectra: (A) wild-type $G\beta_1\gamma_2$ with geranylgeranyl group, (B) soluble $G\beta_1\gamma_2$ without geranylgeranyl group. SFG amide I spectra of the interfacial $G\beta_1\gamma_2$ adsorbed onto a POPG/POPG bilayer. (C) 250 μg of the soluble form of $G\beta_1\gamma_2$ was first injected into the subphase (~ 2 mL) of the bilayer. Only relatively weak signals indicative of β -sheet were observed. (D) After the spectra in panel C were collected, 50 μg wild type $G\beta_1\gamma_2$ was injected into the subphase and stronger SFG signals indicative of α -helix structure were observed. Reproduced with permission from *J. Am. Chem. Soc.*, 2007, 129, 12658–12659. Copyright 2007, American Chemical Society.

(depending on the experimental conditions) (Chen et al., 2002a; Huang and Wu, 1991). A continuous distribution of orientations has also been proposed (Spaar et al., 2004). Because of this lack of agreement in the literature, our lab has also been studying the molecular interactions between alamethicin and lipid bilayers *in situ* and in real time using SFG.

SFG ssp and ppp spectra of alamethicin in a d-DMPC/DMPC bilayer are shown in Fig. 4A. The spectra were collected after 37.5 μg of alamethicin (dissolved in 15 μL methanol) was injected into the water subphase (~ 1.6 mL) of a d-DMPC/DMPC bilayer for 71 min at pH 6.7. The SFG spectra were dominated by two peaks at 1635 and 1670 cm^{-1} . Peak assignments in the literature indicate that the 1635 cm^{-1} peak is due to the 3_{10} -helix, while the peak at 1670 cm^{-1} has contributions by both the 3_{10} helix and the α -helix (Dwivedi and Krimm, 1984; Haris and Chapman, 1988; Haris et al., 2004; Kennedy et al., 1991; Vogel, 1987). The orientation analysis method for a 3_{10} -helix in the membrane, using the SFG amide I

band, has been developed in our lab, similar to the method for an α -helix discussed in Section 3.3.2.1. This method is now being applied to deduce the orientation of alamethicin in a d-DMPC/DMPC bilayer using the spectra shown in Fig. 4A. Detailed results will be reported in the future.

It has been shown that the membrane lipid chain length affects the interaction between alamethicin and the cell membrane (Archer et al, 1991; Barranger-Mathys and Cafiso, 1994; Hall et al., 1984; Marsh et al., 2007a,b; Marsh, 2008). We observed markedly different SFG signal intensities from alamethicin in lipid bilayers with lipids of different chain lengths (Table 3). The length of the lipid chain is one of the factors that determines the physical phase in which the bilayer will exist at room temperature: longer chain lipids tend to exist in the gel phase, whereas shorter chain lipids tend to exist in the fluid phase. Fig. 4B shows the ppp SFG spectra collected after 37.5 μg of alamethicin was injected into the subphase (~ 1.6 mL) of the bilayer for 60–80 min at pH 6.7. In the fluid phase




Fig. 4. (A) The SFG spectra of the alamethicin after 37.5 μg alamethicin was injected into the subphase (~ 1.6 mL) of d-DMPC/DMPC bilayer for 71 min at pH 6.7. (B) The ppp SFG spectra of the alamethicin after 37.5 μg alamethicin was injected into the subphase (~ 1.6 mL) of the bilayers for 60–80 min at pH 6.7. Top: in a POPC/POPC bilayer; Bottom: in a d-DPPC/DPPC bilayer.

bilayers (Fig. 4A and top spectrum in Fig. 4B), the strong SFG signal of alamethicin is dominated by two peaks at 1635 and 1670 cm^{-1} contributed by 3_{10} -helices and α -helices, respectively. When alamethicin was present in gel-phase lipid bilayers (the lower spectrum in Fig. 4B), only two weak peaks at 1685 and 1720 cm^{-1} were observed. The peak at 1685 cm^{-1} was assigned to antiparallel β -sheets or aggregated strands of peptides (Tamm and Tatulian, 1997, see more details in next section), and the 1720 cm^{-1} signal originates from the bilayer. These results suggest that alamethicin is able to insert into fluid-phase membranes, but that it lies or aggregates on the gel-phase membrane surface and does not have significant insertion into these membranes. These results align well with the results obtained using other analytical tools in the literature (Banerjee et al., 1985; Ionov et al., 2000; Mottamal and Lazaridis, 2006).

We also studied the pH effects on alamethicin in the lipid bilayer using a POPC/POPC bilayer as a model. According to the ppp SFG spectra (not shown) of alamethicin in POPC/POPC bilayer at pH 6.7 and pH 11.9 (the pH was adjusted by adding K_3PO_4), after adjusting the pH to 11.9, the SFG amide I intensity from alamethicin increased by ~ 10 -fold and the peak at ~ 1720 cm^{-1} disappeared. These results suggested that higher pH values, which may affect membrane potential, can induce significant insertion or incorporation of alamethicin into membranes. These results may provide molecular information on the voltage dependence of the alamethicin channels formed in membranes.

3.3.4. SFG studies on β -sheet peptides in membrane bilayers: tachyplesin I

After having discussed the α -helices and 3_{10} helices studied by SFG, we will now present SFG results on another important secondary structure, the β -sheet. Tachyplesin I has been widely used as a model antimicrobial peptide (AMP) that has a β -sheet structure (Katsu et al., 1993; Matsuzaki et al., 1993; Mizuguchi et al., 2003; Nakamura et al., 1988). The antiparallel β -sheet structure is held rigidly together by two intra-strand disulfide bonds. The role of the disulfide bonds has been the focus of several research articles. Decreased antimicrobial activity has been reported when the four cysteines are protected by acetamidomethyl groups (Matsuzaki et al., 1993), but another study, which used linear analogs with cysteines mutated to other residues, indicated that the rigidly held disulfide-bonded β -sheet structure may not be absolutely required for antimicrobial activity (Mizuguchi et al., 2003). The exact mode of action for tachyplesin I is therefore still controversial.

Recently, the Chen group discovered that tachyplesin I induced bilayer structural changes that also exhibited concentration dependence (Chen and Chen, 2006). The SFG results indicated that tachyplesin I was very active in disrupting DPPG bilayers (more details in Section 3.4). To carry out the SFG data analysis on β -sheet structures, Chen et al. first studied tachyplesin I at a polystyrene/peptide solution interface (Chen et al., 2005b; Wang et al., 2005b). Due to the D_2 symmetry of β -sheets (Fig. 5), which differs from that

Table 3
The interaction between alamethicin and different lipid bilayers.

Inner layer	Outer layer	Transition temperature of outer layer lipid ($^{\circ}\text{C}$) [*]	Phase of outer layer lipid at experimental condition	SFG signal
POPC	POPC	-2	Fluid	Very strong
POPC	POPG	-2	Fluid	Very strong
d-DMPC	d-DMPC	23	Fluid	Very strong
d-DMPC	DMPC	23	Fluid	Very strong
d-DPPC	DPPC	41	Gel	Weak
d-DPPG	DPPG	41	Gel	Weak
d-DSPC	DSPC	55	Gel	Weak

^{*} <http://www.avantlipids.com/PhaseTransitionTemperaturesForGlycerophospholipids.html>.

of helical structures, one angle θ is not enough to determine β -sheet interfacial orientation because the twist angle ϕ cannot be random. For D_2 symmetry, SFG measured parameters (nonlinear susceptibility tensor components) can be related to molecular properties of β -sheets (hyperpolarizability tensor components) through orientation parameters (Wang et al., 2005b):

B_1 mode:

$$\chi_{zzz} = 2N_s(\langle \cos \theta \sin \phi \cos \phi \rangle - \langle \cos^3 \theta \sin \phi \cos \phi \rangle)\beta_{abc} \quad (8)$$

$$\chi_{xxz} = \chi_{yyz} = \chi_{xzx} = \chi_{yzy} = \chi_{zxx} = \chi_{zyy} = -N_s(\langle \cos \theta \sin \phi \cos \phi \rangle - \langle \cos^3 \theta \sin \phi \cos \phi \rangle)\beta_{abc} \quad (9)$$

$$\chi_{xyy} = -\chi_{zyx} = -\chi_{yzx} = \chi_{xzy} = 0.5N_s(\langle \sin^2 \theta \cos^2 \phi \rangle - \langle \sin^2 \theta \sin^2 \phi \rangle)\beta_{abc} \quad (10)$$

B_2 mode:

$$\chi_{xzy} = -\chi_{zyx} = -\chi_{yzx} = \chi_{xzy} = 0.5N_s(\langle \cos^2 \theta \rangle - \langle \sin^2 \theta \cos^2 \phi \rangle)\beta_{acb} \quad (11)$$

B_3 mode:

$$\chi_{xzy} = -\chi_{zyx} = -\chi_{yzx} = \chi_{xzy} = -0.5N_s(\langle \cos^2 \theta \rangle - \langle \sin^2 \theta \sin^2 \phi \rangle)\beta_{bca} \quad (12)$$

where N_s is the surface number density of the repeating units of the β -sheet. The standard (or achiral) susceptibility components χ_{xxz} , χ_{yyz} , χ_{xzx} , χ_{yzy} , χ_{zxx} , χ_{zyy} and χ_{zzz} can be obtained by fitting achiral SFG spectra, and chiral tensors χ_{zxy} , χ_{zyx} , χ_{yzx} and χ_{xzy} can be deduced from chiral SFG spectra. The achiral susceptibility tensor elements for the B_2 and B_3 modes have the same form as the B_1 mode, except that β_{abc} should be replaced by β_{acb} and β_{bca} , respectively. The molecular hyperpolarizability components of β -sheets can be calculated from literature values in IR and Raman measurements and confirmed by ab-initio calculations. Therefore, for SFG experiments, orientation parameters such as $\langle \cos^2 \theta \rangle$, $\langle \cos \theta \sin \phi \cos \phi \rangle$ and $\langle \sin^2 \theta \cos^2 \phi \rangle$ of β -sheets can be measured.

The Chen group successfully collected SFG amide I signals from tachyplesin I at the polystyrene/solution interface (Fig. 6A) (Chen et al., 2005b). The fitting results showed three major peaks at 1645, 1664 and 1688 cm^{-1} , respectively. A large number of reports have stated that amide I signals at 1688 and 1633 cm^{-1} can be ascribed to the B_1/B_3 and B_2 modes of antiparallel β -sheets, respectively (Barth and Zscherp, 2002; Hilario et al., 2003; Krimm and Bandekar, 1986; Vass et al., 2003). The 1664 and 1645 cm^{-1} peaks in the tachyplesin I SFG spectra are due to turns, random structures or a combination thereof (Barth and Zscherp, 2002; Krimm and Bandekar, 1986; Vass et al., 2003). Following the addition of dithi-




Fig. 5. Molecular coordinate for an antiparallel β -sheet. Reproduced with permission from *Proc. Natl. Acad. Sci. USA*, 2005, 102, 4978–4983. Copyright 2005, The National Academy of Sciences of the USA.

othreitol (DTT), the 1688 cm^{-1} peak disappeared, confirming that this peak was due to β -sheet structure of tachyplesin I at the interface (Fig. 6B). The addition of DTT broke the two disulfide bonds in tachyplesin I, which are essential for it to maintain its β -sheet structure (Li et al., 1998; Matsuzaki et al., 1993). Furthermore, Chen and colleagues also detected very strong SFG chiral spp and psp spectra directly from the polystyrene/tachyplesin I solution interface (Fig. 6C) (Wang et al., 2005b). Their intensities were comparable to those in the ssp spectrum, but with distinct spectral features. For the spp and psp spectra, only two peaks at 1635 and 1685 cm^{-1} were observed from the spectral fitting results, showing that SFG chiral signals were dominated by contributions from β -sheet structures. Such signals can provide more measurements to determine β -sheet orientation.

Chen and colleagues have also collected SFG amide I signals from tachyplesin I in a DPPG/DPPG bilayer. These spectra indicated the presence of antiparallel β -sheet structure, with a dominant band around 1685 cm^{-1} (Fig. 6D) (Chen and Chen, 2006). Some differences in the spectral features in these spectra compared to the spectra in Fig. 6A and C indicated that tachyplesin I adopted a different conformation at the bilayer interface compared to that adsorbed onto a polystyrene surface (Chen et al., 2005b; Wang et al., 2005b). Chen and colleagues observed that the addition of DTT to the solution caused no changes to the spectrum of tachyplesin I already adsorbed onto a bilayer, as opposed to the changes observed when tachyplesin I was adsorbed onto a polystyrene surface, where addition of DTT led to the disappearance of the 1685 cm^{-1} peak of tachyplesin I (Chen and Chen, 2006; Chen et al., 2005b; Wang et al., 2005b). It appeared that the membrane, in contrast to the polystyrene surface, offered protection to the β -sheet structure, either by shielding the interfacial tachyplesin I from the reducing agent or by inducing the β -sheet structure even without the presence of the disulfide bonds. Currently we are investigating the detailed orientation of tachyplesin I in the lipid bilayer using the methods we presented above and the results will be reported in the future.

3.4. Real time monitoring of bilayer perturbation induced by peptides/proteins

In addition to the structural studies of membrane proteins and peptides in lipid bilayers, time-dependent SFG studies have also been applied to monitor the kinetics of changes in bilayers themselves, or as they are interacting with membrane proteins/peptides (Liu and Conboy, 2004a; Chen and Chen, 2006; Chen et al., 2007b). The Conboy group deduced the flip-flop rate for a planar supported lipid bilayer (DSPC/DSPC- d_{83}) by measuring the time-dependent SFG intensity of terminal CH_3 symmetric modes at various temperatures (Liu and Conboy, 2004a). Chen et al. applied SFG to monitor the time-dependent interactions between antimicrobial peptides and lipid bilayers (Chen and Chen, 2006; Chen et al., 2007b). Different modes of actions were observed between lipid bilayers and different molecules such as melittin, tachyplesin I, δ -magainin 2, MSI-843, and a synthetic antibacterial oligomer (Chen and Chen, 2006).

For example, Chen et al. (2007b) monitored the time-dependent and concentration-dependent disruption of a substrate supported lipid bilayer by melittin. In this research, both a symmetric d-DPPG/d-DPPG bilayer and an asymmetric d-DPPG/DPPG bilayer were used. It was found that the extent and kinetics of the bilayer disruption induced by melittin were greatly affected by the peptide concentration. For a symmetric d-DPPG/d-DPPG bilayer, Chen et al. monitored the intensity of the CD_3 symmetric stretching peak at 2070 cm^{-1} as a function of time after injecting melittin at time 0 s (Fig. 7). At high melittin solution concentrations (2.34–7.8 μM), the peak intensity reached a maximum immediately after the injection of melittin, and then decreased. The initial signal in-




Fig. 6. SFG spectra and fitting results for 0.1 mg/mL tachyplesin I adsorbed at solution/polystyrene interface with squares representing the actual spectrum, dotted line the fitted spectrum and solid lines the component peaks used to fit the spectra. (A) SFG spectra before contacting with 10 mM DTT; (B) SFG spectra after contacting with 10 mM DTT. Reproduced with permission from *Langmuir* 2005, 21, 2662–2664. Copyright 2005, American Chemical Society. (C) SFG spectra of adsorbed tachyplesin I at the polystyrene/solution interface with different polarization combinations. Reproduced with permission from *Proc. Natl. Acad. Sci. USA*. 2005, 102, 4978–4983. Copyright 2005, The National Academy of Sciences of the USA. (D) Amide I SFG spectrum of tachyplesin I (5.6 $\mu\text{g/mL}$) adsorbed onto a DPPG/DPPG bilayer collected using ssp polarization combination. Reproduced with permission from *Biochim. Biophys. Acta*. 2006, 1758, 1257–1273. Copyright 2006, Elsevier B.V.

crease was due to the fact that the first leaflet was disrupted more substantially initially and the inversion symmetry of the bilayer was broken. Because the second leaflet was disrupted shortly after, a new inversion symmetry was created causing the SFG signal to decrease. For lower melittin concentrations, the disruption process was much slower. For melittin concentrations at 0.78 μM , the CD_3 peak intensity underwent a gradual increase after the initial increase/decrease pattern. For concentrations even lower (e.g., 0.156 μM), the initial intensity increase process took a much longer time to complete, showing that the disruption process that creates a break in the inversion symmetry was much slower than at higher concentrations of melittin.

For asymmetric d-DPPG/DPPG bilayers (Fig. 8), Chen et al. monitored the SFG intensity of CD_3 and CH_3 symmetric stretching peaks at 2070 and 2875 cm^{-1} , respectively. It was found that the d-DPPG/DPPG bilayer had a similar disruption trend to that observed in the d-DPPG/d-DPPG bilayer. The spectral changes occurred very quickly and then the signal became stable rapidly at higher concentrations, while a much longer time was required for the intensity to become stable for lower melittin concentrations. When 0.78 μM or higher melittin solution concentrations were used, very weak final signals from both leaflets were observed after the peptide-lipid interactions. At a solution concentration of 0.156 μM or

lower, strong SFG signals from the proximal d-DPPG leaflet could still be observed after a prolonged period of time. This experiment also indicated that the lipid bilayer was disrupted at a much faster rate at a higher peptide concentration. In addition, Fig. 8 shows that melittin disrupted the two leaflets at different rates: the distal leaflet was disrupted first, before the disruption of the proximal leaflet.

The Chen group also detected time-dependent SFG signals from the asymmetric d-DPPG-DPPG bilayer/tachyplesin I solution interface with a solution concentration of 5.6 $\mu\text{g/mL}$ (Chen and Chen, 2006). However, unlike the time delay between the onset of SFG C-H and C-D signal decrease observed for melittin, the signal decreases for both leaflets, were coincident with each other. This indicated that there were possibly different interaction patterns among various antimicrobial peptides. With tachyplesin I, the insertion probably occurred more readily and both leaflets were disrupted immediately and at the same rate. Time-dependent interactions between DTT-treated tachyplesin I and a dDPPG/DPPG bilayer were different from those observed for the untreated tachyplesin I. The DTT-treated tachyplesin I induced a decrease in C-D/C-H stretching mode similar to that observed for 0.44 $\mu\text{g/mL}$ of melittin. This shows that the DTT-treated tachyplesin I was actually more potent in causing the signal intensity decrease during




Fig. 7. Signal intensity change at 2070 cm^{-1} monitoring melittin's interaction with a dDPPG/dDPPG bilayer on a CaF_2 prism. Melittin stock solution was injected into the subphase of the bilayer at time 0 s. Four solution concentrations were used and dramatically different kinetics were observed. Reproduced with permission from *Biophys. J.* 2007, 93, 866–875. Copyright 2007, the Biophysical Society.




Fig. 8. Signal intensity changes at 2070 and 2875 cm^{-1} monitoring melittin's interaction with dDPPG/DPPG bilayers on a CaF_2 prism. Melittin stock solution was injected into the subphase of bilayer at time 200 seconds. Four solution concentrations were used and dramatically different kinetics were observed. Reproduced with permission from *Biophys. J.* 2007, 93, 866–875. Copyright 2007, the Biophysical Society.

the initial period of interaction, but the untreated tachyplesin I could induce a more thorough destabilizing effect, which may be the critical factor in killing microbes.

4. Summary

In this paper, we summarized the recent applications of SFG studies on lipid monolayers, HBMs and substrate-supported lipid bilayers, as well as the interactions between peptides/proteins

and lipid monolayers/bilayers, and bilayer perturbation induced by peptides/proteins. We introduced early SFG studies that focused on the C–H stretching frequency region and the more recent SFG studies in the amide I region investigated by our group. As discussed above, the peptide/protein secondary structures (α -helical, 3_{10} -helical and β -sheet) can be determined and their orientation information can be deduced. We discussed examples of studies on the interaction of peptides/proteins (melittin, G proteins, alamethicin, and tachyplesin I) and lipid bilayers using SFG. These

studies showed that SFG is a powerful technique to study the membrane-related peptides and proteins and that SFG can provide a unique understanding of the interactions between a lipid monolayer/bilayer and peptides/proteins without any exogenous labeling in real time and *in situ*.

Acknowledgments

This research is supported by the National Institute of Health (1R01GM081655-01A2) and the Office of Naval Research (N00014-02-1-0832 and N00014-08-1-1211). SVLC acknowledges the Molecular Biophysics Training Grant from the University of Michigan.

Appendix A

The intensity of the sum frequency signal is proportional to the square of the vibration's second-order nonlinear susceptibility $\chi_{eff}^{(2)}$:

$$I(\omega_{SF}) = \frac{8\pi^3 \omega_{SF}^2 \sec^2 \beta}{c^3 n_1(\omega_{SF}) n_1(\omega_{Vis}) n_1(\omega_{IR})} |\chi_{eff}^{(2)}|^2 I_1(\omega_{Vis}) I_2(\omega_{IR}) \quad (A.1)$$

where $n_i(\omega)$ is the refractive index of medium i at frequency ω , β is the reflection angle of the sum-frequency field, $I_1(\omega_{Vis})$ and $I_2(\omega_{IR})$ are the intensities of the two input fields. The effective nonlinear susceptibility $\chi_{eff}^{(2)}$ is:

$$\chi_{eff}^{(2)} = [\hat{e}(\omega_{SF}) \cdot L(\omega_{SF})] \cdot \chi^{(2)} : [L(\omega_{Vis}) \cdot \hat{e}(\omega_{Vis})][L(\omega_{IR}) \cdot \hat{e}(\omega_{IR})] \quad (A.2)$$

with $\hat{e}(\omega)$ being the unit polarization vector and $L(\omega)$ the Fresnel factor at frequency ω . Details about Fresnel factors have been discussed extensively and will not be repeated here (Zhuang et al., 1999).

For Visible–IR SFG, if the IR frequency (ω_{IR}) is near vibrational resonances, the effective surface nonlinear susceptibility $\chi_R^{(2)}$ can be enhanced. The frequency dependence of $\chi_{eff}^{(2)}$ can be written as:

$$\chi_{eff}^{(2)}(\omega) = \chi_{NR}^{(2)} + \chi_R^{(2)} = \chi_{NR}^{(2)} + \sum_q \frac{A_q}{\omega - \omega_q + i\Gamma_q} \quad (A.3)$$

where A_q , ω_q , and Γ_q are the strength, resonant frequency, and damping coefficient of the vibrational mode q , respectively, $\chi_{NR}^{(2)}$ is the nonresonant background, and $\chi_R^{(2)}$ is the resonant contribution (Gautam and Dhinojwala, 2001; Hirose et al., 1992a,b, 1993; Kim and Somorjai, 2003; Shen, 1984; Wang et al., 2001).

The components of $\chi_{eff}^{(2)}$ can be probed by using different polarization combinations of the input and output laser beams, such as ssp (s-polarized SFG signal, s-polarized visible beam, p-polarized IR beam), sps, pss and ppp. Each measured polarization combination is related to the different components of $\chi^{(2)}$ defined in the lab coordinate system. For an azimuthally isotropic interface only four independent components of $\chi^{(2)}$ are nonzero from Eq. (A.2). If the lab coordinate system is defined as the z-axis being along the surface normal and the x-axis being in the incident plane (Fig. 1A), these four components will have the following properties: $\chi_{xxx} = \chi_{yyz}$, $\chi_{xzx} = \chi_{zyy}$, $\chi_{zxx} = \chi_{zyy}$, and χ_{zzz} . The $\chi_{eff}^{(2)}$ expressions under ssp, sps, pss and ppp polarization combinations are then (Miranda and Shen, 1999; Shen, 1984; Wang et al., 2001; Zhuang et al., 1999):

$$\chi_{eff,ssp}^{(2)} = L_{yy}(\omega_{SF}) L_{yy}(\omega_{Vis}) L_{zz}(\omega_{IR}) \sin \beta_{IR} \chi_{yyz}^{(2)} \quad (A.4)$$

$$\chi_{eff,sps}^{(2)} = L_{yy}(\omega_{SF}) L_{zz}(\omega_{Vis}) L_{yy}(\omega_{IR}) \sin \beta_{Vis} \chi_{zyy}^{(2)} \quad (A.5)$$

$$\chi_{eff,pss}^{(2)} = L_{zz}(\omega_{SF}) L_{yy}(\omega_{Vis}) L_{yy}(\omega_{IR}) \sin \beta_{SF} \chi_{zyy}^{(2)} \quad (A.6)$$

$$\begin{aligned} \chi_{eff,ppp}^{(2)} = & -L_{xx}(\omega_{SF}) L_{xx}(\omega_{Vis}) L_{zz}(\omega_{IR}) \cos \beta_{SF} \cos \beta_{Vis} \sin \beta_{IR} \chi_{xxx}^{(2)} \\ & - L_{xx}(\omega_{SF}) L_{zz}(\omega_{Vis}) L_{xx}(\omega_{IR}) \cos \beta_{SF} \sin \beta_{Vis} \cos \beta_{IR} \chi_{xxx}^{(2)} \\ & + L_{zz}(\omega_{SF}) L_{xx}(\omega_{Vis}) L_{xx}(\omega_{IR}) \sin \beta_{SF} \cos \beta_{Vis} \cos \beta_{IR} \chi_{zxx}^{(2)} \\ & + L_{zz}(\omega_{SF}) L_{zz}(\omega_{Vis}) L_{zz}(\omega_{IR}) \sin \beta_{SF} \sin \beta_{Vis} \sin \beta_{IR} \chi_{zzz}^{(2)} \quad (A.7) \end{aligned}$$

where β_{SF} , β_{Vis} and β_{IR} are the angles between the surface normal and the sum frequency beam, the input visible beam, and the input IR beam, respectively; and L_i 's ($i = x, y$ or z) are the Fresnel coefficients.

The tensor components $\chi_{ijk}^{(2)}$ of $\chi^{(2)}$ defined in the lab coordinate system are macroscopic properties of a surface or interface. They can be directly related to the microscopic hyperpolarizability tensor components $\beta_{\lambda\mu\nu}$ of molecules on the surface or interface. In certain papers, microscopic hyperpolarizability tensor components have also been referred to as $\alpha_{\lambda\mu\nu}^{(2)}$. In short, each $\chi_{ijk}^{(2)}$ is a transformation of individual microscopic hyperpolarizability tensor components $\beta_{\lambda\mu\nu}$ in the molecular coordinate system to its macroscopic quantity of the ensemble average over all possible orientations in the lab frame (Miranda and Shen, 1999; Shen, 1984; Wang et al., 2001; Zhuang et al., 1999):

$$\chi_{ijk}^{(2)} = N \sum_{\lambda,\mu,\nu} \langle (\hat{i} \cdot \hat{\lambda})(\hat{j} \cdot \hat{\mu})(\hat{k} \cdot \hat{\nu}) \rangle \beta_{\lambda\mu\nu}, \quad (A.8)$$

where N is the number density of molecules and the brackets denote averaging over the molecular orientation distribution. Both $\chi^{(2)}$ and $\beta_{\lambda\mu\nu}$ are third rank tensors with 27 elements whose values are functions of the frequencies of the three beams involved in the SFG process. For the SFG technique we are discussing here, $\beta_{\lambda\mu\nu}$ is a product of the IR transition dipole moment component and Raman polarizability component, which can be measured experimentally or calculated. Therefore, by measuring $\chi_{ijk}^{(2)}$ in SFG experiments, the orientation and the number density of interfacial molecules in a sample may be deduced.

References

- Alessandrini, A., Facci, P., 2005. AFM: a versatile tool in biophysics. *Meas. Sci. Technol.* 16, R65–R92.
- Allen, H.C., Raymond, E.A., Richmond, G.L., 2000. Non-linear vibrational sum frequency spectroscopy of atmospherically relevant molecules at aqueous solution surfaces. *Curr. Opin. Colloid. Interface Sci.* 5, 74–80.
- Anderson, N.A., Richter, L.J., Stephenson, J.C., Briggman, K.A., 2004. Protein deformation of lipid hybrid bilayer membranes studied by sum frequency generation vibrational spectroscopy. *Langmuir* 20, 8961–8965.
- Anderson, N.A., Richter, L.J., Stephenson, J.C., Briggman, K.A., 2006. Determination of lipid phase transition temperatures in hybrid bilayer membranes. *Langmuir* 22, 8333–8336.
- Anderson, N.A., Richter, L.J., Stephenson, J.C., Briggman, K.A., 2007. Characterization and control of lipid layer fluidity in hybrid bilayer membranes. *J. Am. Chem. Soc.* 129, 2094–2100.
- Andronesi, O.C., Becker, S., Seidel, K., Heise, H., Young, H.S., Baldus, M., 2005. Determination of membrane protein structure and dynamics by magic-angle-spinning solid-state NMR spectroscopy. *J. Am. Chem. Soc.* 127, 12965–12974.
- Anglin, T.C., Liu, J., Conboy, J.C., 2007. Facile lipid flip-flop in a phospholipid bilayer induced by gramicidin A measured by sum-frequency vibrational spectroscopy. *Biophys. J.* 92, L1–L3.
- Anglin, T.C., Conboy, J.C., 2008. Lateral pressure dependence of the phospholipid transmembrane diffusion rate in planar-supported lipid bilayers. *Biophys. J.* 95, 186–193.
- Archer, S.J., Ellena, J.F., Cafiso, D.S., 1991. Dynamics and aggregation of the peptide ion channel alamethicin. *Biophys. J.* 60, 389–398.
- Bader, R., Lerch, M., Zerbe, O., 2003. NMR of membrane-associated peptides and proteins. *Methods and Principles in Medicinal Chemistry: BioNMR in Drug Research*, vol. 16. Wiley, Weinheim, pp. 95–120.
- Bagatoli, L.A., 2006. To see or not to see: lateral organization of biological membranes and fluorescence microscopy. *Biochim. Biophys. Acta* 1758, 1541–1556.
- Bain, C.D., 1995. Sum-frequency vibrational spectroscopy of the solid–liquid interface. *J. Chem. Soc. Faraday Trans.* 91, 1281–1296.
- Bak, M., Baywater, R.P., Hohwy, M., Thomsen, J.K., Adelhorst, K., Jakobsen, H.J., Sorensen, O.W., Nielsen, N.C., 2001. Conformation of alamethicin in oriented phospholipid bilayers determined by N-15 solid-state nuclear magnetic resonance. *Biophys. J.* 81, 1684–1698.

- Baldelli, S., 2008. Surface structure at the ionic liquid-electrified metal interface. *Acc. Chem. Res.* 41, 421–431.
- Banerjee, U., Zidovetzki, R., Birge, R.R., Chan, S.I., 1985. Interaction of alamethicin with lecithin bilayers—a P-31 and H-2 NMR-study. *Biochemistry* 24, 7621–7627.
- Barranger-Mathys, M., Cafiso, D.S., 1994. Collisions between helical peptides in membranes monitored using electron paramagnetic resonance: evidence that alamethicin is monomeric in the absence of a membrane potential. *Biophys. J.* 67, 172–176.
- Barth, A., Zscherp, C., 2002. What vibrations tell us about proteins. *Quart. Rev. Biophys.* 35, 369–430.
- Bechinger, B., 1999. The structure, dynamics and orientation of antimicrobial peptides in membranes by multidimensional solid-state NMR spectroscopy. *Biochim. Biophys. Acta* 1462, 157–183.
- Bechinger, B., Aisenbrey, C., Bertani, P., 2004. The alignment, structure and dynamics of membrane-associated polypeptides by solid-state NMR spectroscopy. *Biochim. Biophys. Acta* 1666, 190–204.
- Belkin, M.A., Shen, Y.R., 2005. Non-linear optical spectroscopy as a novel probe for molecular chirality. *Int. Rev. Phys. Chem.* 24, 257–299.
- Beseničar, M., Maček, P., Lakey, J.H., Anderluh, G., 2006. Surface plasmon resonance in protein-membrane interactions. *Chem. Phys. Lipids* 141, 169–178.
- Bonn, M., Roke, S., Berg, O., Juurlink, L.B.F., Stamouli, A., Muller, M., 2004. A molecular view of cholesterol-induced condensation in a lipid monolayer. *J. Phys. Chem. B* 108, 19083–19085.
- Brasseur, R., Deleu, M., Mingeot-Leclercq, M.P., Francius, G., Dufréne, Y.F., 2008. Probing peptide-membrane interactions using AFM. *Surf. Interface Anal.* 40, 151–156.
- Briggman, K.A., Stephenson, J.C., Wallace, W.E., Richter, L.J., 2001. Absolute molecular orientational distribution of the polystyrene surface. *J. Phys. Chem. B* 2001 (105), 2785–2791.
- Buck, M., Himmelhaus, M., 2001. Vibrational spectroscopy of interfaces by infrared-visible sum frequency generation. *J. Vac. Sci. Technol. A* 19, 2717–2736.
- Cabiaux, V., 2004. PH-sensitive toxins: interactions with membrane bilayers and application to drug delivery. *Adv. Drug Del. Rev.* 56, 987–997.
- Cabrera-Vera, T.M., Vanhauwe, J., Thomas, T.O., Medkova, M., Preininger, A., Mazzoni, M.R., Hamm, H.E., 2003. Insights into G protein structure, function, and regulation. *Endocr. Rev.* 24, 765–781.
- Cafiso, D.S., 1994. Alamethicin—a peptide model for voltage gating and protein membrane interactions. *Annu. Rev. Biophys. Biomol. Struct.* 23, 141–165.
- Castellana, E.T., Cremer, P.S., 2006. Solid supported lipid bilayers: from biophysical studies to sensor design. *Surf. Sci. Rep.* 61, 429–444.
- Chen, F.Y., Lee, M.T., Huang, H.W., 2002a. Sigmoidal concentration dependence of antimicrobial peptide activities: a case study on alamethicin. *Biophys. J.* 82, 908–914.
- Chen, X., Clarke, M.L., Wang, J., Chen, Z., 2005a. Sum frequency generation vibrational spectroscopy studies on molecular conformation and orientation of biological molecules at interfaces. *Int. J. Mod. Phys. B.* 19, 691–713.
- Chen, X., Wang, J., Sniadecki, J., Even, M.A., Chen, Z., 2005b. Probing α -helical and β -sheet structures of peptides at solid/liquid interfaces with SFG. *Langmuir* 21, 2662–2664.
- Chen, X., Chen, Z., 2006. SFG studies on interactions between antimicrobial peptides and supported lipid bilayers. *Biochim. Biophys. Acta* 1758, 1257–1273.
- Chen, X., Tang, H., Even, M.A., Wang, J., Tew, G.N., Chen, Z., 2006. Observing a molecular knife at work. *J. Am. Chem. Soc.* 128, 2711–2714.
- Chen, X., Boughton, A.P., Tesmer, J.J.G., Chen, Z., 2007a. In situ investigation of heterotrimeric G protein β subunit binding and orientation on membrane bilayers. *J. Am. Chem. Soc.* 129, 12658–12659.
- Chen, X., Wang, J., Kristalyn, C.B., Chen, Z., 2007b. Real-time structural investigation of a lipid bilayer during its interaction with melittin using sum frequency generation vibrational spectroscopy. *Biophys. J.* 93, 866–875.
- Chen, X., Wang, J., Boughton, A.P., Kristalyn, C.B., Chen, Z., 2007c. Multiple orientation of melittin inside a single lipid bilayer determined by combined vibrational spectroscopic studies. *J. Am. Chem. Soc.* 129, 1420–1427.
- Chen, Z., 2007. Understanding surfaces and buried interfaces of polymer materials at the molecular level using sum frequency generation vibrational spectroscopy. *Polym. Int.* 56, 577–587.
- Chen, Z., Shen, Y.R., Somorjai, G.A., 2002b. Studies of polymer surfaces by sum frequency generation vibrational spectroscopy. *Annu. Rev. Phys. Chem.* 53, 437–465.
- Clarke, M.L., Wang, J., Chen, Z., 2005. Conformational changes of fibrinogen after adsorption. *J. Phys. Chem. B* 109, 22027–22035.
- Devanathan, S., Salamon, Z., Lindblom, G., Grobner, G., Tollin, G., 2006. Effects of sphingomyelin, cholesterol and zinc ions on the binding, insertion and aggregation of the amyloid β_{1-40} peptide in solid-supported lipid bilayers. *FEBS J.* 273, 1389–1402.
- Doyle, A.W., Fick, J., Himmelhaus, M., Eck, W., Graziani, I., Prudovsky, I., Grunze, M., Maciag, T., Neivandt, D.J., 2004. Protein deformation of lipid hybrid bilayer membranes studied by sum frequency generation vibrational spectroscopy. *Langmuir* 20, 8961–8965.
- Dreesen, L., Humbert, C., Sartenaer, Y., Caudano, Y., Volcke, C., Mani, A.A., Peremans, A., Thiry, P.A., Hanique, S., Frere, J.M., 2004a. Electronic and molecular properties of an adsorbed protein monolayer probed by two-color sum-frequency generation spectroscopy. *Langmuir* 20, 7201–7207.
- Dreesen, L., Sartenaer, Y., Humbert, C., Mani, A.A., Methivier, C., Pradier, C.M., Thiry, P.A., Peremans, A., 2004b. Probing ligand-protein recognition with sum-frequency generation spectroscopy: the avidin-biotin case. *Chem. Phys. Chem.* 5, 1719–1725.
- Duclohier, H., 2004. Helical kink and channel behaviour: a comparative study with the heptaibols alamethicin, trichothoxin and antiameobin. *Eur. Biophys. J.* 33, 169–174.
- Duclohier, H., Wroblewski, H., 2001. Voltage-dependent pore formation and antimicrobial activity by alamethicin and analogues. *J. Membr. Biol.* 184, 1–12.
- Dürr, U.H.N., Yamamoto, K., Im, S.-C., Waskell, L., Ramamoorthy, A., 2007. Solid-state NMR reveals structural and dynamical properties of a membrane-anchored electron-carrier protein, cytochrome *b5*. *J. Am. Chem. Soc.* 129, 6670–6671.
- Dvinskikh, S.V., Dürr, U.H.N., Yamamoto, K., Ramamoorthy, A., 2007. High-resolution 2D NMR spectroscopy of bicelles to measure the membrane interaction of ligands. *J. Am. Chem. Soc.* 129, 794–802.
- Dwivedi, A.M., Krimm, S., 1984. Vibrational analysis of peptides, polypeptides, and proteins. 24. conformation of poly(α -aminoisobutyric acid). *Biopolymers* 23, 2025–2065.
- Eisenenthal, K.B., 1992. Equilibrium and dynamic processes at interfaces by second harmonic and sum frequency generation. *Annu. Rev. Phys. Chem.* 43, 627–661.
- Engel, A., Gaub, H.E., 2008. Structure and mechanics of membrane proteins. *Annu. Rev. Biochem.* 77, 127–148.
- Evans-Nguyen, K.M., Fuierer, R.R., Fitchett, B.D., Tolles, L.R., Conboy, J.C., Schoenfish, M.H., 2006. Changes in adsorbed fibrinogen upon conversion to fibrin. *Langmuir* 22, 5115–5121.
- Faiss, S., Kastl, K., Janshoff, A., Steinem, C., 2008. Formation of irreversibly bound annexin A1 protein domains on POPC/POPS solid supported membranes. *Biochim. Biophys. Acta* 1778, 1601–1610.
- Fernández, C., Adeishvili, K., Wuthrich, K., 2001. Transverse relaxation-optimized NMR spectroscopy with the outer membrane protein OmpX dihexanoyl phosphatidylcholine micelles. *Proc. Natl. Acad. Sci. USA* 98, 2358–2363.
- Fourkas, J.T., Walker, R.A., Can, S.Z., Gershgoren, E., 2007. Effects of reorientation in vibrational sum-frequency spectroscopy. *J. Phys. Chem. C* 111, 8902–8915.
- Fox, R.O., Richards, F.M., 1982. A voltage-gated ion channel model inferred from the crystal-structure of alamethicin at 1.5-Å resolution. *Nature* 300, 325–330.
- Fragneto-Cusani, G., 2001. Neutron reflectivity at the solid/liquid interface: examples of applications in biophysics. *J. Phys. Condens. Matter* 13, 4973–4989.
- Gaudet, R., Savage, J.R., McLaughlin, J.N., Willardson, B.M., Sigler, P.B., 1999. A molecular mechanism for the phosphorylation-dependent regulation of heterotrimeric G proteins by phosphodiacyl. *Mol. Cell* 3, 649–660.
- Gautam, K.S., Schwab, A.D., Dhinojwala, A., Zhang, D., Dougal, S.M., Yeganeh, M.S., 2000. Molecular structure of polystyrene at air/polymer and solid/polymer interfaces. *Phys. Rev. Lett.* 85, 3854–3857.
- Gautam, K.S., Dhinojwala, A., 2001. Molecular structure of hydrophobic alkyl side chains at comb polymer-air interface. *Macromolecules* 34, 1137–1139.
- Gautam, K.S., Dhinojwala, A., 2002. Melting at alkyl side chain comb polymer interfaces. *Phys. Rev. Lett.* 88, 145501.
- Gopalakrishnan, S., Liu, D.F., Allen, H.C., Kuo, M., Shultz, M.J., 2006. Vibrational spectroscopic studies of aqueous interfaces: salts, acids, bases, and nanodrops. *Chem. Rev.* 106, 1155–1175.
- Gracias, D.H., Chen, Z., Shen, Y.R., Somorjai, G.A., 1999. Molecular characterization of polymer and polymer blend surfaces. Combined sum frequency generation surface vibrational spectroscopy and scanning force microscopy studies. *Acc. Chem. Res.* 32, 930–940.
- Guyotsonnest, P., Hunt, J.H., Shen, Y.R., 1987. Sum-frequency vibrational spectroscopy of a Langmuir film—study of molecular-orientation of a two-dimensional system. *Phys. Rev. Lett.* 59, 1597–1600.
- Haas, H., Steitz, R., Fasano, A., Liuzzi, G.M., Polverini, E., Cavatorta, P., Riccio, P., 2007. Laminar order within Langmuir-Blodgett multilayers from phospholipid and Myelin basic protein: a neutron reflectivity study. *Langmuir* 23, 8491–8496.
- Hall, J.E., Vodyanov, I., Balasubramanian, T.M., Marshall, G.R., 1984. Alamethicin—a rich model for channel behavior. *Biophys. J.* 45, 233–247.
- Haris, P.L., Chapman, D., 1988. Fourier-transform infrared-spectra of the polypeptide alamethicin and a possible structural similarity with bacteriorhodopsin. *Biochim. Biophys. Acta* 943, 375–380.
- Haris, P.L., Molle, G., Duclohier, H., 2004. Conformational changes in alamethicin associated with substitution of its α -methylalanines with leucines: a FTIR spectroscopic analysis and correlation with channel kinetics. *Biophys. J.* 86, 248–253.
- Harper, K.L., Allen, H.C., 2007. Competition between DPPC and SDS at the air-aqueous interface. *Langmuir* 23, 8925–8931.
- Heberle, F.A., Buboltz, J.T., Stringer, D., Feigenson, G.W., 2005. Fluorescence methods to detect phase boundaries in lipid bilayer mixtures. *Biochim. Biophys. Acta* 1746, 186–192.
- Hilario, J., Kubelka, J., Keiderling, T.A., 2003. Optical spectroscopic investigations of model β -sheet hairpins in aqueous solution. *J. Am. Chem. Soc.* 125, 7562–7574.
- Hirose, C., Akamatsu, N., Domen, K., 1992a. Formulas for the analysis of the surface SFG spectrum and transformation coefficients of cartesian SFG tensor components. *Appl. Spectrosc.* 46, 1051–1072.
- Hirose, C., Akamatsu, N., Domen, K., 1992b. Formulas for the analysis of surface sum-frequency generation spectrum by CH stretching modes of methyl and methylene groups. *J. Chem. Phys.* 96, 997–1004.
- Hirose, C., Yamamoto, H., Akamatsu, N., Domen, K., 1993. Orientation analysis by simulation of vibrational sum-frequency generation spectrum—CH stretching bands of the methyl-group. *J. Phys. Chem.* 97, 10064–10069.

- Holman, J., Ye, S., Neivandt, D.J., Davies, P.B., 2004. Studying nanoparticle-induced structural changes within fatty acid multilayer films using sum frequency generation vibrational spectroscopy. *J. Am. Chem. Soc.* 126, 14322–14323.
- Hopkins, A.J., McFearin, C.L., Richmond, G.L., 2005. Investigations of the solid-liquid interface with vibrational sum-frequency spectroscopy. *Curr. Opin. Solid State Mater. Sci.* 9, 19–27.
- Huang, H.W., Wu, Y., 1991. Lipid-alamethicin interactions influence alamethicin orientation. *Biophys. J.* 60, 1079–1087.
- Humbert, C., Dreesen, L., Sartenaer, Y., Peremans, A., Thiry, P.A., Volcke, C., 2006. On the protoporphyrin monolayers conformation. *Chem. Phys. Chem.* 7, 569–571.
- Hunt, J.H., Guyot-Sionnest, P., Shen, Y.R., 1987. Observation of C–H stretching vibrations of monolayers of molecules optical sum frequency generation. *Chem. Phys. Lett.* 133, 189–192.
- Ionov, R., El-Abed, A., Angelova, A., Goldmann, M., Peretti, P., 2000. Asymmetrical ion-channel model inferred from two-dimensional crystallization of a peptide antibiotic. *Biophys. J.* 78, 3026–3035.
- Iwahashi, T., Miyamae, T., Kanai, K., Seki, K., Kim, D., Ouchi, Y., 2008. Anion configuration at the air/liquid interface of ionic liquid [bmim]OTf studied by sum-frequency generation spectroscopy. *J. Phys. Chem. B* 112, 11936–11941.
- Johnston, L.J., 2007. Nanoscale imaging of domains in supported lipid membranes. *Langmuir* 23, 5886–5895.
- Kalb, E., Frey, S., Tamm, L.K., 1992. Formation of supported planar bilayers by fusion of vesicles to supported phospholipid monolayers. *Biochim. Biophys. Acta* 1103, 307–316.
- Katsaras, J., Gutberlet, T., 2001. *Lipid Bilayers: Structure and Interactions*. Springer, New York.
- Katsu, T., Nakao, S., Iwanaga, S., 1993. Mode of action of an antimicrobial peptide, tachyplesin I, on biomembranes. *Biol. Pharm. Bull.* 16, 178–181.
- Kennedy, D.F., Crisma, M., Toniolo, C., Chapman, D., 1991. Studies of peptides forming 310- and α -helices and β -bend ribbon structures in organic solution and in model biomembranes by Fourier transform infrared spectroscopy. *Biochemistry* 30, 6541–6548.
- Kessel, A., Cafiso, D.S., Ben-Tal, N., 2000. Continuum solvent model calculations of alamethicin-membrane interactions: thermodynamic aspects. *Biophys. J.* 78, 571–583.
- Kim, J., Cremer, P.S., 2001. Elucidating changes in interfacial water structure upon protein adsorption. *Chem. Phys. Chem.* 2, 543–546.
- Kim, J., Kim, G., Cremer, P.S., 2001. Investigations of water structure at the solid/liquid interface in the presence of supported lipid bilayers by vibrational sum frequency spectroscopy. *Langmuir* 17, 7255–7260.
- Kim, J., Somorjai, G.A., 2003. Molecular packing of lysozyme, fibrinogen, and bovine serum albumin on hydrophilic and hydrophobic surfaces studied by infrared-visible sum frequency generation and fluorescence microscopy. *J. Am. Chem. Soc.* 125, 3150–3158.
- Kim, G., Gurau, M.C., Kim, J., Cremer, P.S., 2002. Investigations of lysozyme adsorption at the air/water and quartz/water interfaces by vibrational sum frequency spectroscopy. *Langmuir* 18, 2807–2811.
- Kim, G., Gurau, M.C., Lim, S.M., Cremer, P.S., 2003. Investigations of the orientation of a membrane peptide by sum frequency spectroscopy. *J. Phys. Chem. B* 107, 1403–1409.
- Kim, C., Gurau, M.C., Cremer, P.S., Yu, H., 2008. Chain conformation of poly(dimethyl siloxane) at the air/water interface by sum frequency generation. *Langmuir* 24, 10155–10160.
- Knoesen, A., Pakalnis, S., Wang, M., Wise, W.D., Lee, N., Frank, C.W., 2004. Sum-frequency spectroscopy and imaging of aligned helical polypeptides. *IEEE J. Sel. Top. Quantum Electron.* 10, 1154–1163.
- Koch, W.J., 2004. Genetic and phenotypic targeting of beta-adrenergic signaling in heart failure. *Mol. Cell. Biochem.* 263, 5–9.
- Koffas, T.S., Amitay-Sadovsky, E., Kim, J., Somorjai, G.A., 2004. Molecular composition and mechanical properties of biopolymer interfaces studied by sum frequency generation vibrational spectroscopy and atomic force microscopy. *J. Biomater. Sci. Polym. Ed.* 4, 475–509.
- Krimm, S., Bandekar, J., 1986. Vibrational spectroscopy and conformation of peptides, polypeptides, and proteins. *Adv. Protein Chem.* 38, 181–364.
- Kučerka, N., Nieh, M.P., Pencic, J., Harroun, T., Katsaras, J., 2007. The study of liposomes, lamellae and membranes using neutrons and X-rays. *Curr. Opin. Coll. Int. Sci.* 12, 17–22.
- Laflamme, E., Badia, A., Lafleur, M., Schwartz, J.L., Laprade, R., 2008. Atomic force microscopy imaging of *Bacillus thuringiensis* Cry1 toxins interacting with insect midgut apical membranes. *J. Membrane Biol.* 222, 127–139.
- Lambert, A.G., Davies, P.B., Neivandt, D.J., 2005. Implementing the theory of sum frequency generation vibrational spectroscopy: a tutorial review. *Appl. Spectrosc. Rev.* 40, 103–145.
- Laver, D.R., 1994. The barrel-stave model as applied to alamethicin and its analogs reevaluated. *Biophys. J.* 66, 355–359.
- Lee, A.G., 2005. How lipids and proteins interact in a membrane: a molecular approach. *Mol. Biosyst.* 1, 203–212.
- Lee, D., Walter, K.F.A., Brückner, A.-K., Hilty, C., Becker, S., Griesinger, C., 2008. Bilayer in small bicelles revealed by lipid-protein interactions using NMR spectroscopy. *J. Am. Chem. Soc.* 130, 13822–13823.
- Lee, S., Krimm, S., 1998a. Ab initio-based vibrational analysis of alpha-poly(L-alanine). *Biopolymers* 46, 283–317.
- Lee, S., Krimm, S., 1998b. Polarized Raman spectra of oriented films of alpha-helical poly(L-alanine) and its N-deuterated analogue. *J. Raman Spectrosc.* 29, 73–80.
- Lee, S.H., Wang, J., Krimm, S., Chen, Z., 2006. Irreducible representation and projection operator application to understanding nonlinear optical phenomena: Hyper-Raman, sum frequency generation, and four-wave mixing spectroscopy. *J. Phys. Chem. A* 110, 7035–7044.
- Leitgeb, B., Szekeres, A., Manczinger, L., Vagvolgyi, C., Kredics, L., 2007. The history of alamethicin: A review of the most extensively studied peptaibol. *Chem. Biodivers.* 4, 1027–1051.
- Levy, D., Briggman, K.A., 2007. Cholesterol/phospholipid interactions in hybrid bilayer membranes. *Langmuir* 23, 7155–7161.
- Li, Q., Hua, R., Chou, K.C., 2008. Electronic and conformational properties of the conjugated polymer MEH-PPV at a buried film/solid interface investigated by two-dimensional IR-visible sum frequency generation. *J. Phys. Chem. B* 112, 2315–2318.
- Li, Y.-J., Rothwarf, D.M., Scheraga, H.A., 1998. An unusual adduct of dithiothreitol with a pair of cysteine residues of a protein as a stable folding intermediate. *J. Am. Chem. Soc.* 120, 2668–2669.
- Lindblom, G., Grobner, G., 2006. NMR on lipid membranes and their proteins. *Curr. Opin. Colloid Interface Sci.* 11, 24–29.
- Liu, J., Conboy, J.C., 2004a. Direct measurement of the transbilayer movement of phospholipids by sum-frequency vibrational spectroscopy. *J. Am. Chem. Soc.* 126, 8376–8377.
- Liu, J., Conboy, J.C., 2004b. Phase transition of a single lipid bilayer measured by sum-frequency vibrational spectroscopy. *J. Am. Chem. Soc.* 126, 8894–8895.
- Liu, J., Conboy, J.C., 2005a. Structure of a gel phase lipid bilayer prepared by the Langmuir–Blodgett/Langmuir–Schaefer method characterized by sum-frequency vibrational spectroscopy. *Langmuir* 21, 9091–9097.
- Liu, J., Conboy, J.C., 2005b. 1,2-Diacyl-phosphatidylcholine flip-flop measured directly by sum-frequency vibrational spectroscopy. *Biophys. J.* 89, 2522–2532.
- Liu, J., Conboy, J.C., 2007. Asymmetric distribution of lipids in a phase segregated phospholipid bilayer observed by sum-frequency vibrational spectroscopy. *J. Phys. Chem. C* 111, 8988–8999.
- Lobau, J., Sassi, M., Pohle, W., Selle, C., Koch, M.H.J., Wolfrum, K., 1999. Chain fluidity and phase behaviour of phospholipids as revealed by FTIR and sum-frequency spectroscopy. *J. Mol. Struct.* 481, 407–411.
- Lodowski, D.T., Pitcher, J.A., Capel, W.D., Lefkowitz, R.J., Tesmer, J.J.G., 2003. Keeping G proteins at bay: A complex between G protein-coupled receptor kinase 2 and G beta gamma. *Science* 300, 1256–1262.
- Ma, G., Allen, H.C., 2006. DPPC Langmuir monolayer at the air–water interface: probing the tail and head groups by vibrational sum frequency generation spectroscopy. *Langmuir* 22, 5341–5349.
- Ma, G., Allen, H.C., 2007. Condensing effect of palmitic acid on DPPC in mixed Langmuir monolayers. *Langmuir* 23, 589–597.
- Marsh, D., Muller, M., Schmitt, F.J., 2000. Orientation of the infrared transition moments for an alpha-helix. *Biophys. J.* 78, 2499–2510.
- Marsh, D., Jost, M., Peggion, C., Toniolo, C., 2007a. TOAC spin labels in the backbone of alamethicin: EPR studies in lipid membranes. *Biophys. J.* 92, 473–481.
- Marsh, D., Jost, M., Peggion, C., Toniolo, C., 2007b. Lipid chainlength dependence for incorporation of alamethicin in membranes: EPR studies on TOAC-spin labelled analogues. *Biophys. J.* 92, 4002–4011.
- Marsh, D., 2008. Energetics of hydrophobic matching in lipid-protein interactions. *Biophys. J.* 94, 3996–4013.
- Mateo, C.R., Gómez, J., Villalán, J., González Ros, J.M., 2006. *Protein-lipid Interaction: New Approaches and Emerging Concepts*. Springer, Berlin.
- Mathew, M.K., Balaran, P., 1983a. Alamethicin and related membrane channel forming polypeptides. *Mol. Cell. Biochem.* 50, 47–64.
- Mathew, M.K., Balaran, P., 1983b. A helix dipole model for alamethicin and related transmembrane channels. *FEBS Lett.* 157, 1–5.
- Matsuzaki, K., Nakayama, M., Fukui, M., Otaka, A., Funakoshi, S., Fujii, N., Bessho, K., Miyajima, K., 1993. Role of disulfide linkages in tachyplesin-lipid interactions. *Biochemistry* 32, 11704–11710.
- McConnell, H.M., Watts, T.H., Weis, R.M., Brian, A.A., 1986. Supported planar membranes in studies of cell-cell recognition in the immune system. *Biochim. Biophys. Acta* 864, 95–106.
- McIntosh, T.J., Simon, S.A., 2006. Roles of bilayer material properties in function and distribution of membrane proteins. *Annu. Rev. Biophys. Biomol. Struct.* 35, 177–198.
- Mermut, O., Phillips, D.C., York, R.L., McCrea, K.R., Ward, R.S., Somorjai, G.A., 2006. In situ adsorption studies of a 14-amino acid leucine-lysine peptide onto hydrophobic polystyrene and hydrophilic silica surfaces using quartz crystal microbalance, atomic force microscopy, and sum frequency generation vibrational spectroscopy. *J. Am. Chem. Soc.* 128, 3598–3607.
- Miranda, P.B., Shen, Y.R., 1999. Liquid interfaces: a study by sum-frequency vibrational spectroscopy. *J. Phys. Chem. B* 103, 3292–3307.
- Mizuguchi, M., Kamata, S., Kawabat, S., Kawan, K., 2003. Structure of horseshoe crab antimicrobial peptide tachyplesin I in dodecylphosphocholine micelles. *Peptide Sci.* 39, 281–282.
- Moore, F.G., Richmond, G.L., 2008. Integration or segregation: how do molecules behave at oil/water interfaces? *Acc. Chem. Res.* 41, 739–748.
- Mottamal, M., Lazaridis, T., 2006. Voltage-dependent energetics of alamethicin monomers in the membrane. *Biophys. Chem.* 122, 50–57.
- Nagaraj, R., Balaran, P., 1981. Alamethicin. A transmembrane channel. *Acc. Chem. Res.* 14, 356–362.
- Naito, A., Kawamura, I., 2007. Solid-state NMR as a method to reveal structure and membrane-interaction of amyloidogenic proteins and peptides. *Biochim. Biophys. Acta* 1768, 1900–1912.
- Nakamura, T., Furunaka, H., Miyata, T., Tokunaga, F., Muta, T., Iwanaga, S., Niwa, M., Takao, T., Shimomishi, Y., 1988. Tachyplesin, a class of antimicrobial peptide

- from the hemocytes of the horseshoe crab (*tachypleus tridentatus*). Isolation and chemical structure. *J. Biol. Chem.* 263, 16709–16713.
- Neves, S.R., Ram, P.T., Iyengar, R., 2002. G protein pathways. *Science* 296, 1636–1639.
- Nickolov, Z.S., Britt, D.W., Miller, J.D., 2006. Sum-frequency spectroscopy analysis of two-component Langmuir monolayers and the associated interfacial water structure. *J. Phys. Chem.* 110, 15506–15513.
- North, C.L., Barranger-Mathys, M., Cafiso, D.S., 1995. Membrane orientation of the N-terminal segment of alamethicin determined by solid-state N-15 NMR. *Biophys. J.* 69, 2392–2397.
- Ohe, C., Ida, Y., Matsumoto, S., Sasaki, T., Goto, Y., Noi, A., Tsurumaru, T., Itoh, K., 2004. Investigations of polymyxin B-phospholipid interactions by vibrational sum frequency generation spectroscopy. *J. Phys. Chem. B* 108, 18081–18087.
- Ohe, C., Sasaki, T., Noi, M., Goto, Y., Itoh, K., 2007a. Sum frequency generation spectroscopic study of the condensation effect of cholesterol on a lipid monolayer. *Anal. Bioanal. Chem.* 388, 73–79.
- Ohe, C., Goto, Y., Noi, M., Arai, M., Kamijo, H., Itoh, K., 2007b. Sum frequency generation spectroscopic studies on phase transitions of phospholipid monolayers containing poly(ethylene oxide) lipids at the air–water interface. *J. Phys. Chem. B* 111, 1693–1700.
- Oh-e, M., Hong, S.C., Shen, Y.R., 2002. Orientations of phenyl sidegroups and liquid crystal molecules on a rubbed polystyrene surface. *Appl. Phys. Lett.* 80, 784–786.
- Oldham, W.M., Hamm, H.E., 2006. Structural basis of function in heterotrimeric G proteins. *Quart. Rev. Biophys.* 39, 117–166.
- Opdahl, A., Koffas, T.S., Amitay-Sadovsky, E., Kim, J., Somorjai, G.A., 2004. Characterization of polymer surface structure and surface mechanical behaviour by sum frequency generation surface vibrational spectroscopy and atomic force microscopy. *J. Phys. Condens. Matter.* 16, R659–R677.
- Perry, A., Neupert, C., Space, B., Moore, P.B., 2006. Theoretical modeling of interface specific vibrational spectroscopy: Methods and applications to aqueous interfaces. *Chem. Rev.* 106, 1234–1258.
- Petralli-Mallow, T.P., Briggman, K.A., Richter, L.J., Stephenson, J.C., Plant, A.L., 1999. Nonlinear optics as a detection scheme for biomimetic sensors: SFG spectroscopy of hybrid bilayer membrane formation. *Proc. SPIE* 3858, 25–31.
- Pitcher, J.A., Ingles, J., Higgins, J.B., Arriza, J.L., Casey, P.J., Kim, C., Benovic, J.L., Kwatra, M.M., Caron, M.G., Lefkowitz, R.J., 1992. Role of beta-gamma-subunits of G-proteins in targeting the beta-adrenergic-receptor kinase to membrane-bound receptors. *Science* 257, 1264–1267.
- Richmond, G.L., 2001. Structure and bonding of molecules at aqueous surfaces. *Annu. Rev. Phys. Chem.* 52, 357–389.
- Richmond, G.L., 2002. Molecular bonding and interactions at aqueous surfaces as probed by vibrational sum frequency spectroscopy. *Chem. Rev.* 102, 2693–2724.
- Richter, R.P., Bérat, R., Brisson, A.R., 2006. Formation of solid-supported lipid bilayers: an integrated view. *Langmuir* 22, 3497–3505.
- Rintoul, L., Carter, E.A., Stewart, S.D., Fredericks, P.M., 2000. Keratin orientation in wool and feathers by polarized Raman spectroscopy. *Biopolymers* 57, 19–28.
- Rocha-Mendoza, I., Yankelevich, D.R., Wang, M., Reiser, K.M., Frank, C.W., Knoesen, A., 2007. Sum frequency vibrational spectroscopy: The molecular origins of the optical second-order nonlinearity of collagen. *Biophys. J.* 93, 4433–4444.
- Roke, S., Schins, J., Muller, M., Bonn, M., 2003. Vibrational spectroscopic investigation of the phase diagram of a biomimetic lipid monolayer. *Phys. Rev. Lett.* 90, 128101.
- Ruppel, G., Weilach, C., 2008. Spectroscopic studies of surface-gas interactions and catalyst restructuring at ambient pressure: mind the gap! *J. Phys. Condens. Matter.* 20, 184019.
- Sackmann, E., 1996. Supported membranes: scientific and practical applications. *Science* 271, 43–48.
- Salamon, Z., Macleod, H.A., Tollin, G., 1997. Surface plasmon resonance spectroscopy as a tool for investigating the biochemical and biophysical properties of membrane protein systems. II: applications to biological systems. *Biochim. Biophys. Acta* 1331, 131–152.
- Sansom, M.S.P., 1993a. Alamethicin and related peptaibols—model ion channels. *Eur. Biophys. J.* 22, 105–124.
- Sansom, M.S.P., 1993b. Structure and function of channel-forming peptaibols. *Quart. Rev. Biophys.* 26, 365–421.
- Sartenaer, Y., Tourillon, G., Dreesen, L., Lis, D., Mani, A.A., Thiry, P.A., Peremans, A., 2007. Sum-frequency generation spectroscopy of DNA monolayers. *Biosens. Bioelectron.* 22, 2179–2183.
- Shen, Y.R., 1984. *The Principles of Nonlinear Optics*. John Wiley & Sons, New York.
- Shen, Y.R., 1989. Surface-properties probed by 2nd-harmonic and sum-frequency generation. *Nature* 337, 519–525.
- Shen, Y.R., Ostroverkhov, V., 2006. Sum-frequency vibrational spectroscopy on water interfaces: polar orientation of water molecules at interfaces. *Chem. Rev.* 106, 1140–1154.
- Shultz, M.J., Schnitzer, C., Simonelli, D., Baldelli, S., 2000. Sum frequency generation spectroscopy of the aqueous interface: ionic and soluble molecular solutions. *Inter. Rev. Phys. Chem.* 19, 123–153.
- Shultz, M.J., Baldelli, S., Schnitzer, C., Simonelli, D., 2002. Aqueous solution/air interfaces probed with sum frequency generation spectroscopy. *J. Phys. Chem. B* 106, 5313–5324.
- Sovago, M., Wurpel, G.W.H., Smits, M., Muller, M., Bonn, M., 2007. Calcium-induced phospholipid ordering depends on surface pressure. *J. Am. Chem. Soc.* 129, 11079–11084.
- Spaar, A., Munster, C., Salditt, T., 2004. Conformation of peptides in lipid membranes studied by X-ray grazing incidence scattering. *Biophys. J.* 87, 396–407.
- Steinem, C., Galla, H., Janshoff, A., 2000. Interaction of melittin with solid supported membranes. *Phys. Chem. Chem. Phys.* 2, 4580–4585.
- Stella, L., Burattini, M., Mazzuca, C., Palleschi, A., Venanzi, M., Coin, I., Peggion, C., Toniolo, C., Pispisa, B., 2007. Alamethicin interaction with lipid membranes: A spectroscopic study on synthetic analogues. *Chem. Biodivers.* 4, 1299–1312.
- Stiopin, I.V., Jayathilake, H.D., Bordenyuk, A.N., Benderskii, A.V., 2008. Heterodyne-detected vibrational sum frequency generation spectroscopy. *J. Am. Chem. Soc.* 130, 2271–2275.
- Tadjeddine, A., Peremans, A., 1996. Vibrational spectroscopy of the electrochemical interface by visible infrared sum-frequency generation. *Surf. Sci.* 368, 377–383.
- Tamm, L.K., McConnell, H.M., 1985. Supported phospholipid-bilayers. *Biophys. J.* 47, 105–113.
- Tamm, L.K., 1988. Lateral diffusion and fluorescence microscope studies on a monoclonal-antibody specifically bound to supported phospholipid bilayers. *Biochemistry* 27, 1450–1457.
- Tamm, L.K., Tatulian, U.A., 1997. Infrared spectroscopy of proteins and peptides in lipid bilayers. *Quart. Rev. Biophys.* 30, 365–429.
- Tanaka, M., Sackmann, E., 2005. Polymer-supported membranes as models of the cell surface. *Nature* 437, 656–663.
- Tesmer, V.M., Kawano, T., Shankaranarayanan, A., Kozasa, T., Tesmer, J.J.G., 2005. Snapshot of activated G proteins at the membrane: the Gq-GRK2-G complex. *Science* 310, 1686–1690.
- Thompson, N.L., Palmer III, A.G., 1988. Model cell membranes on planar substrates. *Comments Mol. Cell. Biophys.* 5, 39–56.
- Tyrode, E., Johnson, C.M., Baldelli, S., Leygraf, C., Rutland, M.W., 2005. A vibrational sum frequency spectroscopy study of the liquid–gas interface of acetic acid–water mixtures: 2. Orientation analysis. *J. Phys. Chem. B* 109, 329–341.
- Vass, E., Hollosi, M., Besson, F., Buchet, R., 2003. Vibrational spectroscopic detection of beta- and gamma-turns in synthetic and natural peptides and proteins. *Chem. Rev.* 103, 1917–1954.
- Vogel, H., 1987. Comparison of the conformation and orientation of alamethicin and melittin in lipid-membranes. *Biochemistry* 26, 4562–4572.
- Voges, A.B., Stokes, G.Y., Gibbs-Davis, J.M., Lettan, R.B., Bertin, P.A., Pike, R.C., Nguyen, S.T., Scheidt, K.A., Geiger, F.M., 2007. Insights into heterogeneous atmospheric oxidation chemistry: Development of a tailor-made synthetic model for studying tropospheric surface chemistry. *J. Phys. Chem. C* 111, 1567–1578.
- Wall, M.A., Coleman, D.E., Lee, E., Iniguez-Lluhi, J.A., Posner, B.A., Gilman, A.G., Sprang, S.R., 1995. The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2. *Cell* 83, 1047–1058.
- Wang, G.S., 2008. NMR of membrane-associated peptides and proteins. *Curr. Protein Pept. Sci.* 9, 50–69.
- Wang, H.F., Gan, W., Lu, R., Rao, Y., Wu, B.H., 2005a. Quantitative spectral and orientational analysis in surface sum frequency generation vibrational spectroscopy (SFG-VS). *Int. Rev. Phys. Chem.* 24, 191–256.
- Wang, J., Chen, C.Y., Buck, S.M., Chen, Z., 2001. Molecular chemical structure on poly(methyl methacrylate) (PMMA) surface studied by sum frequency generation (SFG) vibrational spectroscopy. *J. Phys. Chem. B* 105, 12118–12125.
- Wang, J., Buck, S.M., Chen, Z., 2003a. The effect of surface coverage on conformation changes of bovine serum albumin molecules at the air–solution interface detected by sum frequency generation vibrational spectroscopy. *Analyst* 128, 773–778.
- Wang, J., Even, M.A., Chen, X., Schmaier, A.H., Waite, J.H., Chen, Z., 2003b. Detection of amide I signals of interfacial proteins in situ using SFG. *J. Am. Chem. Soc.* 125, 9914–9915.
- Wang, J., Clarke, M.L., Zhang, Y.B., Chen, X., Chen, Z., 2003c. Using isotope-labeled proteins and sum frequency generation vibrational spectroscopy to study protein adsorption. *Langmuir* 19, 7862–7866.
- Wang, J., Chen, X., Clarke, M.L., Chen, Z., 2005b. Detection of chiral sum frequency generation vibrational spectra of proteins and peptides at interfaces in situ. *Proc. Natl. Acad. Sci. USA* 102, 4978–4983.
- Wang, J., Clarke, M.L., Chen, X.Y., Even, M.A., Johnson, W.C., Chen, Z., 2005c. Molecular studies on protein conformations at polymer/liquid interfaces using sum frequency generation vibrational spectroscopy. *Surf. Sci.* 587, 1–11.
- Wang, J., Chen, X., Clarke, M.L., Chen, Z., 2006. Vibrational spectroscopic studies on fibrinogen adsorption at polystyrene/protein solution interfaces: hydrophobic side chain and secondary structure changes. *J. Phys. Chem. B.* 110, 5017–5024.
- Wang, J., Paszti, Z., Clarke, M.L., Chen, X.Y., Chen, Z., 2007. Deduction of structural information of interfacial proteins by combined vibrational spectroscopic methods. *J. Phys. Chem. B* 111, 6088–6095.
- Wang, J., Lee, S.H., Chen, Z., 2008. Quantifying the ordering of adsorbed proteins in situ. *J. Phys. Chem. B* 112, 2281–2290.
- Watry, M.R., Tarbuck, T.L., Richmond, G.L., 2003. Vibrational sum-frequency studies of a series of phospholipid monolayers and the associated water structure at the vapor/water interface. *J. Phys. Chem.* 107, 512–518.
- White, R.J., Zhang, B., Daniel, S., Tang, J.M., Ervin, E.N., Cremer, P.S., White, H.S., 2006. Ionic conductivity of the aqueous layer separating a lipid bilayer membrane and a glass support. *Langmuir* 22, 10777–10783.
- Williams, C.T., Beattie, D.A., 2002. Probing buried interfaces with non-linear optical spectroscopy. *Surf. Sci.* 500, 545–576.
- Woolley, G.A., Wallace, B.A., 1992. Model ion channels—gramicidin and alamethicin. *J. Membr. Biol.* 129, 109–136.

- Yang, C.S.C., Richter, L.J., Stephenson, J.C., Briggman, K.A., 2002. In situ, vibrationally resonant sum frequency spectroscopy study of the self-assembly of dioctadecyl disulfide on gold. *Langmuir* 18, 7549–7556.
- Yang, L., Harroun, T.A., Weiss, T.M., Ding, L., Huang, H.W., 2001. Barrel-stave model or toroidal model? A case study on melittin pores. *Biophys. J.* 81, 1475–1485.
- Ye, S.J., McClelland, A., Majumdar, P., Staflieni, S.J., Daniels, J., Chisholm, B., Chen, Z., 2008. Detection of tethered biocide moiety segregation to silicone surface using sum frequency generation vibrational spectroscopy. *Langmuir* 24, 9686–9694.
- Yeagle, P.L., 2005. *The Structure of Biological Membranes*. CRC Press, Boca Raton, FL.
- York, R.L., Holinga, G.J., Guyer, D.R., McCrea, K.R., Ward, R.S., Somorjai, G.A., 2008. A new optical parametric amplifier based on lithium thioindate used for sum frequency generation vibrational spectroscopic studies of the amide I mode of an interfacial model peptide. *Appl. Spectrosc.* 62, 937–1047.
- Zhao, J., Tamm, L.K., 2003. FTIR and fluorescence studies of interactions of synaptic fusion proteins in polymer-supported bilayers. *Langmuir* 19, 1838–1846.
- Zhu, X.D., Suhr, H., Shen, Y.R., 1987. Surface vibrational spectroscopy by infrared-visible sum frequency generation. *Phys. Rev. B* 35, 3047–3050.
- Zhuang, X., Miranda, P.B., Kim, D., Shen, Y.R., 1999. Mapping molecular orientation and conformation at interfaces by surface nonlinear optics. *Phys. Rev. B* 59, 12632–12640.
- Zhuang, X., Shen, Y.R., 1996. The application of nonlinear optics to the study of polymers at interfaces. *Trends Polym. Sci.* 4, 258–264.