Effects of parity violation on non-Gaussianity of primordial gravitational waves in Hořava-Lifshitz gravity

Tao Zhu,^{1,2,*} Wen Zhao,^{3,†} Yongqing Huang,^{1,‡} Anzhong Wang,^{1,§} and Qiang Wu^{2,||}

¹GCAP-CASPER, Physics Department, Baylor University, Waco, Texas 76798-7316, USA

²Institute for Advanced Physics and Mathematics, Zhejiang University of Technology, Hangzhou 310032, China

³Department of Astronomy, University of Science and Technology of China, Hefei 230026, China

(Received 11 May 2013; published 4 September 2013)

In this paper, we study the effects of parity violation on non-Gaussianities of primordial gravitational waves in the framework of the Hořava-Lifshitz theory of gravity, in which high-order spatial derivative operators, including the ones violating parity, generically appear. By calculating the three-point function, we find that the leading-order contributions to the non-Gaussianities come from the usual second-order derivative terms, which produce the same bispectrum as that found in general relativity. The contributions from high-order spatial *n*th derivative terms are always suppressed by a factor $(H/M_*)^{n-2}$ ($n \ge 3$), where *H* denotes the inflationary energy and M_* the suppression mass scale of the high-order spatial derivative operators of the theory. Therefore, the next leading-order contributions come from the three-dimensional gravitational Chern-Simons term. With some reasonable arguments, it is shown that this three-dimensional operator is the only one that violates the parity and in the meantime has nonvanishing contributions to non-Gaussianities.

DOI: 10.1103/PhysRevD.88.063508

PACS numbers: 98.80.Cq, 98.80.-k, 04.50.-h

I. INTRODUCTION

Primordial gravitational waves (PGWs), which are expected to be generated during inflation, have attracted a great deal of attention recently, as their detections would be the direct evidence of inflation, and more important the existence of gravitational waves in the Universe. From the properties of PGWs, such as their power spectra and non-Gaussianities, we can extract useful information about the theory of inflation and gravity. In particular, the PGWs produce not only the temperature anisotropy, but also a distinguishable signature in the polarization of the cosmic microwave background (CMB) [1]. Decomposing the polarization into two modes: one is curl-free, the E-mode, and the other is divergence-free, the B-mode, one finds that the B-mode pattern cannot be produced by density fluctuations. Thus, its detection would provide a unique signature for the existence of PGWs [2].

In addition, PGWs normally produce the TT, EE, BB and TE spectra of CMB, but the spectra of TB and EB vanish when the parity of the PGWs is conserved [1]. However, if the theory is chiral, the power spectra of right-hand and left-hand PGWs can have different amplitudes, and then induce nonvanishing TB and EB correlation in large scales [3]. This provides the opportunity to directly detect the chiral asymmetry of the theory by observations [3–5]. Recently, in [6,7] the above-mentioned problem was addressed in the framework of Hořava-Lifshitz (HL) theory of gravity [8], in which the Lorentz

symmetry is broken in the ultraviolet (UV), and parityviolating operators generically appear. In particular, it was shown that, because of the parity violation and nonadiabatic evolution of the modes, a large polarization of PGWs can be produced, and could be well within the range of detection of the forthcoming CMB observations [7].

The effects of the parity violation on non-Gaussianities of PGWs were also studied [9,10] in the theory with the general covariance, and shown that, because of the symmetry of the pure de Sitter background, the parity violation from Weyl cubic terms has no contributions to the non-Gaussianities, although this is no longer true when the coupling of Weyl cubic terms is time dependent [11]. It should be noted that in all these studies the symmetry of the general diffeomorphisms of the underlaid theories plays a crucial role. On the other hand, in the HL theory the symmetry is reduced to the foliation-preserving diffeomorphisms [8], and the parity-violating operators allowed by such a symmetry are quite different from those with the general diffeomorphisms. Thus, it is expected that in the HL theory some distinguishable features of non-Gaussianities of PGWs due to these parity-violating operators should exist, which may provide a smoking gun for the tests of the HL theory in the forthcoming CMB observations. With these motivations, in this paper we study the non-Gaussianities of PGWs in the HL theory, and focus ourselves mainly on the effects of the high-order spatial operators on the non-Gaussianities of PGWs, especially on the ones that violate the parity.

The rest of the paper is organized as follows: In Sec. II we first give a very brief review on the HL theory, and then restrict ourselves to the model recently proposed in [12,13], where an extra U(1) symmetry is enforced in the nonprojectable case, in order to eliminate the spin-0 gravitons

^{*}Tao_Zhu@baylor.edu

wzhao7@mail.ustc.edu.cn

[‡]Yongqing_Huang@baylor.edu

Anzhong_Wang@baylor.edu

wuq@zjut.edu.cn

usually appearing in the HL theory. In this section, we also present the linearized equation of motion of the tensor perturbations, originally derived in [7]. In Sec. III, from the cubic action of tensor perturbations, we calculate the three-point correlation function and obtain the bispectrum of the PGWs, while in Sec. IV, we plot the shapes of bispectrum produced by both the second-order derivative operators and the three-dimensional parity-violating Chern-Simons one. In Sec. V, we summarize our main results. There are also two appendices, A and B, in which the cubic action is given explicitly.

Before processing further, we would like to note that, although in this paper we restrict ourselves only to the model of the HL theory proposed recently in [12,13], our results can be easily generalized to other models, as the tensor perturbations are quite similar in all of these models [7,14]. In addition, non-Gaussianities of PGWs in the framework of the general covariant theory with the projectability condition was also studied in [15], and several remarkable features were found. In particular, it was found that the terms $R_{ij}R^{ij}$ and $(\nabla^i R^{jk})$ $(\nabla_i R_{ik})$ exhibit a peak at the squeezed limit, while the one $R_i^i R_k^j R_i^k$ favors the equilateral shape when spins of the three tensor fields are the same, but peaks in between the equilateral and squeezed limits when spins are mixed, where R_{ii} denotes the three-dimensional Ricci tensor made of the threedimensional metric g_{ij} of the leaves t = Constant, and ∇_i denotes the covariant derivative with respect to g_{ii} . The consistency with the recently released Planck observations [16] was also discussed. However, in [15] the parity-violating operators were excluded. Therefore, in this paper we shall focus mainly on the effects of these operators on non-Gaussianities, as mentioned above. Moreover, non-Gaussianities of scalar perturbations were also studied in the framework of the HL theory, one in the curvaton scenario [17] and the other in the inflationary model [18], and some remarkable features were obtained.

II. NONPROJECTABLE GENERAL COVARIANT HL GRAVITY AND LINEAR TENSOR PERTURBATIONS

By construction, the HL theory is power-counting renormalizable [8]. This is achieved by breaking the symmetry of the general covariance in the UV, and including only high-order spatial derivative operators, so that it remains also unitary, a problem that has been faced for a long time in the quantization of gravity [19]. In the low energy, low dimensional operators take over, and it is expected that the Lorentz symmetry is "accidentally" restored [20]. Since Hořava first proposed it in 2009, the theory has attracted a lot of attention, partially because of various remarkable features when applied to cosmology [21], and partially because of some challenging questions, such as ghosts, instability, and strong coupling. To overcome these questions, various models have been proposed [20], including the ones with an additional local U(1) symmetry [12,13,22], in which the problems mentioned above can be avoided by properly choosing the coupling constants appearing in the theory. Since in all of those models, the tensor perturbations are almost the same [7,14,15], without loss of the generality, in this paper we shall work with the model proposed in [12,13].

A. Action of the nonprojectable general covariant HL gravity

The fundamental variables in the nonprojectable general covariant HL gravity proposed in [12,13] are

$$(N, N^i, g_{ij}, A, \varphi),$$

where N and Nⁱ denote, respectively, the lapse function and shift vector in the Arnowitt-Deser-Misner decompositions [23], and A and φ are, respectively, the U(1) gauge field and Newtonian prepotential [22]. Then, the corresponding total action can be cast in the form

$$S = \zeta^2 \int dt d^3x \sqrt{g} N(\mathcal{L}_K - \mathcal{L}_V + \mathcal{L}_A + \mathcal{L}_{\varphi} + \zeta^{-2} \mathcal{L}_M),$$
(2.1)

where $\zeta^2 = 1/(16\pi G)$ with G being the Newtonian constant, \mathcal{L}_M describes matter fields, and

$$\mathcal{L}_{K} = K_{ij}K^{ij} - \lambda K^{2}, \qquad \mathcal{L}_{V} = \mathcal{L}_{V}^{R} + \mathcal{L}_{V}^{a},$$

$$\mathcal{L}_{A} = \frac{A}{N}(2\Lambda_{g} - R),$$

$$\mathcal{L}_{\varphi} = \varphi G^{ij}(2K_{ij} + \nabla_{i}\nabla_{j}\varphi + a_{i}\nabla_{j}\varphi)$$

$$+ (1 - \lambda)[(\Delta\varphi + a_{i}\nabla^{i}\varphi)^{2} + 2(\Delta\varphi + a_{i}\nabla^{i}\varphi)K]$$

$$+ \frac{1}{3}\hat{G}^{ijlk}[4(\nabla_{i}\nabla_{j}\varphi)a_{(k}\nabla_{l)}\varphi + 5(a_{(i}\nabla_{j)}\varphi)a_{(k}\nabla_{l)}\varphi$$

$$+ 2(\nabla_{(i}\varphi)a_{j)(k}\nabla_{l)}\varphi + 6K_{ij}a_{(l}\nabla_{k)}\varphi], \qquad (2.2)$$

with $\Delta \equiv \nabla^2$, and

$$\begin{split} K_{ij} &= \frac{1}{2N} (-\dot{g}_{ij} + \nabla_i N_j + \nabla_j N_i), \qquad a_i = \frac{N_{,i}}{N}, \\ a_{ij} &= \nabla_j a_i, \qquad \hat{G}^{ijlk} = g^{il} g^{jk} - g^{ij} g^{kl}, \\ G_{ij} &= R_{ij} - \frac{1}{2} g_{ij} R + \Lambda_g g_{ij}, \\ \mathcal{L}_V^R &= \gamma_0 \zeta^2 + \gamma_1 R + \frac{\gamma_2 R^2 + \gamma_3 R_{ij} R^{ij}}{\zeta^2} + \frac{\gamma_5}{\zeta^4} C_{ij} C^{ij}, \\ \mathcal{L}_V^a &= -\beta_0 a_i a^i + \frac{1}{\zeta^2} [\beta_1 (a_i a^i)^2 + \beta_2 (a^i_i)^2 \\ &+ \beta_3 (a_i a^i) a^j_j + \beta_4 a^{ij} a_{ij} + \beta_5 (a_i a^i) R \\ &+ \beta_6 a_i a_j R^{ij} + \beta_7 R a^i_i] + \frac{1}{\zeta^4} \beta_8 (\Delta a^i)^2. \end{split}$$
(2.3)

Here *R* denotes the Ricci scalar, and C_{ij} the Cotton tensor, defined by

$$C^{ij} = \frac{e^{ikl}}{\sqrt{g}} \nabla_k \left(R^j_l - \frac{1}{4} R \delta^j_l \right), \qquad (2.4)$$

with $e^{123} = 1$, etc. λ , γ_n , β_s , and Λ_g are the coupling constants of the theory. In terms of R_{ij} , we have [13]

$$C_{ij}C^{ij} = \frac{1}{2}R^3 - \frac{5}{2}RR_{ij}R^{ij} + 3R_j^i R_k^j R_k^k + \frac{3}{8}R\Delta R + (\nabla_i R_{jk})(\nabla^i R^{jk}) + \nabla_k G^k, \quad (2.5)$$

where

$$G^{k} = \frac{1}{2} R^{jk} \nabla_{j} R - R_{ij} \nabla^{j} R^{ik} - \frac{3}{8} R \nabla^{k} R.$$
 (2.6)

It should be noted that in writing the above action, we have excluded all the terms that violate the parity [12,13]. For our current purpose, we add the fifth- and third-order spatial derivative operators to the potential \mathcal{L}_V [7],

$$\Delta \mathcal{L}_{V} = \frac{1}{M_{*}^{3}} (\alpha_{0} K_{ij} R_{ij} + \alpha_{2} \epsilon^{ijk} R_{il} \Delta_{j} R_{k}^{l}) + \frac{\alpha_{1} \omega_{3}(\Gamma)}{M_{*}} + \cdots$$
(2.7)

Here the coupling constant α_0 , α_1 , α_2 are dimensionless and arbitrary, $\epsilon^{ijk} = e^{ijk}/\sqrt{g}$ is the total antisymmetric tensor, and $\omega_3(\Gamma)$ the three-dimensional gravitational Chern-Simons term.¹ "···" denotes the rest of the fifthorder operators given in Eq. (2.6) of [13]. Since they have no contributions to tensor perturbations, in this paper we shall not write them out explicitly. As shown in [7], because of the additional parity violation terms of Eq. (2.7), the nonadiabatic evolution of modes lead to a large polarization of PGWs, and it could be well within the detection of CMB observations, as mentioned above. In this paper, we investigate their effects on the non-Gaussianities of PGWs.

B. The linearized tensor perturbations

The general formulas of the linearized tensor perturbations were given in [7], so in the rest of this section we give a very brief summary of the main results obtained there, in order to initiate our studies of the non-Gaussianities of PGWs in the next section. For details, we refer readers to [7]. Consider a flat Friedmann-Robertson-Walker (FRW) universe,

$$\hat{N} = a(\eta), \qquad \hat{N}^{i} = \hat{A} = \hat{\varphi} = 0,$$

$$\hat{g}_{ij} dx^{i} dx^{j} = a(\eta)^{2} \delta_{ij} dx^{i} dx^{j},$$
(2.8)

where quantities with hats denote the background of the FRW universe in the coordinates $(\eta, x^i) = (\eta, x, y, z)$. Then, the tensor perturbations are given by

$$\delta N = \delta N^i = \delta A = \delta \varphi = 0, \quad \delta g_{ij} = a^2 h_{ij}(\eta, \mathbf{x}). \quad (2.9)$$

Assuming that matter fields have no contributions to tensor perturbations, we find that the quadratic part of the total action can be cast in the form

$$S_{g}^{(2)} = \zeta^{2} \int d\eta d^{3}x \bigg\{ \frac{a^{2}}{4} (h_{ij}^{\prime})^{2} - \frac{1}{4} a^{2} (\partial_{k} h_{ij})^{2} \\ - \frac{\hat{\gamma}_{3}}{4M_{*}^{2}} (\partial^{2} h_{ij})^{2} - \frac{\hat{\gamma}_{5}}{4M_{*}^{4}a^{2}} (\partial^{2} \partial_{k} h_{ij})^{2} \\ - \frac{\alpha_{1} a e^{ijk}}{2M_{*}} (\partial_{l} h_{i}^{m} \partial_{m} \partial_{j} h_{k}^{l} - \partial_{l} h_{im} \partial^{l} \partial_{j} h_{k}^{m}) \\ - \frac{\alpha_{2} e^{ijk}}{4M_{*}^{3}a} \partial^{2} h_{il} (\partial^{2} h_{k}^{l})_{,j} - \frac{3\alpha_{0} \mathcal{H}}{8M_{*}a} (\partial_{k} h_{ij})^{2} \bigg\}, \quad (2.10)$$

where $h'_{ij} \equiv \partial h_{ij} / \partial \eta$, $\partial^2 \equiv \delta^{ij} \partial_i \partial_j$, $\mathcal{H} = a'/a$, and

$$\gamma_3 \equiv \left(\frac{M_{pl}}{2M_*}\right)^2 \hat{\gamma}_3, \qquad \gamma_5 \equiv \left(\frac{M_{pl}}{2M_*}\right)^4 \hat{\gamma}_5.$$

To avoid fine-tuning, α_n and $\hat{\gamma}_n$ are expected to be of the same order. Then, the field equations for h_{ij} read

$$h_{ij}'' + 2\mathcal{H}h_{ij}' - \alpha^2 \partial^2 h_{ij} + \frac{\hat{\gamma}_3}{a^2 M_*^2} \partial^4 h_{ij} - \frac{\hat{\gamma}_5}{a^4 M_*^4} \partial^6 h_{ij} + e_i^{\ lk} \left(\frac{2\alpha_1}{M_* a} + \frac{\alpha_2}{M_*^3 a^3} \partial^2\right) (\partial^2 h_{jk})_{,l} = 0, \qquad (2.11)$$

where $\alpha^2 \equiv 1 + 3\alpha_0 \mathcal{H}/(2M_*^3 a)$.

To study the evolution of h_{ij} , we expand it over spatial Fourier harmonics,

$$h_{ij}(\boldsymbol{\eta}, \mathbf{x}) = \sum_{s=R,L} \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \psi_k^s(\boldsymbol{\eta}) e^{i\mathbf{k}\cdot\mathbf{x}} P_{ij}^{(s)}(\hat{\mathbf{k}}), \quad (2.12)$$

where $P_{ij}^{(s)}(\hat{\mathbf{k}})$ are the circular polarization tensors and satisfy the relations: $ik_m e^{rmj} P_{ij}^{(s)} = k\rho^s P_i^{r(s)}$ with $\rho^R = 1$, $\rho^L = -1$, and $P^{*i(s)}_{ij} P_i^{j(s')} = \delta^{ss'}$ [6]. Define $u_k^s(\eta) = \frac{1}{2}a(\eta)M_{pl}\psi_k^s(\eta)$ and with the de Sitter background $a = -1/(H\eta)$, we obtain

$$u_{k}^{s}(\eta)'' + \left[\omega_{s}^{2}(k,\eta) - \frac{2}{\eta^{2}}\right]u_{k}^{s}(\eta) = 0, \quad (2.13)$$

where

$$\omega_s^2(k,\eta) \equiv \alpha^2 k^2 [1 - \delta_1 \rho^s(\epsilon_* \alpha k \eta) + \delta_2(\epsilon_* \alpha k \eta)^2 + \delta_3 \rho^s(\epsilon_* \alpha k \eta)^3 + \delta_4(\epsilon_* \alpha k \eta)^4], \qquad (2.14)$$

with $\epsilon_* \equiv H/M_* \ll 1$, and

$$\delta_1 \equiv \frac{2\alpha_1}{\alpha^3}, \qquad \delta_2 \equiv \frac{\hat{\gamma}_3}{\alpha^4}, \qquad \delta_3 \equiv \frac{\alpha_2}{\alpha^5}, \qquad \delta_4 \equiv \frac{\hat{\gamma}_5}{\alpha^6}.$$
(2.15)

¹To take quantum effects into account, it was proposed to add boundary terms $\Delta S_3 = \sum_i \beta_i M^{3-\Delta_i} \int_{t=t_*} d^3x \sqrt{g} \mathcal{O}^i$ into the Einstein-Hilbert action at the moment $t = t_*$, right before the inflation started [24]. Clearly, one choice of \mathcal{O}^i is $\mathcal{O}^i \propto \omega(\Gamma)$. We thank Jiro Soda for pointing it out to us.

Following [25,26], we choose the initial conditions at $\eta = \eta_i$ as

$$u_{k}^{s}(\eta_{i}) = \frac{1}{\sqrt{2\omega_{s}(k,\eta_{i})}}, \quad u_{k}^{s}(\eta_{i})' = i\sqrt{\frac{\omega_{s}(k,\eta_{i})}{2}}.$$
 (2.16)

Then, if one assumes that $\omega_s(k, \eta)$ is slowly varying, i.e.,

$$Q \equiv \left| \frac{\omega_s(k, \eta)'}{\omega_s^2(k, \eta)} \right| \ll 1, \qquad (2.17)$$

one can approximatively treat $\omega_s(k, \eta)$ as constant, and get the approximate solution of the mode function $u_k^s(\eta)$,

$$u_k^s(\eta) \simeq \frac{1}{\sqrt{2\omega_s(k,\eta)}} \left(1 - \frac{i}{\omega_s \eta}\right) e^{-i \int \omega_s d\eta}, \qquad (2.18)$$

or

$$\psi_k^s(\eta) \simeq -2 \frac{iH}{M_{pl}} \frac{1}{\sqrt{2\omega_s^3}} (1 + i\omega_s \eta) e^{-i \int \omega_s d\eta}.$$
 (2.19)

Promoting $\psi_k^s(\eta)$ to a quantum operator,

$$\psi_{k}^{s}(\eta) = \psi_{k}^{s}a_{s}(k) + \psi_{k}^{*s}a_{s}^{\dagger}(k), \qquad (2.20)$$

one finds that the power spectrum of the tensor perturbations is given by

$$\Delta_{\rm T}^2 \equiv \frac{k^3 (|\psi_k^R|^2 + |\psi_k^L|^2)}{2\pi^2} \simeq \frac{2H^2}{\pi^2 M_{pl}^2}, \qquad (2.21)$$

which has the same expression as that given in general relativity (GR). Here $a_s(k)$ and $a_s^{\dagger}(-k)$ are annihilation and creation operators, and their commutation relation is given by

$$[a_{s}(k), a_{s'}^{\dagger}(k')] = (2\pi)^{3}\delta_{ss'}\delta(k - k')$$

It should be noted that, in our calculations we have assumed (2.17). This condition implies that the adiabatic condition is always satisfied (before the modes exit the horizon), and thus, there is no important modification in the power spectrum of PGWs. Once this condition is violated, as shown in [7], some interesting modifications on power spectrum and polarization of PGWs become possible. For simplification, in this paper we assume that (2.17) always holds.

III. THE INTERACTION HAMILTONIAN AND BISPECTRUM

In this section, we turn to the cubic action and the bispectrum of the tensor perturbations. The cubic action $S_g^{(3)}$ is given by Eq. (A1), which can be written in the form

$$S_{\rm g}^{(3)} = -\int d\eta H_{\rm int}(\eta). \tag{3.1}$$

Then the 3-point correlation function can be computed by employing the in-in formalism [27],

$$\langle \psi_{k_1}^{s_1}(\eta) \psi_{k_2}^{s_2}(\eta) \psi_{k_3}^{s_3}(\eta) \rangle = -i \int_{\eta_i}^{\eta} d\eta' \langle [\psi_{k_1}^{s_1}(\eta) \psi_{k_2}^{s_2}(\eta) \psi_{k_3}^{s_3}(\eta), H_{\text{int}}(\eta')] \rangle, \quad (3.2)$$

where η_i represents the early time when inflation starts, and η is a time when the bispectrum is evaluated. A good approximation is to extend the integral into the whole half axis, $\eta \in (-\infty, 0)$. After some simple but very tedious calculations, it can be shown that the 3-point correlation function can be rewritten in the form

$$\begin{split} \psi_{k_{1}}^{s_{1}}(0)\psi_{k_{2}}^{s_{2}}(0)\psi_{k_{3}}^{s_{3}}(0)\rangle \\ &=i(2\pi)^{3}\delta^{3}(k_{1}+k_{2}+k_{3})\zeta^{2} \\ &\times \int_{-\infty}^{0}a^{2}(\eta')d\eta'F_{k_{1}k_{2}k_{3}}^{s_{1}s_{2}s_{3}}(\eta')\Big[W_{k_{1}k_{2}k_{3}}^{s_{1}s_{2}s_{3}}(\eta') \\ &- W_{k_{1}k_{2}k_{3}}^{*s_{1}s_{2}s_{3}}(\eta')\Big], \end{split}$$
(3.3)

where $F_{k_1k_2k_3}^{s_1s_2s_3}(\eta')$ is given in Appendix B, and $W_{k_1k_2k_3}^{s_1s_2s_3}(\eta')$ is defined as

$$\begin{aligned}
W^{s_1s_2s_3}_{k_1k_2k_3}(\eta') \\
&\equiv \psi^{s_1}_{k_1}(0)\psi^{s_2}_{k_2}(0)\psi^{s_3}_{k_3}(0)\psi^{*s_1}_{k_1}(\eta')\psi^{*s_2}_{k_2}(\eta')\psi^{*s_3}_{k_3}(\eta').
\end{aligned}$$
(3.4)

In the de Sitter background, the 3-point correlation function reduces to

$$\langle \psi_{k_1}^{s_1}(0)\psi_{k_2}^{s_2}(0)\psi_{k_3}^{s_3}(0) \rangle$$

= $(2\pi)^7 \delta^3 (k_1 + k_2 + k_3) \frac{\Delta_{\rm T}^4}{2^3 k_1^3 k_2^2 k_3^3} B_{k_1,k_2,k_3}^{s_1 s_2 s_3}, \quad (3.5)$

where

$$B_{k_{1}k_{2}k_{3}}^{s_{1}s_{2}s_{3}} \equiv \sum_{n=0}^{4} \delta_{n} \epsilon_{*}^{n} F_{n} I_{n}, \qquad (3.6)$$

with
$$\delta_0 = 1$$
, and I_n is given by
 $I_n \equiv \operatorname{Im} \left\{ \int_{-\infty}^0 d\eta (-\eta)^{n-2} e^{i \int (\omega_{s_1} + \omega_{s_2} + \omega_{s_3}) d\eta} \\
\times \sqrt{\frac{k_1^3 k_2^3 k_3^3}{\omega_{s_1}^3 \omega_{s_2}^3 \omega_{s_3}^3}} (1 - i\omega_{s_1} \eta) (1 - i\omega_{s_2} \eta) (1 - i\omega_{s_3} \eta) \right\}.$
(3.7)

Usually, the *k* dependence of the bispectrum receives contributions from both the interaction Hamiltonian $H_{int}(\sim \sum F_n)$ and the mode function integration I_n . For the former, one can see from (3.6) that the high order spatial derivative terms do have contributions in bispectrum, but are suppressed by the factor ϵ_* . Then, the leading-order contributions come from the two derivative term F_0 , which has the same expression as that given in general relativity.

On the other hand, the mode function integration I_n in the current case involved very complicated expression, and thus it is very hard to perform the integration explicitly. In general relativity, to minimize the errors, one usually splits the integrals into three different regions: one outside the horizon, one around the horizon, and one inside the horizon. The contribution from the last region vanishes due to

the high-frequency oscillate [28]. A similar conclusion is also applicable to the current case, because when the highorder derivative terms dominate, the oscillation becomes more rapid than that in general relativity. Thus, using the same arguments, one can safely neglect the effects from the high-order derivative terms in the last region.

In the first region, the mode is outside the horizon and the effective mass term $-2/\eta^2$ in Eq. (2.13) dominates. Then, in this region the corresponding results are also the same as those given in general relativity. Therefore, the effects from the parity-violating operators come only from the region around the horizon. In this region, although the k^2 term dominates, the high-derivative terms still have non-negligible contributions. In order to take these effects into account, we expand the integration in terms of ϵ_* . In particular, to the zeroth order of ϵ_* , we have

$$\operatorname{Im}\left\{i\int_{-\infty}^{0}d\eta(-\eta)^{n-2}e^{i(k_{1}+k_{2}+k_{3})\eta}\times(1+ik_{1}\eta)(1+ik_{2}\eta)(1+ik_{3}\eta)\right\},\qquad(3.8)$$

which coincides with the mode integration in general relativity, as expected. Thus, one immediately obtains the bispectrum of the leading order,

$$B_{(GR)}^{s_1s_2s_3}(k_1, k_2, k_3) = \left(-K + \frac{k_1k_2 + k_1k_3 + k_2k_3}{K} + \frac{k_1k_2k_3}{K}\right)F_0, \quad (3.9)$$

where $K \equiv k_1 + k_2 + k_3$, and which is precisely the bispectrum of PGWs given in GR.

Now let us turn to the first order contributions of ϵ_* . Ignoring all the detailed calculations, it can be shown that it takes the form

$$B_{(PV)}^{s_{1}s_{2}s_{3}}(k_{1}, k_{2}, k_{3}) = -\frac{\pi}{2}\delta_{1}\epsilon_{*}\left[F_{1} + \frac{3}{4}(s_{1}k_{1} + s_{2}k_{2} + s_{3}k_{3})F_{0}\right].$$
 (3.10)

. . .

Since this term is directly proportional to δ_1 , from Eqs. (2.7) and (2.15) we find that it represents the contributions of the three-dimensional Chern-Simons term.

For operators with dimensions $n \ge 4$, their contributions can be written in the form

$$\epsilon_{*}^{n-2} \sum_{r} F_{r} \operatorname{Im} \left\{ \int_{-\infty}^{0} d\eta (-\eta)^{n-4} e^{iK\eta} f_{n-r-2}(\eta) \right\}, \quad (3.11)$$

where r = (0, n - 2), and $f_{n-r-2}(K, \eta)$ can be expressed as

$$f_{n-r-2}(K, \eta) \equiv a_0 + a_1(i\eta) + a_2(i\eta)^2 + \dots + a_{n-r+1}(i\eta)^{n-r+1}.$$
(3.12)

Here a_r are functions of k_1 , k_2 , k_3 . In particular, the effects from fifth-order derivative terms should contribute to the bispectrum at the order of ε_*^3 . But, a careful analysis over the integration shows that their contributions vanish identically. This result can be easily generalized to higher order terms. In fact, for n = 2j + 1 with j = 2, 3, 4, ..., the bispectrum of PGWs at the order ϵ_*^{n-2} always vanishes. This implies that $B_{(PV)}^{s_1s_2s_3}(k_1, k_2, k_3)$ given by Eq. (3.10) represents the only contribution from parity violation operators.

In Ref. [10], Soda *et al.* calculated the bispectrum of PGWs from the Weyl cubic terms, W^3 and $\tilde{W}W^2$, and proved that no contributions from parity violation appear in the non-Gaussianity of PGWs in pure de Sitter background. This is consistent with our current results. However, it must be noted that in their considerations, the symmetry of the theory is still of the general diffeomorphisms. As a result, only the parity-violating terms W^3 and $\tilde{W}W^2$ are allowed. These terms are both P-odd and T-odd. Thus, when one calculates the bispectrum, the two terms produce an integral similar to (3.11) with n - 2 = 2j - 1 as an odd number that is greater than two [10,11]. Hence, with the arguments given above, they indeed have no contributions to the bispectrum of PGWs.



FIG. 1 (color online). (a) Shape of $(k_1k_2k_3)^{-1}B_{(\text{GR})}^{+++}(k_1, k_2, k_3)$. (b) Shape of $(k_1k_2k_3)^{-1}B_{(\text{GR})}^{++-}(k_1, k_2, k_3)$. All are normalized to unity in the equilateral limit.



FIG. 2 (color online). Shapes of $(k_1k_2k_3)^{-1}B_{(PV)}^{s_1s_2s_3}(k_1, k_2, k_3)$ for various spin products: (a) + + +; (b) + + -; (c) - - +; (d) - - -. All are normalized to unity in the equilateral limit.

IV. SHAPE OF THE BISPECTRUM

We are now ready to plot the shapes of the bispectrum. For $s_1 = s_2 = s_3 = 1$ and $s_1 = s_2 = -s_3 = 1$, we plot the shapes of the bispectrum of the leading order contributions $(k_1k_2k_3)^{-1}B_{(GR)}^{s_1s_2s_3}(k_1, k_2, k_3)$ in Fig. 1. Because there is no parity violation in the leading order, we have $B_{(GR)}^{+++}(k_1, k_2, k_3) = B_{(GR)}^{---}(k_1, k_2, k_3)$ and $B_{(GR)}^{++-}(k_1, k_2, k_3) = B_{(GR)}^{---+}(k_1, k_2, k_3)$. Thus, there are only two possible configurations, which both peak at the squeezed limit $(k_3/k_1 \rightarrow 0)$. As pointed out in [29], the second configuration [Fig. 1(b)] is subdominant, in comparison with the first one in the equilateral limit $(k_1 \simeq k_2 \simeq k_3)$, that is, $B_{(GR)}^{++-} \simeq B_{(GR)}^{+++-}/81$.

Now we turn to the contributions from the parityviolating Chern-Simons term. In this case, because of the violation of the parity, for different spin products we have four independent configurations. We plot the shapes of these four configurations in Fig. 2, from which it can be seen that all the configurations peak in the squeezed limit [Note that for the (++-) and (---)cases they peak in the negative direction.] More specifically, we have $B_{(PV)}^{+++}(k_1, k_2, k_3) = -B_{(PV)}^{---}(k_1, k_2, k_3)$, and $B_{PV}^{++-}(k_1, k_2, k_3) = -B_{(PV)}^{--++}(k_1, k_2, k_3)$.

V. CONCLUSIONS AND REMARKS

In this paper, we have investigated the non-Gaussianities of PGWs generated during the de Sitter expansion of the Universe in the framework of the HL theory, and paid particular attention on the effects of the operators that violate the parity. Because of the restricted foliationpreserving diffeomorphisms of the theory, the parityviolating third- and fifth-dimensional operators exist generically. By calculating the three-point correlation function of the PGWs, we have shown that the leading-order contributions still come from the second-order spatial derivative terms, and are the same as those given in general relativity. The high-order *n*th spatial derivative terms of the theory also contribute to the bispectrum, although their contributions are suppressed by a factor ϵ_*^{n-2} .

More remarkably, we have also found that the three-dimensional gravitational Chern-Simons operator $\omega(\Gamma)$ is the only one that violates the parity and meantime has nonvanishing contributions to the non-Gaussianities of PGWs. In comparison with the contributions of the second-order operators that produce the same non-Gaussianity as given in general relativity, its contributions are suppressed by the factor ϵ_* . In addition, operators with odd order and higher than three have no contributions to the non-Gaussianities of PGWs.

It should be noted that in obtaining the above results, we have assumed that the adiabatic condition (2.17) is always satisfied before the modes exit the horizon. If this condition fails to hold, the nonadiabatic evolution of the modes

becomes possible, and hence the integral history of the mode function will be dramatically altered. Then, large non-Gaussianities are expected, although it is still an open question how one can extract information of PGWs in this case.

ACKNOWLEDGMENTS

We would like to express our gratitude to Shinji Mukohyama and Jiro Soda for valuable discussions and comments. This work was supported in part by DOE, DE-FG02-10ER41692 (A. W.); by the NSFC with Grants No. 11205133 (Q. W. and T. Z.), No. 11047008 (Q. W. and T. Z.), No. 11105120 (T. Z.), No. 11173021 (W. Z. and A. W.), No. 11075141 (W. Z. and A. W.), No. 11375153 (A. W.), No. 11322324 (W. Z.); Ciência Sem Fronteiras, No. 004/2013-DRI/CAPES (A. W.); and by the project of Knowledge Innovation Program of Chinese Academy of Science (W. Z.).

APPENDIX A: THE CUBIC ACTION OF PGWS

The cubic action for the tensor perturbations can be written in the form

$$S_{\rm g}^{(3)} = \zeta^2 \int d\eta d^3 x a^2 \bigg\{ \frac{\alpha^2}{4} L_2 + \frac{\alpha_1}{aM_*} L_3 + \frac{\hat{\gamma}_3}{a^2 M_*^2} L_4 + \frac{\alpha_2}{a^3 M_*^3} L_5 + \frac{\hat{\gamma}_5}{a^4 M_*^4} L_6 \bigg\},\tag{A1}$$

where

$$L_{2} = h^{mk, ij} (2h_{mi}h_{kj} - h_{ij}h_{mk}),$$

$$L_{3} = -\frac{1}{2} e^{ijk} (\partial^{2}h_{lk})_{l}h_{im}h^{lm} - \frac{e^{ijk}}{4} \Big[4h_{i[m,l]j}h^{lp}(h_{p,k}^{m} + h_{pk}^{m} - h_{k,p}^{m}) + h_{i,l}^{n}h_{j,m}^{l} \Big(\frac{2}{3}h_{k,n}^{m} - 2h_{kn}^{m} \Big) + 2h_{l,i}^{n}h_{m,j}^{l}h_{k,n}^{m} \Big],$$

$$L_{4} = -\frac{1}{2} (\partial^{2}h_{ij})_{,n}^{m}h_{m}^{i}h_{n}^{j} - \frac{1}{2} (\partial^{2}h_{ij})h_{mn} \Big(2h^{im,jn} - \frac{h^{mn,ij}}{2} - h^{ij,mn} \Big),$$

$$L_{5} = \frac{e^{ijk}}{2} \Big[-(\partial^{2}h_{il})h_{,j}^{mp} \Big(h_{p,mk}^{l} - \frac{1}{2}h_{mp,k}^{l}\Big) + (\partial^{2}h_{il,j})h^{mp}(h_{mk,p}^{l} - h_{k,mp}^{l}) + 2(\partial^{2}h_{il,j})h_{mk,p}h^{l[p,m]} - \frac{1}{2}(\partial^{2}h_{il,j})h^{lp}\partial^{2}h_{pk} \Big] \\ - \frac{1}{8} e^{ijk} (\partial^{2}h_{il}) (\partial^{2}h_{k}^{m})(h_{m,j}^{l} - h_{j,m}^{l} - h_{jm}^{l}) - \frac{1}{4} e^{ijk} (\partial^{4}h_{kl,j})h_{im}h^{lm},$$

$$L_{6} = \frac{3}{8} (\partial^{2}h_{j}^{l}) (\partial^{2}h_{k}^{l}) (\partial^{2}h_{k}^{l}) + \frac{1}{2} \Big[(\partial^{4}h_{ij})h_{mn} \Big(2h^{im,jn} - \frac{1}{2}h^{mn,ij} - h^{ij,mn} \Big) + (\partial^{4}h_{ij,n}^{m})h_{m}^{i}h^{jn} + 2(\partial^{2}h_{jk})_{,il} (\partial^{2}h^{lk})h^{ij} \Big].$$
(A2)

Then the interaction Hamiltonian is

$$H_{\rm int}(\eta) = \int d^3x \mathcal{H}_{\rm int}(\eta, x), \qquad \mathcal{H}_{\rm int}(\eta, x) = -\zeta^2 a^2 \left[\frac{\alpha^2}{4} L_2 + \frac{\alpha_1}{aM_*} L_3 + \frac{\hat{\gamma}_3}{a^2 M_*^2} L_4 + \frac{\alpha_2}{a^3 M_*^3} L_5 + \frac{\hat{\gamma}_5}{a^4 M_*^4} L_6 \right].$$
(A3)

APPENDIX B: EXPRESSION OF $F_{k_1k_2k_3}^{s_1s_2s_3}(\eta)$

 $F_{k_1k_2k_3}^{s_1s_2s_3}(\eta)$ is given by

$$F_{k_1k_2k_3}^{s_1s_2s_3}(\eta) = \alpha^2 F_0 + \frac{\alpha_1}{aM_*} F_1 + \frac{\hat{\gamma}_3}{a^2M_*^2} F_2 + \frac{\alpha_2}{a^3M_*^3} F_3 + \frac{\hat{\gamma}_5}{a^4M_*^4} F_4 = \alpha^2 \sum_{n=0}^4 \delta_n \epsilon_*^n (-\eta)^n F_n, \tag{B1}$$

where the even parity terms are

$$F_0 = \widetilde{\mathfrak{G}}_k (k_1 s_1 + k_2 s_2 + k_3 s_3)^4, \qquad F_2 = \widetilde{\mathfrak{G}}_k (k_1 s_1 + k_2 s_2 + k_3 s_3)^4 \alpha^2 (k_1^2 + k_2^2 + k_3^2), \tag{B2}$$

$$F_{4} = \alpha^{4} \mathfrak{F}_{k} \{k_{1}^{8} + 4k_{1}^{7}s_{1}(k_{2}s_{2} + k_{3}s_{3}) + 6k_{1}^{6}(k_{2}s_{2} + k_{3}s_{3})^{2} + 4k_{1}^{5}s_{1}(k_{2}^{3}s_{2} + 4k_{2}k_{3}^{2}s_{2} + 4k_{2}^{2}k_{3}s_{3} + k_{3}^{4}s_{3}) \\ + k_{1}^{4}(2k_{2}^{4} + 19k_{2}^{2}k_{3}^{2} + 2k_{3}^{4} + 16k_{2}k_{3}s_{2}s_{3}(k_{2}^{2} + k_{3}^{2})) + 2k_{1}^{3}s_{1}(2k_{2}^{5}s_{2} + 8k_{2}^{4}k_{3}s_{3} + 9k_{2}^{3}k_{3}^{2}s_{2} + 9k_{2}^{2}k_{3}^{3}s_{3}) \\ + 8k_{2}k_{3}^{4}s_{2} + 2k_{3}^{5}s_{3}) + k_{1}^{2}[6k_{2}^{6} + 19k_{2}^{4}k_{3}^{2} + 19k_{2}^{2}k_{3}^{4} + 6k_{3}^{6} + 2k_{3}s_{3}s_{2}(8k_{2}^{5} + 9k_{2}^{3}k_{3}^{2} + 8k_{2}k_{3}^{4})] \\ + 4k_{1}s_{1}(k_{2}^{4} + k_{2}^{2}k_{3}^{2} + k_{3}^{4})(k_{2}s_{2} + k_{3}s_{3})^{3} + (k_{2}^{4} + k_{3}^{4})(k_{2}s_{2} + k_{3}s_{3})^{4}\},$$
(B3)

$$\mathfrak{F}_{k} = -\frac{(k_{1} - k_{2} - k_{3})(k_{1} + k_{2} - k_{3})(k_{1} - k_{2} + k_{3})(k_{1} + k_{2} + k_{3})}{256k_{1}^{2}k_{2}^{2}k_{3}^{2}},\tag{B4}$$

and the odd parity terms are

$$\begin{split} F_{1} &= -\alpha \widetilde{\mathfrak{G}}_{k} \{ k_{1}^{5}(4s_{1} - 4s_{2} - 5s_{3}) + k_{2}^{5}(4s_{2} - 4s_{3} - 5s_{1}) + k_{3}^{5}(4s_{3} - 4s_{1} - 5s_{2}) - 2k_{1}^{3}k_{2}^{2}(s_{1} - 5s_{2} - s_{3} + 2s_{1}s_{2}s_{3}) \\ &- 2k_{1}^{3}k_{3}^{2}(s_{1} - 4s_{3} + 2s_{1}s_{2}s_{3}) - 2k_{2}^{3}k_{3}^{2}(s_{2} - 5s_{3} - s_{1} + 2s_{1}s_{2}s_{3}) - 2k_{2}^{3}k_{1}^{2}(s_{2} - 4s_{1} + 2s_{1}s_{2}s_{3}) \\ &- 2k_{3}^{3}k_{1}^{2}(s_{3} - 5s_{1} - s_{2} + 2s_{1}s_{2}s_{3}) - 2k_{3}^{3}k_{2}^{2}(s_{3} - 4s_{2} + 2s_{1}s_{2}s_{3}) + k_{1}^{4}k_{2}(-3s_{1} + 6s_{2} - 4s_{3} - 4s_{1}s_{2}s_{3}) \\ &+ k_{1}^{4}k_{3}(-6s_{1} - 5s_{2} + 6s_{3} - 4s_{1}s_{2}s_{3}) + k_{2}^{4}k_{3}(-3s_{2} + 6s_{3} - 4s_{1} - 4s_{1}s_{2}s_{3}) + k_{2}^{4}k_{1}(-6s_{2} - 5s_{3} + 6s_{1} - 4s_{1}s_{2}s_{3}) \\ &+ k_{3}^{4}k_{1}(-3s_{3} + 6s_{1} - 4s_{2} - 4s_{1}s_{2}s_{3}) + k_{3}^{4}k_{2}(-6s_{3} - 5s_{1} + 6s_{2} - 4s_{1}s_{2}s_{3}) \\ &+ 2k_{1}^{2}k_{2}^{2}k_{3}(5s_{1} + 4s_{2} - 2s_{3} + 18s_{1}s_{2}s_{3}) + 2k_{1}^{3}k_{2}k_{3}(-7s_{1} + 7s_{2} + 7s_{3} + 2s_{1}s_{2}s_{3}) \\ &+ 2k_{2}^{2}k_{3}^{2}k_{1}(5s_{2} + 4s_{3} - 2s_{1} + 18s_{1}s_{2}s_{3}) + 2k_{3}^{3}k_{1}k_{2}(-7s_{3} + 7s_{1} + 7s_{2} + 2s_{1}s_{2}s_{3}) \}, \end{split}$$

$$\begin{split} F_{3} &= \frac{a^{3}\widetilde{k}_{k}}{2k_{1}^{2}k_{2}^{2}k_{3}^{2}} \{k_{1}^{11}k_{3}^{2}[2s_{2}] + k_{1}^{9}k_{2}k_{3}^{3}[-8s_{3}] + k_{1}^{9}k_{2}^{2}k_{3}^{2}[5s_{1} - 8s_{2} - 2s_{3}] + k_{1}^{9}k_{3}^{4}[-4s_{2}] + k_{2}^{11}k_{1}^{2}[2s_{3}] + k_{2}^{9}k_{3}k_{1}^{3}[-8s_{1}] \\ &+ k_{2}^{9}k_{3}^{2}k_{1}^{2}[5s_{2} - 8s_{3} - 2s_{1}] + k_{2}^{9}k_{1}^{4}[-4s_{3}] + k_{1}^{11}k_{2}^{2}[2s_{1}] + k_{3}^{9}k_{1}k_{2}^{3}[-8s_{2}] + k_{3}^{9}k_{1}^{2}k_{2}^{2}[5s_{3} - 8s_{1} - 2s_{2}] \\ &+ k_{3}^{9}k_{2}^{4}[-4s_{1}] + k_{1}^{8}k_{2}^{9}k_{3}^{3}[8s_{1}s_{3}(s_{1} - s_{3})] + k_{1}^{8}k_{2}^{3}k_{2}^{3}[9 + 8s_{1}(-s_{2} + s_{3})] + k_{1}^{7}k_{2}k_{3}^{5}[8s_{3}] + k_{1}^{7}k_{3}^{6}[2s_{2}] \\ &+ k_{3}^{8}k_{3}^{3}k_{1}^{2}[2s_{1} - s_{1}] + k_{3}^{8}k_{3}^{3}k_{1}^{2}[9 + 8s_{2}(-s_{3} + s_{1})] + k_{2}^{7}k_{3}k_{5}^{5}[8s_{1}] + k_{2}^{7}k_{1}^{6}[2s_{3}] + k_{3}^{8}k_{1}^{3}k_{2}^{2}[8s_{3}s_{2}(s_{3} - s_{2})] \\ &+ k_{3}^{8}k_{1}^{3}k_{2}^{2}[9 + 8s_{3}(-s_{1} + s_{2})] + k_{3}^{3}k_{1}^{5}[8s_{2}] + k_{7}^{7}k_{2}^{5}[2s_{1}] + k_{1}^{7}k_{2}^{4}k_{1}^{6}[2s_{3}] + k_{3}^{8}k_{1}^{3}k_{2}^{2}[8s_{3}s_{2}(s_{3} - s_{2})] \\ &+ k_{1}^{7}k_{2}^{4}k_{3}^{2}[2(-3s_{1} + s_{2} + 10s_{3})] + k_{1}^{7}k_{2}^{3}k_{3}^{3}[4s_{2}s_{3}(3s_{1} + 4s_{2} + 2s_{3})] + k_{1}^{7}k_{2}^{3}k_{3}^{4}[-5s_{2} + 20s_{3} - 4s_{1}] \\ &+ k_{1}^{7}k_{4}^{3}k_{1}^{2}[2(-3s_{3} + s_{1} + 10s_{1})] + k_{1}^{7}k_{3}^{3}k_{1}^{3}[4s_{1}s_{1}(3s_{2} + 4s_{3} + 2s_{1})] + k_{1}^{7}k_{2}^{3}k_{1}^{3}[1s_{1} - 2s_{2} - 23s_{3}] \\ &+ k_{1}^{6}k_{2}^{5}k_{3}^{2}[14s_{1} - 25s_{2} - 4s_{3} + 8s_{1}s_{2}s_{3}] + k_{2}^{6}k_{2}^{3}k_{1}^{2}[12s_{2} - 2s_{3} - 23s_{1}] + k_{1}^{6}k_{2}^{3}k_{1}^{4}[(2s_{2})(1 + 4s_{1}(s_{2} + s_{3}))] \\ &+ k_{1}^{6}k_{2}^{3}k_{1}^{3}[1s_{1}(-3 + 8s_{1}(2s_{2} + s_{3})] + k_{1}^{5}k_{2}^{3}k_{1}^{3}[-12s_{1} + 8s_{2} + 16s_{3} - 8s_{1}s_{2}s_{3}] + k_{1}^{6}k_{3}^{3}k_{1}^{4}[(2s_{1})(1 + 4s_{3}(s_{1} + s_{2}))] \\ &+ k_{1}^{6}k_{2}^{3}k_{1}^{3}[(s_{3})(2 - s_{1}s_{2} - s_{1}s_{3} - s_{2}s_{3}] + k_{1}^{6}k_{2}^{5}k_{1}$$

- L. M. Krauss, S. Dodelson, and S. Meyer, Science 328, 989 (2010); J. Garcia-Bellido, Prog. Theor. Phys. Suppl. 190, 322 (2011).
- U. Seljak and M. Zaldarriaga, Phys. Rev. Lett. 78, 2054 (1997); M. Kamionkowski, A. Kosowsky, and A. Stebbins, *ibid.* 78, 2058 (1997); J. Bock *et al.*, arXiv:astro-ph/0604101.
- [3] A. Lue, L. Wang, and M. Kamionkowski, Phys. Rev. Lett.
 83, 1506 (1999); N. Seto and A. Taruya, *ibid.* 99, 121101 (2007).
- [4] S. Saito, K. Ichiki, and A. Taruya, J. Cosmol. Astropart. Phys. 09 (2007) 002; C. Bischoff *et al.* Quiet Collaboration, Astrophys. J. 741, 111 (2011).
- [5] V. Gluscevic and M. Kamionkowski, Phys. Rev. D 81, 123529 (2010).
- [6] T. Takahashi and J. Soda, Phys. Rev. Lett. 102, 231301 (2009).
- [7] A. Wang, Q. Wu, W. Zhao, and T. Zhu, Phys. Rev. D 87, 103512 (2013).
- [8] P. Hořava, Phys. Rev. D 79, 084008 (2009).

- [9] J. M. Maldacena and G. L. Pimentel, J. High Energy Phys. 09 (2011) 045.
- [10] J. Soda, H. Kodama, and M. Nozawa, J. High Energy Phys. 08 (2011) 067.
- [11] M. Shiraishi, D. Nitta, and S. Yokoyama, Prog. Theor. Phys. **126**, 937 (2011).
- [12] T. Zhu, Q. Wu, A. Wang, and F.-W. Shu, Phys. Rev. D 84, 101502(R) (2011).
- [13] T. Zhu, F.-W. Shu, Q. Wu, and A. Wang, Phys. Rev. D 85, 044053 (2012).
- [14] A. Wang, Phys. Rev. D 82, 124063 (2010).
- [15] Y.-Q. Huang, A. Wang, R. Yousefi, and T. Zhu, Phys. Rev. D 88, 023523 (2013).
- [16] P. Ade *et al.* Planck Collaboration, arXiv:1303.5076.
- [17] K. Izumi, T. Kobayashi, and S. Mukohyama, J. Cosmol. Astropart. Phys. 10 (2010) 031.
- [18] Y.-Q. Huang and A. Wang, Phys. Rev. D 86, 103523 (2012).
- [19] K. S. Stelle, Phys. Rev. D 16, 953 (1977).
- [20] D. Blas, O. Pujolas, and S. Sibiryakov, J. High Energy Phys. 04 (2011) 018; P. Hořava, Classical Quantum Gravity 28, 114012 (2011); T. Clifton, P.G. Ferreira, A. Padilla, and C. Skordis, Phys. Rep. 513, 1 (2012).

- [21] S. Mukohyama, Classical Quantum Gravity 27, 223101 (2010).
- [22] P. Hořava and C. M. Melby-Thompson, Phys. Rev. D 82, 064027 (2010); A. Wang and Y. Wu, Phys. Rev. D 83, 044031 (2011); A. M. da Silva, Classical Quantum Gravity 28, 055011 (2011).
- [23] R. Arnowitt, S. Deser, and C. W. Misner, Gen. Relativ. Gravit. 40, 1997 (2008); C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman and Company, San Francisco, 1973), pp. 484–528.
- [24] M. Porrati, arXiv:hep-th/0409210.
- [25] J. Martin and R. Brandenberger, Phys. Rev. D 63, 123501 (2001).
- [26] J. Martin and R. Brandenberger, Phys. Rev. D 68, 063513 (2003); R. Brandenberger and J. Martin, Phys. Rev. D 71, 023504 (2005).
- [27] J. Maldacena, J. High Energy Phys. 05 (2003) 013;
 S. Weinberg, Phys. Rev. D 72, 043514 (2005).
- [28] J. Maldacena, J. High Energy Phys. 05 (2003) 013.
- [29] X. Gao, T. Kobayashi, M. Yamaguchi, and J. Yokoyama, Phys. Rev. Lett. **107**, 211301 (2011).