

A Note on Directed 5-Cycles in Digraphs*

Hao Liang^{1#}, Junming Xu²

¹Department of Mathematics, Southwestern University of Finance and Economics, Chengdu, China

²School of Mathematical Sciences, University of Science and Technology of China,

Wentsun Wu Key Laboratory of CAS, Hefei, China

Email: [#]lianghao@mail.ustc.edu.cn

Received May 18, 2012; revised June 18, 2012; accepted June 25, 2012

ABSTRACT

In this note, it is proved that if $\alpha \geq 0.24817$, then any digraph on n vertices with minimum outdegree at least αn contains a directed cycle of length at most 5.

Keywords: Digraph; Directed Cycle

1. Introduction

Let $G = (V, E)$ be a digraph without loops or parallel edges, where $V = V(G)$ is the vertex-set and $E = E(G)$ is the arc-set. In 1978, Caccetta and Häggkvist [1] made the following conjecture:

Conjecture 1.1 Any digraph on n vertices with minimum outdegree at least r contains a directed cycle of length at most $\lceil n/r \rceil$.

Trivially, this conjecture is true for $r = 1$, and it has been proved for $r = 2$ by Caccetta and Häggkvist [1], $r = 3$ by Hamildoune [2], $r = 4$ and $r = 5$ by Hoáng and Reed [3], $r < \sqrt{n/2}$ by Shen [4]. While the general conjecture is still open, some weaker statements have been obtained. A summary of results and problems related to the Caccetta-Häggkvist conjecture sees Sullivan [5].

For the conjecture, the case $r = n/2$ is trivial, the case $r = n/3$ has received much attention, but this special case is still open. To prove the conjecture, one may seek as small a constant α as possible such that any digraph on n vertices with minimum outdegree at least αn contains a directed triangle. The conjecture is that $\alpha = 1/3$. Caccetta and Häggkvist [1] obtained

$$\alpha \leq (3 - \sqrt{5})/2 \approx 0.3819, \text{ Bondy [6] showed}$$

$$\alpha \leq (2\sqrt{6} - 3)/5 \approx 0.3797, \text{ Shen [7] gave}$$

$\alpha \leq 3 - \sqrt{7} \approx 0.3542$, Hamburger, Haxell, and Kostochka [8] improved it to 0.35312. Hladký, Král' and Norin [9] further improved this bound to 0.3465. Namely, any digraph on n vertices with minimum out-degree at least $0.3465n$ contains a directed triangle. Very recently, Li-

chiardopol [10] showed that for $\beta \geq 0.343545$, any digraph on n vertices with both minimum out-degree and minimum in-degree at least βn contains a cycle of length at most 3.

In this note, we consider the minimum constant α such that any digraph on n vertices with minimum out-degree at least αn contains a directed cycle of length at most 5. The conjecture is that $\alpha = 1/5$. By refining the combinatorial techniques in [6,7,11], we prove the following result.

Theorem 1.2 If $\alpha \geq 0.24817$, then any digraph on n vertices with minimum outdegree at least αn contains a directed cycle of length at most 5.

2. Proof of Theorem 1.2

We prove Theorem 1.2 by induction on n . The theorem holds for $n \leq 5$ clearly. Now assume that the theorem holds for all digraphs with fewer than n vertices for $n \geq 5$. Let G be a digraph on n vertices with minimum outdegree at least αn . Suppose G contains no directed cycles with length at most 5. We can, without loss of generality, suppose that G is r -outregular, where $r = \lceil \alpha n \rceil$, that is, every vertex is of the outdegree r in G . We will try to deduce a contradiction. First we present some notations following [7].

For any $v \in V(G)$, let

$$N^+(v) = \{u \in V(G) : (v, u) \in E(G)\},$$

and $\deg^+(v) = |N^+(v)|$, the outdegree of v ;

$$N^-(v) = \{u \in V(G) : (u, v) \in E(G)\},$$

and $\deg^-(v) = |N^-(v)|$, the indegree of v .

We say $\langle u, v, w \rangle$ a *transitive triangle* if

*Supported by NNSF of China (No. 11071233).

#Corresponding author.

$(u, v), (v, w), (u, w) \in E(G)$. The arc (u, v) is called the base of the transitive triangle.

For any $(u, v) \in E(G)$, let

$$P(u, v) = N^+(v) \setminus N^+(u),$$

and $p(u, v) = |N^+(v) \setminus N^+(u)|$, the number of induced 2-path with the first arc (u, v) ;

$$Q(u, v) = N^-(u) \setminus N^-(v),$$

and $q(u, v) = |N^-(u) \setminus N^-(v)|$, the number of induced 2-path with the last arc (u, v) ;

$$T(u, v) = N^+(u) \cap N^+(v),$$

and $t(u, v) = |N^+(u) \cap N^+(v)|$, the number of transitive triangles with base (u, v) .

Lemma 2.1 For any $(u, v) \in E(G)$,

$$n > r + (1 - \alpha)r + (1 - \alpha)^2 r + (1 - \alpha)^3 t(u, v) + \text{deg}^-(v) + q(u, v). \tag{1}$$

Proof: To prove this inequality, we consider two cases according to $t(u, v) = 0$ or $t(u, v) > 0$.

If $t(u, v) = 0$, then substituting it into (1) yields

$$n > r + (1 - \alpha)r + (1 - \alpha)^2 r + \text{deg}^-(v) + q(u, v). \tag{2}$$

There exists some $w \in N^+(v)$ with outdegree less than αr in the subdigraph of G induced by $N^+(v)$ (Otherwise, this subdigraph would contain a directed 4-cycle by the induction hypothesis). Thus

$$|N^+(w) \setminus N^+(v)| \geq r - \alpha r.$$

Consider the subdigraph of G induced by $N^+(v) \cup N^+(w)$, by the induction hypothesis, some vertex $x \in N^+(v) \cup N^+(w)$ has outdegree less than $\alpha |N^+(v) \cup N^+(w)|$ in this subdigraph. Thus, the set of outneighbors of x not in $N^+(v) \cup N^+(w)$ satisfies

$$\begin{aligned} & |N^+(x) \setminus (N^+(v) \cup N^+(w))| \\ & \geq r - \alpha |N^+(v) \cup N^+(w)| \\ & = r - \alpha (|N^+(v)| + |N^+(w) \setminus N^+(v)|) \\ & = (1 - \alpha)r - \alpha |N^+(w) \setminus N^+(v)|, \end{aligned}$$

Since G has no directed 5-cycle, then $N^+(v)$, $N^+(w) \setminus N^+(v)$, $N^+(x) \setminus (N^+(v) \cup N^+(w))$, $N^-(v)$ and $N^-(u) \setminus N^-(v)$ are pairwise-disjoint sets with cardinalities r , $|N^+(w) \setminus N^+(v)|$,

$|N^+(x) \setminus (N^+(v) \cup N^+(w))|$, $\text{deg}^-(v)$ and $q(u, v)$, we have that

$$\begin{aligned} n & > r + |N^+(w) \setminus N^+(v)| + |N^+(x) \setminus (N^+(v) \cup N^+(w))| \\ & \quad + \text{deg}^-(v) + q(u, v) \\ & \geq r + (1 - \alpha)r + (1 - \alpha) |N^+(w) \setminus N^+(v)| \\ & \quad + \text{deg}^-(v) + q(u, v) \\ & \geq r + (1 - \alpha)r + (1 - \alpha)^2 r + \text{deg}^-(v) + q(u, v), \end{aligned}$$

Thus, the inequality (1) holds for $t(u, v) = 0$.

We now assume $t(u, v) > 0$. By the induction hypothesis, there is some vertex $w \in N^+(u) \cap N^+(v)$ that has outdegree less than $\alpha t(u, v)$ in the subdigraph of G induced by $N^+(u) \cap N^+(v)$, otherwise, this subdigraph would contain a directed 5-cycle. Also, w has not more than $p(u, v)$ outneighbors in the subdigraph of G induced by $N^+(v) \setminus N^+(u)$. Let $N^+(w) \setminus N^+(v)$ be the outneighbors of w which is not in $N^+(v)$. Noting that $t(u, v) = r - p(u, v)$, we have that

$$\begin{aligned} |N^+(w) \setminus N^+(v)| & \geq r - p(u, v) - \alpha t(u, v) \\ & = (1 - \alpha)t(u, v). \end{aligned} \tag{3}$$

Because G has no directed triangle, all outneighbors of w are neither in $N^+(v)$ nor in $N^-(u) \setminus N^-(v)$. Consider the subdigraph of G induced by $N^+(v) \cup N^+(w)$, by the induction hypothesis, there is some vertex $x \in N^+(v) \cup N^+(w)$ that has outdegree less than $\alpha |N^+(v) \cup N^+(w)|$ in this subdigraph. Thus, the set of outneighbors of x not in $N^+(v) \cup N^+(w)$ satisfies

$$\begin{aligned} & |N^+(x) \setminus (N^+(v) \cup N^+(w))| \\ & \geq r - \alpha |N^+(v) \cup N^+(w)| \\ & = r - \alpha (|N^+(v)| + |N^+(w) \setminus N^+(v)|) \\ & = (1 - \alpha)r - \alpha |N^+(w) \setminus N^+(v)|, \end{aligned} \tag{4}$$

Since G has no directed 4-cycle, all outneighbors of w are neither in $N^-(v)$ nor in $N^-(u) \setminus N^-(v)$. Consider the subdigraph of G induced by

$$N^+(v) \cup N^+(w) \cup N^+(x),$$

by the induction hypothesis, there is some vertex

$$y \in N^+(v) \cup N^+(w) \cup N^+(x)$$

that has outdegree less than

$$\alpha |N^+(v) \cup N^+(w) \cup N^+(x)|$$

in this subdigraph. Thus, the set of outneighbors of y not in $N^+(v) \cup N^+(w) \cup N^+(x)$ satisfies

$$\begin{aligned}
 & \left| N^+(y) \setminus (N^+(v) \cup N^+(w) \cup N^+(x)) \right| \\
 & \geq r - \alpha \left| N^+(v) \cup N^+(w) \cup N^+(x) \right| \\
 & = r - \alpha \left(\left| N^+(v) \cup N^+(w) \right| \right. \\
 & \quad \left. + \left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right| \right) \\
 & = (1 - \alpha)r - \alpha \left| N^+(w) \setminus N^+(v) \right| \\
 & \quad - \alpha \left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right|,
 \end{aligned} \tag{5}$$

Because G has no directed cycle of length at most 5, then $N^+(v)$, $N^+(w) \setminus N^+(v)$,

$$N^+(x) \setminus (N^+(v) \cup N^+(w)),$$

$$N^+(y) \setminus (N^+(v) \cup N^+(w) \cup N^+(x)),$$

$N^-(v)$ and $N^-(u) \setminus N^-(v)$ are pairwise-disjoint sets of cardinalities r , $|N^+(w) \setminus N^+(v)|$,

$$\left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right|,$$

$$\left| N^+(y) \setminus (N^+(v) \cup N^+(w) \cup N^+(x)) \right|,$$

$\deg^-(v)$ and $q(u, v)$, we have that

$$\begin{aligned}
 n & > r + \left| N^+(w) \setminus N^+(v) \right| \\
 & \quad + \left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right| \\
 & \quad + \left| N^+(y) \setminus (N^+(v) \cup N^+(w) \cup N^+(x)) \right| \\
 & \quad + \deg^-(v) + q(u, v)
 \end{aligned}$$

Substituting (3), (4) and (5) into this inequalities yields

$$\begin{aligned}
 n & > r + \left| N^+(w) \setminus N^+(v) \right| \\
 & \quad + \left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right| \\
 & \quad + \left| N^+(y) \setminus (N^+(v) \cup N^+(w) \cup N^+(x)) \right| \\
 & \quad + \deg^-(v) + q(u, v) \\
 & = r + |N^+(w) \setminus N^+(v)| + (1 - \alpha)r \\
 & \quad - \alpha \left| N^+(w) \setminus N^+(v) \right| \\
 & \quad + (1 - \alpha) \left| N^+(x) \setminus (N^+(v) \cup N^+(w)) \right| \\
 & \quad + \deg^-(v) + q(u, v) \\
 & \geq r + (1 - \alpha)r + (1 - \alpha)^2 r \\
 & \quad + (1 - \alpha)^2 \left| N^+(w) \setminus N^+(v) \right| + \deg^-(v) + q(u, v) \\
 & \geq r + (1 - \alpha)r + (1 - \alpha)^2 r \\
 & \quad + (1 - \alpha)^3 t(u, v) + \deg^-(v) + q(u, v)
 \end{aligned}$$

as desired, and so the lemma follows.

Connect to Proof of Theorem 1.2

Recalling that $t(u, v) = r - p(u, v)$, we can rewrite the inequality (1) as

$$\begin{aligned}
 & (3\alpha - 3\alpha^2 + \alpha^3)t(u, v) \\
 & > (4 - 3\alpha + \alpha^2)r - n + \deg^-(v) + q(u, v) - p(u, v).
 \end{aligned} \tag{6}$$

Summing over all $(u, v) \in E(G)$, we have that

$$\sum_{(u,v) \in E(G)} t(u, v) = t, \tag{7}$$

where t is the number of transitive triangles in G , and

$$\sum_{(u,v) \in E(G)} (4 - 3\alpha + \alpha^2)r - n = nr \left[(4 - 3\alpha + \alpha^2)r - n \right]. \tag{8}$$

By Cauchy's inequality and the first theorem on graph theory (see, for example, Theorem 1.1 in [12]), we have that

$$\begin{aligned}
 \sum_{(u,v) \in E(G)} \deg^-(v) & = \sum_{v \in V(G)} (\deg^-(v))^2 \\
 & \geq \frac{1}{n} \left(\sum_{v \in V(G)} \deg^-(v) \right)^2 = nr^2,
 \end{aligned}$$

that is

$$\sum_{(u,v) \in E(G)} \deg^-(v) \geq nr^2. \tag{9}$$

Because $\sum_{(u,v) \in E(G)} p(u, v)$ and $\sum_{(u,v) \in E(G)} q(u, v)$ are both equal to the number of induced directed 2-paths in G , it follows that

$$\sum_{(u,v) \in E(G)} p(u, v) = \sum_{(u,v) \in E(G)} q(u, v). \tag{10}$$

Summing over all $(u, v) \in E(G)$ for the inequality (6) and substituting inequalities (7)-(10) into that inequality yields,

$$(3\alpha - 3\alpha^2 + \alpha^3)t > (5 - 3\alpha + \alpha^2)nr^2 - n^2r. \tag{11}$$

Noting that $t \leq n \binom{r}{2}$ (see Shen [7]), we have that

$$\begin{aligned}
 t(3\alpha - 3\alpha^2 + \alpha^3) & \leq n \binom{r}{2} (3\alpha - 3\alpha^2 + \alpha^3) \\
 & < \frac{nr^2}{2} (3\alpha - 3\alpha^2 + \alpha^3).
 \end{aligned} \tag{12}$$

Combining (11) with (12) yields

$$(5 - 3\alpha + \alpha^2)nr^2 - n^2r < \frac{nr^2}{2} (3\alpha - 3\alpha^2 + \alpha^3). \tag{13}$$

Dividing both sides of the inequality (13) by $\frac{nr^2}{2}$,

and noting that $r = \lceil \alpha n \rceil \geq \alpha n$, we get

$$2(5 - 3\alpha + \alpha^2) - \frac{2}{\alpha} < (3\alpha - 3\alpha^2 + \alpha^3),$$

that is

$$\alpha^4 - 5\alpha^3 + 9\alpha^2 - 10\alpha + 2 > 0.$$

We obtain that $\alpha < 0.248164$, a contradiction. This completes the proof of the theorem.

REFERENCES

- [1] L. Caccetta and R. Häggkvist, "On Minimal Digraphs with Given Girth," *Proceedings of the 9th Southeast Conference on Combinatorics, Graph Theory, and Computing*, Boca Raton, 1978, pp. 181-187.
- [2] Y. O. Hamidoune, "A Note on Minimal Directed Graphs with Given Girth," *Journal of Combinatorial Theory, Series B*, Vol. 43, No. 3, 1987, pp. 343-348.
- [3] C. Hoàng and B. Reed, "A Note on Short Cycles in Digraphs," *Discrete Mathematics*, Vol. 66, No. 1-2, 1987, pp. 103-107. [doi:10.1016/0012-365X\(87\)90122-1](https://doi.org/10.1016/0012-365X(87)90122-1)
- [4] J. Shen, "On the Girth of Digraphs," *Discrete Mathematics*, Vol. 211, No. 1-3, 2000, pp. 167-181. [doi:10.1016/S0012-365X\(99\)00323-4](https://doi.org/10.1016/S0012-365X(99)00323-4)
- [5] B. D. Sullivan, "A Summary of Results and Problems Related to the Caccetta-Häggkvist Conjecture," 2006. <http://www.aimath.org/WWN/caccetta/caccetta.pdf>
- [6] J. A. Bondy, "Counting Subgraphs: A New Approach to the Caccetta-Häggkvist Conjecture," *Discrete Mathematics*, Vol. 165-166, 1997, pp. 71-80. [doi:10.1016/S0012-365X\(96\)00162-8](https://doi.org/10.1016/S0012-365X(96)00162-8)
- [7] J. Shen, "Directed Triangles in Digraphs," *Journal of Combinatorial Theory, Series B*, Vol. 74, 1998, pp. 405-407.
- [8] P. Hamburger, P. Haxell and A. Kostochka, "On the Directed Triangles in Digraphs," *Electronic Journal of Combinatorics*, Vol. 14, No. 19, 2007.
- [9] J. Hladký, D. Král' and S. Norin, "Counting Flags in Triangle-Free Digraphs," *Electronic Notes in Discrete Mathematics*, Vol. 34, 2009, pp. 621-625. [doi:10.1016/j.endm.2009.07.105](https://doi.org/10.1016/j.endm.2009.07.105)
- [10] N. Lichiardopol, "A New Bound for a Particular Case of the Caccetta-Häggkvist Conjecture," *Discrete Mathematics*, Vol. 310, No. 23, 2010, pp. 3368-3372. [doi:10.1016/j.disc.2010.07.026](https://doi.org/10.1016/j.disc.2010.07.026)
- [11] Q. Li and R. A. Brualdi, "On Minimal Regular Digraphs with Girth 4," *Czechoslovak Mathematical Journal*, Vol. 33, 1983, pp. 439-447.
- [12] J.-M. Xu, "Theory and Application of Graphs," Kluwer Academic Publishers, Dordrecht/Boston/London, 2003.